

LENGTH IN THE CREMONA GROUP

JÉRÉMY BLANC AND JEAN-PHILIPPE FURTER

ABSTRACT. The Cremona group is the group of birational transformations of the plane. By the famous Noether-Castelnuovo theorem, every element f of the group is a product of birational transformations, usually called Jonquières transformations, sending a pencil of lines onto another pencil of lines. The minimal number of such elements will be called the length, and written $\text{lgth}(f)$. Even if this length is rather unpredictable, we provide an explicit algorithm to compute it, which only depends on the multiplicities of the linear system of f .

As an application of this computation, we give a few properties of the dynamical length of f defined as the limit of the sequence $n \mapsto \text{lgth}(f^n)/n$. It follows for example that all distorted elements of the Cremona group are algebraic. The computation of the length may also be applied to the so called Wright complex associated with the Cremona group: This is made by Lonjou in [Lon2018]. Moreover, we show that the restriction of the length to the automorphism group of the affine plane is the classical length of this latter group (the length coming from its amalgamated structure). We also compute the lengths and dynamical lengths of all monoidal transformations, and of some Halphen transformations. Finally, we show that the length is a lower semicontinuous map on the Cremona group, endowed with its Zariski topology.

CONTENTS

1. Introduction	2
1.1. The length of elements of the Cremona group	2
1.2. Dynamical length	5
1.3. Distorted elements	5
1.4. Lower semicontinuity of the length	6
2. Reminders	7
2.1. The bubble space and the Picard-Manin space	7
2.2. The infinite Weyl group	10
2.3. Jonquières elements viewed in the Weyl group	14
3. The algorithm that computes the length and the proof of Theorem 1	17
3.1. Degree, maximal multiplicity and comultiplicity	17
3.2. Predecessors	19
3.3. From the Weyl group to the Cremona group	26
4. Examples and applications	31
4.1. Length of birational maps of small degree	31
4.2. Automorphisms of the affine plane	33

Date: March 16, 2018.

The first author acknowledges support by the Swiss National Science Foundation Grant “Birational Geometry” PP00P2_153026. The second author acknowledges support by the Région Poitou-Charentes



Both authors were also supported by the ANR Grant “BirPol” ANR-11-JS01-004-01.

4.3.	Decreasing the length and increasing the degree via a single Jonquières element	34
4.4.	The number of predecessors is not uniformly bounded	35
4.5.	Reduced decompositions of arbitrary lengths	36
4.6.	Examples of dynamical lengths	38
4.7.	Length of monomial transformations	40
4.8.	Dynamical length of regularisable elements and the proof of Theorem 2	46
5.	Lower semicontinuity of the length: the proof of Theorem 3	49
5.1.	Variables	49
5.2.	Definition of the Zariski topology on $\text{Bir}(\mathbb{P}^2)$ and basic properties	50
5.3.	The use of a valuative criterion	52
5.4.	The end of the proof of Theorem 3	55
	References	58

1. INTRODUCTION

1.1. The length of elements of the Cremona group. Let us fix an algebraically closed field k . The Cremona group over k , often written $\text{Cr}_2(k)$, is the group $\text{Bir}(\mathbb{P}^2)$ of k -birational transformations of the projective plane \mathbb{P}^2 . Such transformations can be written under the form

$$[x : y : z] \dashrightarrow [p_0(x, y, z) : p_1(x, y, z) : p_2(x, y, z)]$$

where $p_0, p_1, p_2 \in k[x, y, z]$ are homogeneous polynomials of the same degree, and this degree is the degree of the map, if the polynomials have no common factor. The Cremona transformations of degree 1 are the automorphisms of \mathbb{P}^2 , i.e. elements of the group $\text{Aut}(\mathbb{P}^2) = \text{PGL}_3(k)$, and Cremona transformations of degree 2 are called quadratic maps.

The group $\text{Bir}(\mathbb{P}^2)$ is generated by the automorphism group $\text{Aut}(\mathbb{P}^2)$ and by the single involution $\sigma : [x : y : z] \dashrightarrow [yz : xz : xy]$, called the *standard quadratic transformation*. The proof of this result, made by G. Castelnuovo [Cas1901], first decomposes an element of $\text{Bir}(\mathbb{P}^2)$ into a product of *Jonquières* elements (also called *Jonquières transformations*).

These latter maps are defined as the birational maps f for which there exist points $p, q \in \mathbb{P}^2$ such that f sends the pencil of lines passing through p to the pencil of lines passing through q . In this text, the group of Jonquières transformations preserving the pencil of lines passing through a given point $p \in \mathbb{P}^2$ is denoted by $\text{Jonq}_p \subseteq \text{Bir}(\mathbb{P}^2)$. The set of all Jonquières transformations is then equal to

$$\text{Jonq} = \bigcup_{p \in \mathbb{P}^2} \text{Aut}(\mathbb{P}^2) \text{Jonq}_p \text{Aut}(\mathbb{P}^2) = \bigcup_{p \in \mathbb{P}^2} \text{Aut}(\mathbb{P}^2) \text{Jonq}_p = \text{Aut}(\mathbb{P}^2) \text{Jonq}_{p_0} \text{Aut}(\mathbb{P}^2),$$

for any fixed point $p_0 \in \mathbb{P}^2$. The above equalities follow from the equality $\alpha \circ \text{Jonq}_p \circ \alpha^{-1} = \text{Jonq}_{\alpha(p)}$, which holds for each $\alpha \in \text{Aut}(\mathbb{P}^2)$.

Nowadays, one can also see the proof of Castelnuovo theorem by using the Sarkisov program (see [Cor1995]), and the number of Jonquières transformations needed corresponds to the number of links involved, which do not preserve a fibration. The proof

that every Jonquières transformation is a product of linear maps and σ is then an easy exercise (see for example [Alb2002, §8.4]).

In order to study the complexity of an element of $\text{Bir}(\mathbb{P}^2)$, according to the above decomposition, one naturally defines the following:

Definition 1.1. For each Cremona transformation $f \in \text{Bir}(\mathbb{P}^2)$ we define its *length* $\text{lgth}(f)$ in the following manner. If $f \in \text{Aut}(\mathbb{P}^2)$, we set $\text{lgth}(f) = 0$. Otherwise, we set $\text{lgth}(f) = n$ where n is the least positive integer for which f admits a decomposition

$$f = \varphi_n \circ \cdots \circ \varphi_1, \quad \forall i, \varphi_i \in \text{Jonq}.$$

In the above definition, note that we have $\text{lgth}(f^{-1}) = \text{lgth}(f)$.

For any fixed point $p_0 \in \mathbb{P}^2$ the group $\text{Bir}(\mathbb{P}^2)$ is generated by its two subgroups $\text{Aut}(\mathbb{P}^2)$ and Jonq_{p_0} . The length of f might also be seen as the least nonnegative integer n for which f admits a decomposition

$$f = \alpha_n \circ \varphi_n \circ \cdots \circ \alpha_1 \circ \varphi_1 \circ \alpha_0, \quad \forall i, \alpha_i \in \text{Aut}(\mathbb{P}^2) \text{ and } \forall j, \varphi_j \in \text{Jonq}_{p_0}.$$

In particular, this least integer n is independent from p_0 . All this follows from the equality $\text{Jonq} = \text{Aut}(\mathbb{P}^2)\text{Jonq}_{p_0}\text{Aut}(\mathbb{P}^2)$.

This notion of length is similar to the case of automorphisms of the affine plane. Taking a linear embedding $\mathbb{A}^2 \hookrightarrow \mathbb{P}^2$, the classical Jung-Van der Kulk theorem says that $\text{Aut}(\mathbb{A}^2)$ is generated by $\text{Aff}_2 := \text{Aut}(\mathbb{P}^2) \cap \text{Aut}(\mathbb{A}^2)$ and $\text{Jonq}_{p, \mathbb{A}^2} := \text{Jonq}_p \cap \text{Aut}(\mathbb{A}^2)$ for any $p \in \mathbb{P}^2 \setminus \mathbb{A}^2$ [Jun1942, vdK53, Lam2002]. Moreover, there is no relation except the trivial ones, i.e. the group $\text{Aut}(\mathbb{A}^2)$ is the amalgamated product of Aff_2 and $\text{Jonq}_{p, \mathbb{A}^2}$ over their intersection.

The length in $\text{Aut}(\mathbb{A}^2)$ is then easy to compute, by writing an element in a reduced form (i.e. in a product of elements of Aff_2 and $\text{Jonq}_{p, \mathbb{A}^2}$ where two consecutive elements do not belong to the same group). It has moreover natural properties, namely it is lower semicontinuous for the Zariski topology on $\text{Aut}(\mathbb{A}^2)$, as shown by [Fur2002] when $\text{char}(\mathbb{k}) = 0$ (in fact this result also holds in positive characteristic by Theorem 3 and Proposition 4.2 below).

The case of $\text{Bir}(\mathbb{P}^2)$ is more complicated, as $\text{Bir}(\mathbb{P}^2)$ is not the amalgamated product of $\text{Aut}(\mathbb{P}^2)$ and Jonq_{p_0} . There is only one relation, of very small length [Bla2012], which makes the group $\text{Bir}(\mathbb{P}^2)$ more complicated than the group $\text{Aut}(\mathbb{A}^2)$ (see also [Giz1982, Isk1985] for other presentations with generators and relations of $\text{Bir}(\mathbb{P}^2)$). In particular, there exist elements of $\text{Bir}(\mathbb{P}^2)$ of finite order (finitely many families up to conjugacy) which are neither conjugate to an element of $\text{Aut}(\mathbb{P}^2)$ nor to an element of Jonq_p [Bla2011], contrary to the case of amalgamated products. Another way to see the difference is that $\text{Aut}(\mathbb{A}^2)$ acts on a tree thanks to its amalgamated structure [Ser1980, Lam2001], but $\text{Bir}(\mathbb{P}^2)$ only acts on a simply connected simplicial complex of dimension two [Wri1992]. The group $\text{Bir}(\mathbb{P}^2)$ does not act (non-trivially) on a tree because it is not a non-trivial amalgamated product [Cor2013].

Computing the length of an element $f \in \text{Bir}(\mathbb{P}^2)$ is then more tricky than the case of $\text{Aut}(\mathbb{A}^2)$ and we cannot only take a reduced decomposition (i.e. a product $f = \varphi_n \circ \cdots \circ \varphi_1$ of Jonquières elements such that $\varphi_{i+1} \circ \varphi_i$ is not Jonquières for $i = 1, \dots, n-1$). The length of such reduced decompositions is unbounded (Proposition 4.10). One way to give an upper bound for the length of an element is to follow the proof of Castelnuovo and to apply successive Jonquières elements to decrease the degree (this is detailed in

Algorithm 3.25). It is a priori not clear that it gives the length. However, this is one of our main results and will be proven in Corollary 3.29.

Multiplying an element $f \in \text{Bir}(\mathbb{P}^2)$ with a Jonquières element φ , we have $\text{lgth}(f \circ \varphi) \in \{\text{lgth}(f) - 1, \text{lgth}(f), \text{lgth}(f) + 1\}$. The possibilities occur in a rather chaotic way since there are examples where $\deg(f \circ \varphi) > \deg(f)$ but $\text{lgth}(f \circ \varphi) = \text{lgth}(f) - 1$ (Proposition 4.3(3)). Moreover, the number of Jonquières elements φ such that $\text{lgth}(f \circ \varphi) = \text{lgth}(f) - 1$ can be infinite, even up to right multiplication with an element of $\text{Aut}(\mathbb{P}^2)$ (Proposition 4.3(4)).

We will however show that there is a natural algorithm that yields the length. Moreover, we will show that the length only depends on combinatorial properties of the maps, namely the multiplicities of the base-points. Recall that if $f \in \text{Bir}(\mathbb{P}^2)$ is of degree $d \geq 1$, then its linear system consists of curves of degree d passing through finitely many points p_1, \dots, p_r (lying on \mathbb{P}^2 or infinitely near) with some multiplicities m_1, \dots, m_r such that $\sum m_i = 3(d - 1)$, $\sum (m_i)^2 = d^2 - 1$ (see below, in particular Remark 2.11 and Lemmas 2.14 and 2.20). In particular, maps of degree 1 have no base-points and maps of degree 2 have three base-points of multiplicity 1. We say that (m_1, \dots, m_r) is the *homaloidal type* of f ; it is a finite sequence up to permutation, or equivalently a set with multiple entries. We often write $(d; m_1, \dots, m_r)$ to see the degree, but this one is of course uniquely determined by the multiplicities. We also define the *comultiplicity* of f to be $\deg(f) - \max_i m_i$. This notion is sometimes used in the literature, for instance in the proof of the Noether-Castelnuovo theorem given by Alexander [Ale1916]. One can observe that $\text{comult}(f) = 1$ if and only if f is a Jonquières element (follows from Lemma 2.42 and Definition 2.40). To state our main result, we use the following notion:

Definition 1.2. Let $f \in \text{Bir}(\mathbb{P}^2)$. A *predecessor* of f is an element of minimal degree among the elements of the form $f \circ \varphi$ where φ is a Jonquières transformation.

A precise description of the predecessors of an element of $\text{Bir}(\mathbb{P}^2)$ is algorithmic and not very difficult to obtain. In particular, the following holds:

Lemma 1.3. Let $f \in \text{Bir}(\mathbb{P}^2)$.

- (1) The homaloidal type of a predecessor of f is uniquely determined by the homaloidal type of f .
- (2) There are infinitely many predecessors of f , but only finitely many classes up to right composition with an element of $\text{Aut}(\mathbb{P}^2)$.
- (3) If φ is a Jonquières transformation such that $f \circ \varphi$ is a predecessor of f , then $\text{Base}(\varphi^{-1}) \subseteq \text{Base}(f)$.

Remark 1.4. Lemma 1.3(2) asserts that any Cremona transformation $f \in \text{Bir}(\mathbb{P}^2)$ admits finitely many predecessors up to right multiplication by an element of $\text{Aut}(\mathbb{P}^2)$. However, we will see in Lemma 4.6 that this number is not uniformly bounded on $\text{Bir}(\mathbb{P}^2)$.

Computing a sequence of predecessors (which is algorithmic, as said before, and whose homaloidal types are uniquely determined by the one of the map we start with) yields then a finite algorithm to compute the length of any element of $\text{Bir}(\mathbb{P}^2)$, as our main theorem states:

Theorem 1. Let $f_0 \in \text{Bir}(\mathbb{P}^2)$, let $n \geq 1$ be an integer, and let $(f_i)_{i \in \mathbb{N}}$ be a sequence of elements of $\text{Bir}(\mathbb{P}^2)$ such that f_i is a predecessor of f_{i-1} for each $i \geq 1$. For all Jonquières elements $\varphi_1, \dots, \varphi_n \in \text{Bir}(\mathbb{P}^2)$, the element $g_n = f \circ \varphi_1 \circ \dots \circ \varphi_n$ satisfies

- (1) $\deg(f_n) \leq \deg(g_n)$;
- (2) $\text{comult}(f_n) \leq \text{comult}(g_n)$;
- (3) If $\deg(f_n) = \deg(g_n)$, then f_n and g_n have the same homaloidal type.

In particular, $\text{lgth}(f) = \min\{n \mid \deg(f_n) = 1\}$.

This allows to easily compute the length of all maps of some given degree (see §4.1 for tables).

Another consequence of Theorem 1 is that the length of an element of $\text{Aut}(\mathbb{A}^2)$, viewed as an element of $\text{Bir}(\mathbb{P}^2)$, is the same as the classical length given by the amalgamated product (Proposition 4.2).

Also, the distance in the connected simplicial complex (or simply the graph associated) of [Wri1992] is given by the length in $\text{Bir}(\mathbb{P}^2)$. Hence, Theorem 1 provides a way to compute the distance in this graph. In particular, the graph of Wright is of unbounded length (follows for instance from Lemma 4.11), a fact that does not follow directly from the definition of the graph. Another application of Theorem 1, made by Lonjou is that the graph of Wright is not hyperbolic, in the sense of Gromov [Lon2018].

In §4.7, we compute the length of all monomial elements of $\text{Bir}(\mathbb{P}^2)$, and relate it to the decomposition of elements in $\text{SL}_2(\mathbb{Z})$ and continued fractions.

1.2. Dynamical length. Since $\text{lgth}(f \circ g) \leq \text{lgth}(f) + \text{lgth}(g)$ for all $f, g \in \text{Bir}(\mathbb{P}^2)$, the sequence $n \mapsto \text{lgth}(f^n)$ is subadditive so that the sequence $n \mapsto \frac{\text{lgth}(f^n)}{n}$ admits a limit when n goes to infinity. This allows the following definition:

Definition 1.5. For each $f \in \text{Bir}(\mathbb{P}^2)$, the *dynamical length* is defined as

$$\mathfrak{d}_{\text{lgth}}(f) := \lim_{n \rightarrow \infty} \frac{\text{lgth}(f^n)}{n} \in \mathbb{R}_+.$$

Note that $\mathfrak{d}_{\text{lgth}}(f)$ is invariant under conjugation (contrary to the length), and satisfies $0 \leq \mathfrak{d}_{\text{lgth}}(f) \leq \text{lgth}(f)$. It is not very easy to compute $\mathfrak{d}_{\text{lgth}}(f)$ in general, but we will do it precisely for all monomial elements of $\text{Bir}(\mathbb{P}^2)$, and relate this to continued fractions and decompositions in $\text{SL}_2(\mathbb{Z})$ (Section 4.7). We will also show that

$$\frac{1}{2}\mathbb{Z}_{\geq 0} \cup \frac{1}{3}\mathbb{Z}_{\geq 0} \subseteq \mathfrak{d}_{\text{lgth}}(\text{Bir}(\mathbb{P}^2)) = \{\mathfrak{d}_{\text{lgth}}(f) \mid f \in \text{Bir}(\mathbb{P}^2)\}$$

(Corollary 4.13), but do not have any example of a Cremona transformation $f \in \text{Bir}(\mathbb{P}^2)$ such that $\mathfrak{d}_{\text{lgth}}(f) \notin \frac{1}{2}\mathbb{Z}_{\geq 0} \cup \frac{1}{3}\mathbb{Z}_{\geq 0}$. In particular, every monomial map of $\text{Bir}(\mathbb{P}^2)$ has a dynamical length in $\frac{1}{2}\mathbb{Z}_{\geq 0}$ (follows from Proposition 4.25).

Question 1.6. What does the set $\mathfrak{d}_{\text{lgth}}(\text{Bir}(\mathbb{P}^2)) = \{\mathfrak{d}_{\text{lgth}}(f) \mid f \in \text{Bir}(\mathbb{P}^2)\}$ look like?

1.3. Distorted elements. We begin with the two following definitions.

Definition 1.7. If G is a group generated by a finite set $F \subseteq G$, an element $g \in G$ is said to be *distorted* if $\lim_{n \rightarrow \infty} \frac{|g^n|}{n} = 0$, where $|g^n|$ denotes the length of g^n according to F , i.e. the minimal number of elements of F necessary to write g^n (word length). This notion actually does not depend on the chosen F , but only on the pair (g, G) .

If G is any group, an element $g \in G$ is said to be *distorted* if it is distorted in some finitely generated subgroup of G .

Definition 1.8. An element $f \in \text{Bir}(\mathbb{P}^2)$ is said to be *algebraic* (or *elliptic*) if it is contained in an algebraic subgroup of $\text{Bir}(\mathbb{P}^2)$, or equivalently if the sequence $n \mapsto \deg(f^n)$ is bounded ([BlaFur2013, §2.6]). By [BlaDés2015, Proposition 2.3], this is also equivalent to saying that f is of finite order or conjugate to an element of $\text{Aut}(\mathbb{P}^2)$.

It is proven in [CC2018] that any algebraic element of $\text{Bir}(\mathbb{P}^2)$ is distorted. Using the dynamical length, we will prove the converse statement:

Theorem 2. *Any distorted element of $\text{Bir}(\mathbb{P}^2)$ is algebraic.*

A function $S: \text{Bir}(\mathbb{P}^2) \rightarrow \mathbb{R}_{\geq 0}$ is said to be *subadditive* if it satisfies $S(f \circ g) \leq S(f) + S(g)$ for all $f, g \in \text{Bir}(\mathbb{P}^2)$. For such a function, the limit $\lim_{n \rightarrow \infty} \frac{S(f^n)}{n} \in \mathbb{R}_{\geq 0}$ exists for any f , and is moreover equal to zero when f is distorted.

It turns out that the three following functions are subadditive on $\text{Bir}(\mathbb{P}^2)$: the length, the number of base-points, and the logarithm of the degree. For any such S , the corresponding limit $\lim_{n \rightarrow \infty} \frac{S(f^n)}{n}$ is: the dynamical length $\mathfrak{d}_{\text{lgth}}(f)$, the dynamical number of base-points (written $\mu(f)$ in [BlaDés2015]), and the logarithm of the dynamical degree $\log(\lambda(f))$, where the dynamical degree is $\lambda(f) = \lim_{n \rightarrow \infty} (\deg(f^n))^{1/n}$.

We will show that f is algebraic if and only if $\mathfrak{d}_{\text{lgth}}(f) = \mu(f) = \log(\lambda(f)) = 0$, thus proving Theorem 2. More precisely, we can decompose elements of $\text{Bir}(\mathbb{P}^2)$ into five disjoint subsets of elements (see §4.8), and the situation is as in Figure 1: In particular,

f	$\mathfrak{d}_{\text{lgth}}(f)$	$\mu(f)$	$\log(\lambda(f))$
Algebraic elements	0	0	0
Jonquières twists	0	> 0	0
Halphen twists	> 0 (Corollary 4.32)	0	0
Regularisable loxodromic elements	> 0 (Proposition 4.34)	0	> 0
Non-regularisable loxodromic elements	sometimes > 0 (Lemma 4.11)	> 0	> 0

FIGURE 1. Positivity of $\mathfrak{d}_{\text{lgth}}(f)$, $\mu(f)$, $\log(\lambda(f))$ for elements $f \in \text{Bir}(\mathbb{P}^2)$

Corollary 4.32 is sufficient for showing that an element f of $\text{Bir}(\mathbb{P}^2)$ is algebraic if and only if $\mathfrak{d}_{\text{lgth}}(f) = \mu(f) = \log(\lambda(f)) = 0$.

1.4. Lower semicontinuity of the length. Even if $\text{Bir}(\mathbb{P}^2)$ is not naturally an ind-group ([BlaFur2013, Theorem 1]), following [Dem1970, Ser2010], it admits a natural Zariski topology (see Definition 5.8). We prove that the length is compatible with this topology, and thus behaves well in families:

Theorem 3. *The length map $\text{lgth}: \text{Bir}(\mathbb{P}^2) \rightarrow \mathbb{N}$, $f \mapsto \text{lgth}(f)$ is lower semicontinuous. In other words, for each integer $\ell \geq 0$, the set $\{f \in \text{Bir}(\mathbb{P}^2) \mid \text{lgth}(f) \leq \ell\}$ is closed.*

As explained before, this implies the same result for automorphisms of \mathbb{A}^2 , already proven in [Fur2002] when $\text{char}(\mathbb{k}) = 0$. It also shows that some degenerations of birational maps are not possible. For instance, it is not possible to have a family of birational maps with homaloidal type $(7; 3^4; 2^3)$ which degenerates to a birational map of homaloidal type $(7; 5; 2^5, 1^3)$ (see [BCM2015, BlaCal2016] for more details on this question).

2. REMINDERS

When we want to decompose a birational transformation of \mathbb{P}^2 , we have to study the multiplicities of the linear system at points, and also the position of the points (if one is infinitely near to another, if they are on the same line, \dots). A very fruitful approach consists of looking at the (linear and faithful) action of $\text{Bir}(\mathbb{P}^2)$ on the so called Picard-Manin space. Forgetting the position of the points and studying only the arithmetic part can be done by studying an *infinite Weyl group* W_∞ , as done in [BlaCan2016]. This group still acts on the Picard-Manin space and contains $\text{Bir}(\mathbb{P}^2)$. An analogous Weyl group is used in [BlaCal2016], but with a slightly different definition.

2.1. The bubble space and the Picard-Manin space. Let us recall the following classical notions.

Definition 2.1. Let Y be a smooth projective rational surface. We denote by $\mathcal{B}(Y)$ the *bubble space* of Y . It is the set of points that belong, as proper or infinitely near points to Y . More precisely, an element of $\mathcal{B}(Y)$ is the equivalence class of a triple (p, X, π) , where X is a smooth projective surface, $\pi: X \rightarrow Y$ is a birational morphism (a sequence of blow-ups) and $p \in X$. Two triples (p, X, π) and (p', X', π') are equivalent if $(\pi')^{-1} \circ \pi: X \dashrightarrow X'$ restricts to an isomorphism $U \rightarrow U'$, where $U \subseteq X$, $U' \subseteq X'$ are two open neighbourhoods of p and p' , and if p is sent to p' by this isomorphism.

Definition 2.2. There is a natural order on $\mathcal{B}(Y)$. We say that $(p, X, \pi) \geq (p', X', \pi')$ if $(\pi')^{-1} \circ \pi: X \dashrightarrow X'$ restricts to a morphism $U \rightarrow X'$, where $U \subseteq X$ is an open subset containing p and if p is sent on p' by this morphism.

Remark 2.3. We have an inclusion $\mathbb{P}^2 \hookrightarrow \mathcal{B}(\mathbb{P}^2)$, that sends a point $p \in \mathbb{P}^2$ onto the equivalence class of $(p, \mathbb{P}^2, \text{id})$. We will also see elements of $\mathcal{B}(\mathbb{P}^2)$ as points, the surfaces and the morphisms being then implicit.

Every birational map $\varphi: \mathbb{P}^2 \dashrightarrow \mathbb{P}^2$ has a finite number of base-points. The set of all such points is denoted $\text{Base}(\varphi) \subseteq \mathcal{B}(\mathbb{P}^2)$. Moreover, φ induces a bijection $\mathcal{B}(\mathbb{P}^2) \setminus \text{Base}(\varphi) \rightarrow \mathcal{B}(\mathbb{P}^2) \setminus \text{Base}(\varphi^{-1})$.

Let us recall the following classical notions. See for example [Alb2002] and references there.

Definition 2.4. Let $p, q \in \mathcal{B}(\mathbb{P}^2)$. We say that p is *infinitely near* q if $p \geq q$ (for the order defined above). We say that p is in the *first neighbourhood* of q if $p > q$ and if there is no $r \in \mathcal{B}(\mathbb{P}^2)$ with $p > r > q$. We say that a point $p \in \mathcal{B}(\mathbb{P}^2)$ is a *proper point* of \mathbb{P}^2 if p is minimal. This corresponds to saying that $p \in \mathbb{P}^2 \subseteq \mathcal{B}(\mathbb{P}^2)$.

Definition 2.5. Let Y be a smooth projective rational surface. Its Picard-Manin space \mathcal{Z}_Y is defined as the inductive limit of all the Picard groups $\text{Pic}(X)$, where X is a smooth projective rational surface and $X \rightarrow Y$ is a birational morphism.

More precisely, an element $c \in \mathcal{Z}_Y$ corresponds to an equivalence class of triples (C, X, π) , where X is a smooth projective rational surface, $\pi: X \rightarrow Y$ is a birational morphism and $C \in \text{Pic}(X)$. Two triples (C_1, X_1, π_1) and (C_2, X_2, π_2) are identified if one can find another smooth projective rational X_3 together with birational morphisms $\pi'_1: X_3 \rightarrow X_1$, $\pi'_2: X_3 \rightarrow X_2$ such that $\pi_1 \circ \pi'_1 = \pi_2 \circ \pi'_2$, and such that $(\pi'_1)^*(C_1) = (\pi'_2)^*(C_2)$.

The \mathbb{Z} -module \mathcal{Z}_Y is endowed with an intersection form and canonical form $\omega: \mathcal{Z}_Y \rightarrow \mathbb{Z}$. The canonical form sends a triple (C, X, π) onto $K_X \cdot C \in \mathbb{Z}$, where K_X is the canonical class of X . To intersect two classes, we take representants (C_1, X, π) and (C_2, X, π) on the same surface by taking a common resolution and compute $C_1 \cdot C_2 \in \mathbb{Z}$.

Remark 2.6. If (C, X, π) is a triple as in the above definition and $\epsilon: X' \rightarrow X$ is a birational morphism, then $(\epsilon^*(C), X', \pi \circ \epsilon)$ is equivalent to (C, X, π) . Moreover, $K_X \cdot C = \epsilon^*(K_{X'}) \cdot \epsilon^*(C) = K_{X'} \cdot \epsilon^*(C)$.

Using this remark we obtain that the canonical form $\omega: \mathcal{Z}_Y \rightarrow \mathbb{Z}$ defined above is independent of the choice of the triple in the equivalence class of an element of \mathcal{Z}_Y . The intersection form is also well-defined, as $\epsilon^*(C) \cdot \epsilon^*(D) = C \cdot D$ for all $C, D \in \text{Pic}(X)$.

Definition 2.7. Let Y be a smooth projective rational surface. For each point $q \in \mathcal{B}(Y)$, we define an element $e_q \in \mathcal{Z}_Y$ as follows: the point q is the class of (p, X, π) , and $e_q \in \mathcal{Z}_Y$ is the class of $(E_p, \hat{X}, \pi \circ \epsilon)$, where $\epsilon: \hat{X} \rightarrow X$ is the blow-up of $p \in X$, and $E_p = \epsilon^{-1}(p) \in \text{Pic}(X)$ is the exceptional divisor.

Lemma 2.8. *Let Y be a smooth projective rational surface. The group \mathcal{Z}_Y is naturally isomorphic to*

$$\mathcal{Z}_Y \simeq \text{Pic}(Y) \oplus \bigoplus_{p \in \mathcal{B}(Y)} \mathbb{Z}e_p.$$

Moreover, the restriction of the intersection form of \mathcal{Z}_Y on $\text{Pic}(Y)$ is the classical one, and we have

$$C \cdot e_p = 0, e_p^2 = -1, e_p \cdot e_q = 0, \omega(C) = C \cdot K_Y, \omega(e_p) = -1.$$

for all $p, q \in \mathcal{B}(Y)$, $C \in \text{Pic}(Y)$, $p \neq q$.

Proof. The map sending $C \in \text{Pic}(Y)$ onto the class of (C, Y, id) yields an inclusion $\text{Pic}(Y) \hookrightarrow \mathcal{Z}_Y$. By definition, the restriction of the intersection form and the canonical form of \mathcal{Z}_Y on $\text{Pic}(Y)$ are the classical ones.

If $\epsilon: \hat{X} \rightarrow X$ is the blow-up of a point $p \in X$, then $\text{Pic}(\hat{X}) = \epsilon^* \text{Pic}(X) \oplus \mathbb{Z}e_p$, where the exceptional divisor $e_p \in \hat{X}$ satisfies $e_p^2 = -1$, $e_p \cdot R = 0$ for each $R \in \epsilon^*(\text{Pic}(X))$. Moreover, $K_{\hat{X}} = \pi^*(K_X) + e_p$. This provides the result, as every birational morphism $X \rightarrow Y$, where X is a smooth projective rational surface, is a sequence of blow-ups of finitely many points of $\mathcal{B}(Y)$. \square

Corollary 2.9. *The group $\mathcal{Z}_{\mathbb{P}^2}$ is naturally isomorphic to*

$$\mathbb{Z}e_0 \oplus \bigoplus_{p \in \mathcal{B}(\mathbb{P}^2)} \mathbb{Z}e_p,$$

where $e_0 \in \text{Pic}(\mathbb{P}^2)$ is the class of a line and e_p corresponds to the exceptional divisor of $p \in \mathcal{B}(\mathbb{P}^2)$. Moreover, we have

$$(e_0)^2 = 1, e_p^2 = -1, \omega(e_p) = -1, \omega(e_0) = -3 \text{ and } e_p \cdot e_q = 0$$

for all $p, q \in \mathcal{B}(\mathbb{P}^2)$, $p \neq q$.

Proof. Follows from Lemma 2.8 and the fact that $\text{Pic}(\mathbb{P}^2) = \mathbb{Z}e_0$, $(e_0)^2 = 1$, $K_{\mathbb{P}^2} = -3e_0$, so $\omega(e_0) = -3$. \square

Definition 2.10. Let $a \in \mathcal{Z}_{\mathbb{P}^2}$ and $q \in \mathcal{B}(\mathbb{P}^2)$. We define the *degree* of a to be $\deg(a) = e_0 \cdot a \in \mathbb{Z}$ and the *multiplicity of a at q* to be $m_q(a) = e_q \cdot a \in \mathbb{Z}$. We then define the set of *base-points* $\text{Base}(a)$ of a to be $\{q \in \mathcal{B}(\mathbb{P}^2) \mid m_q(a) \neq 0\}$.

Remark 2.11. Let Λ be a linear system on \mathbb{P}^2 which we assume of positive dimension and without fixed component. Denote by p_1, \dots, p_r its base-points and by $\pi: X \rightarrow \mathbb{P}^2$ the blow-up of the points p_i on \mathbb{P}^2 . Then, the strict transform $\tilde{\Lambda}$ of Λ on X is a base-point free linear system and we have

$$(\diamond) \quad \tilde{\Lambda} = d\pi^*(e_0) - \sum_{i=1}^r m_i e_{p_i} \in \text{Pic}(X),$$

where d is the degree of Λ , $m_1, \dots, m_r \geq 1$ are its multiplicities at the points p_1, \dots, p_r and $e_{p_i} \in \text{Pic}(X)$ is the pull-back (or total transform) of the exceptional curve produced by blowing-up p_i .

Indeed, since we have $\text{Pic}(X) = \pi^*(\text{Pic}(\mathbb{P}^2)) \oplus (\bigoplus_{i=1}^r \mathbb{Z}e_{p_i})$, there exist integers d, m_1, \dots, m_r for which the equality (\diamond) holds. We then compute

$$d = \pi^*(e_0) \cdot \tilde{\Lambda} = \pi^*(e_0) \cdot \pi^*(\Lambda) = e_0 \cdot \Lambda = \deg(\Lambda)$$

and we see that the multiplicity of Λ at p_i is $e_{p_i} \cdot \tilde{\Lambda} = m_i$. Hence, the definition of base-points, degree and multiplicities coincides with the classical one.

Definition 2.12. Let $\varphi: Y_1 \dashrightarrow Y_2$ be a birational map between two smooth projective rational surfaces. We define an isomorphism $\varphi_\bullet: \mathcal{Z}_{Y_1} \rightarrow \mathcal{Z}_{Y_2}$ in the following way:

An element $c \in \mathcal{Z}_{Y_1}$ corresponds to the class of a triple (C, X, π_1) . By blowing-up more points if necessary, we may assume that π_1 is such that $\pi_2 := \varphi \circ \pi_1: X \rightarrow Y_2$ is a birational morphism. We then define $\varphi_\bullet(c) \in \mathcal{Z}_{Y_2}$ to be the class of (C, X, π_2) .

Remark 2.13. If $\varphi: Y_1 \dashrightarrow Y_2$ and $\psi: Y_2 \dashrightarrow Y_3$ are two birational maps between smooth projective rational surfaces, then $(\psi \circ \varphi)_\bullet = \psi_\bullet \circ \varphi_\bullet$. This implies that φ and ψ are isomorphisms of \mathbb{Z} -modules. They moreover preserve the intersection form and the canonical form (which can be checked on blowing-ups).

We then obtain the following result:

Lemma 2.14. *The group $\text{Bir}(\mathbb{P}^2)$ acts faithfully on $\mathcal{Z}_{\mathbb{P}^2}$ and preserves the intersection form and the canonical form. Moreover, if $f \in \text{Bir}(\mathbb{P}^2)$, then $(f_\bullet)^{-1}(e_0) = de_0 - \sum_{i=1}^r m_i e_{p_i}$, where $d = \deg f$, $p_1, \dots, p_r \in \mathcal{B}(\mathbb{P}^2)$ are the base-points of f and $m_1, \dots, m_r \geq 1$ are their multiplicities.*

Proof. We decompose every $f \in \text{Bir}(\mathbb{P}^2)$ into $f = \eta \circ \pi^{-1}$, where $\eta: X \rightarrow \mathbb{P}^2$, $\pi: X \rightarrow \mathbb{P}^2$ are blow-ups of the base-points of f^{-1} and f respectively.

$$\begin{array}{ccc} & X & \\ \pi \swarrow & & \searrow \eta \\ \mathbb{P}^2 & \overset{f}{\dashrightarrow} & \mathbb{P}^2 \end{array}$$

We have $\text{Pic}(X) = \pi^*(\text{Pic}(\mathbb{P}^2)) \oplus (\bigoplus_{i=1}^r \mathbb{Z}e_{p_i})$, where e_{p_1}, \dots, e_{p_r} are the pull-backs in $\text{Pic}(X)$ of the exceptional divisors of the base-points p_1, \dots, p_r of f (or equivalently of π) and can thus write $\eta^*(e_0) \in \text{Pic}(X)$ as $\eta^*(e_0) = d\pi^*(e_0) - \sum m_i e_{p_i}$, where d is the degree

of the linear system and $m_1, \dots, m_r \geq 1$ are the multiplicities of the linear system at the points p_1, \dots, p_r (see Remark 2.11). The fact that for each non-trivial $f \in \text{Bir}(\mathbb{P}^2)$, we can choose a general point $p \in \mathbb{P}^2$, sent by f onto another point q , yields $f(e_p) = e_q \neq e_p$ and shows that the action is faithful. \square

Example 2.15. Let $\sigma: [x : y : z] \dashrightarrow [yz : xz : xy]$ be the standard quadratic transformation of \mathbb{P}^2 . Its base-points are $p_1 = [1 : 0 : 0]$, $p_2 = [0 : 1 : 0]$, $p_3 = [0 : 0 : 1]$. We then write

$$X = \{([x_0 : x_1 : x_2], [y_0 : y_1 : y_2]) \mid x_0 y_0 = x_1 y_1 = x_2 y_2\}$$

and denote by $\pi: X \rightarrow \mathbb{P}^2$ and $\eta: X \rightarrow \mathbb{P}^2$ the first and second projections, which are blow-ups of p_1, p_2, p_3 and satisfy $\eta = \sigma \circ \pi$. There are six (-1) -curves $E_1, E_2, E_3, F_1, F_2, F_3$ on X , where $E_i = \pi^{-1}(p_i)$ and $F_i = \eta^{-1}(p_i)$ for $i = 1, 2, 3$.

The action of σ on $\mathcal{Z}_{\mathbb{P}^2}$ is given as follows. Firstly we have $\sigma_{\bullet}(e_0) = 2e_0 - e_{p_1} - e_{p_2} - e_{p_3}$ (Lemma 2.14). Secondly we have $\sigma_{\bullet}(e_{p_1}) = e_0 - e_{p_2} - e_{p_3}$ thanks to the following correspondant equality $E_1 = \eta^*(e_0) - F_2 - F_3$ holding in $\text{Pic}(X)$. This latter equality holds because E_1 is the strict transform of the line through p_2, p_3 by η . Similarly, we obtain $\sigma_{\bullet}(e_{p_2}) = e_0 - e_{p_1} - e_{p_3}$ and $\sigma_{\bullet}(e_{p_3}) = e_0 - e_{p_1} - e_{p_2}$.

For all other points $q \in \mathcal{B}(\mathbb{P}^2) \setminus \{p_1, p_2, p_3\}$, we have $\sigma(e_q) = e_{q'}$, for some $q' \in \mathcal{B}(\mathbb{P}^2)$.

2.2. The infinite Weyl group.

Definition 2.16. Denote by $\text{Aut}(\mathcal{Z}_{\mathbb{P}^2})$ the group of linear automorphisms of the \mathbb{Z} -module $\mathcal{Z}_{\mathbb{P}^2}$ and define $\text{Sym}_{\mathbb{P}^2} \subseteq \text{Aut}(\mathcal{Z}_{\mathbb{P}^2})$ to be the subgroup of elements that fix e_0 and permute the e_p , $p \in \mathcal{B}(\mathbb{P}^2)$.

We define $W_{\infty} \subseteq \text{Aut}(\mathcal{Z}_{\mathbb{P}^2})$ to be the *infinite Weyl group* generated by $\text{Bir}(\mathbb{P}^2)$ and the group $\text{Sym}_{\mathbb{P}^2}$.

Remark 2.17. Note that $\text{Aut}(\mathbb{P}^2) = \text{Sym}_{\mathbb{P}^2} \cap \text{Bir}(\mathbb{P}^2)$. Moreover, the Noether-Castelnuovo theorem yields $\text{Bir}(\mathbb{P}^2) = \langle \text{Aut}(\mathbb{P}^2), \sigma \rangle$, which implies that $W_{\infty} = \langle \text{Sym}_{\mathbb{P}^2}, \sigma \rangle$. Later on (see Corollary 2.28), we will prove that $W_{\infty} = \text{Sym}_{\mathbb{P}^2} \text{Bir}(\mathbb{P}^2) \text{Sym}_{\mathbb{P}^2}$.

Definition 2.18. Let $f \in W_{\infty}$ and $q \in \mathcal{B}(\mathbb{P}^2)$. We define the *degree* of f to be $\deg f = e_0 \cdot f^{-1}(e_0) \in \mathbb{Z}$ and the *multiplicity of f at q* to be $m_q(f) = e_q \cdot f^{-1}(e_0) \in \mathbb{Z}$. We denote $\text{Base}(f) \subseteq \mathcal{B}(\mathbb{P}^2)$ the set of points q such that $m_q(f) \neq 0$.

Remark 2.19. By construction, the degree, base-points and multiplicities of $f \in W_{\infty}$ are the same as for $f^{-1}(e_0) \in \mathcal{Z}_{\mathbb{P}^2}$ (which were defined in Definition 2.10). By Lemma 2.14, this definition coincides with the classical definition if $f \in \text{Bir}(\mathbb{P}^2)$.

Lemma 2.20.

- (1) Every element of W_{∞} preserves the intersection form and the canonical form.
- (2) For each $f \in W_{\infty}$ we have

$$f^{-1}(e_0) = (\deg f) \cdot e_0 - \sum_{q \in \text{Base}(f)} m_q(f) \cdot e_q$$

and the following equalities hold (Noether equalities), where $d = \deg f$:

$$\sum_{q \in \text{Base}(f)} m_q(f) = 3(d - 1), \quad \sum_{q \in \text{Base}(f)} (m_q(f))^2 = d^2 - 1.$$

- (3) For each $f \in W_{\infty}$, we have $\deg f^{-1} = \deg f$.

- (4) $\text{Sym}_{\mathbb{P}^2} = \{f \in W_\infty \mid f(e_0) = e_0\} = \{f \in W_\infty \mid \deg(f) = 1\} = \{f \in W_\infty \mid \deg(f) = \pm 1\}$.
- (5) For each $\alpha \in \text{Sym}_{\mathbb{P}^2}$, we have $\sigma \circ \alpha \circ \sigma \in \text{Sym}_{\mathbb{P}^2}$ if and only if α preserves the set $\{e_{[1:0:0]}, e_{[0:1:0]}, e_{[0:0:1]}\}$.

Proof. (1): Follows from the fact that $\text{Bir}(\mathbb{P}^2)$ and $\text{Sym}_{\mathbb{P}^2}$ preserve the intersection form and the canonical form.

(2): The first equality follows from Definition 2.18 and from the next identity:

$$\forall a \in \mathcal{Z}_{\mathbb{P}^2}, a = (a \cdot e_0) e_0 - \sum_{q \in \mathcal{B}(\mathbb{P}^2)} (a \cdot e_q) e_q.$$

The Noether equalities follows from (1), since $f^{-1}(e_0)^2 = d^2 - \sum (m_i)^2 = (e_0)^2 = 1$ and $\omega(f^{-1}(e_0)) = -3d + \sum m_i = \omega(e_0) = -3$.

(3): We have $\deg f = e_0 \cdot f^{-1}(e_0) = f(e_0) \cdot e_0 = \deg f^{-1}$.

(4): Let $f \in W_\infty$ be such that $\deg f = d = \pm 1$. It follows successively from the Noether equalities that all multiplicities $m_q(f)$ are zero, $d = 1$ and $f(e_0) = e_0$. For each $p \in \mathcal{B}(\mathbb{P}^2)$ we have $f(e_p) \cdot e_0 = 0$, so $f(e_p) = \sum_{i=1}^r a_i e_{q_i}$, for some $q_1, \dots, q_r \in \mathcal{B}(\mathbb{P}^2)$. As $-1 = (e_p)^2 = (f(e_p))^2 = -\sum (a_i)^2$, we find that $f(e_p) = \pm e_{q_i}$ for some i . Since $\omega(e_p) = \omega(f(e_p))$, we get $f(e_p) = e_{q_i}$. Hence, $f \in \text{Sym}_{\mathbb{P}^2}$. (5): By (4), we have $\sigma \circ \alpha \circ \sigma \in \text{Sym}_{\mathbb{P}^2}$ if and only if $\sigma \circ \alpha \circ \sigma(e_0) = e_0$. Since this corresponds to $\alpha \circ \sigma(e_0) = \sigma(e_0)$, the result follows from the equality $\sigma(e_0) = 2e_0 - e_{[1:0:0]} - e_{[0:1:0]} - e_{[0:0:1]}$ (Example 2.15). \square

Corollary 2.21. Let $f, g \in W_\infty$. The following conditions are equivalent:

- (1) $f^{-1}(e_0) = g^{-1}(e_0)$.
- (2) There exists $\alpha \in \text{Sym}_{\mathbb{P}^2}$ such that $g = \alpha \circ f$.

Proof. Let us write $\alpha = g \circ f^{-1}$. By Lemma 2.20(4), $\alpha \in \text{Sym}_{\mathbb{P}^2}$ if and only if $\alpha(e_0) = e_0$. Applying g^{-1} , this condition is equivalent to $f^{-1}(e_0) = g^{-1}(e_0)$. \square

Corollary 2.22. If $f, g \in W_\infty$, we have

$$\deg f \circ g^{-1} = (\deg f)(\deg g) - \sum_{q \in \mathcal{B}(\mathbb{P}^2)} m_q(f) m_q(g).$$

Proof. We have $\deg f \circ g^{-1} = e_0 \cdot (f \circ g^{-1})^{-1}(e_0) = f^{-1}(e_0) \cdot g^{-1}(e_0)$, so that the result follows from Lemma 2.20(2). \square

Lemma 2.23. Let $g \in W_\infty$ and let $q \in \mathcal{B}(\mathbb{P}^2)$.

- (1) If $q \in \text{Base}(g)$, then $g(e_q) = m_q(g) e_0 - \sum_{p \in \text{Base}(g^{-1})} a_p e_p$, $a_p \in \mathbb{Z}$.
- (2) If $q \notin \text{Base}(g)$, then $g(e_q) = e_{\tilde{q}}$ for some $\tilde{q} \in \mathcal{B}(\mathbb{P}^2) \setminus \text{Base}(g^{-1})$.

In particular, g induces a bijection $\mathcal{B}(\mathbb{P}^2) \setminus \text{Base}(g) \rightarrow \mathcal{B}(\mathbb{P}^2) \setminus \text{Base}(g^{-1})$.

Proof. We write $g(e_q) = d e_0 + \sum a_i e_{p_i}$, for some $\{p_1, \dots, p_n\} \subseteq \mathcal{B}(\mathbb{P}^2)$. We then observe that $d = e_0 \cdot g(e_q) = g^{-1}(e_0) \cdot e_q = m_q(g)$.

If $q \notin \text{Base}(g)$, we then obtain $g(e_q) = \sum a_i e_{p_i}$. Since $1 = -\omega(e_q) = -\omega(\sum a_i e_{p_i}) = \sum a_i$ and $1 = -(e_q)^2 = \sum (a_i)^2$, we find that $g(e_q)$ is equal to $e_{\tilde{q}}$ for some $\tilde{q} \in \mathcal{B}(\mathbb{P}^2)$. Moreover, $\tilde{q} \notin \text{Base}(g^{-1})$, since $m_{\tilde{q}}(g^{-1}) = e_{\tilde{q}} \cdot g(e_0) = e_q \cdot e_0 = 0$. This yields (2).

To get (1), we consider the case $q \in \text{Base}(g)$ and need to show that if $p_i \notin \text{Base}(g^{-1})$, then $a_i = 0$. This is because $a_i = e_{p_i} \cdot g(e_q) = g^{-1}(e_{p_i}) \cdot e_q$ and because $g^{-1}(e_{p_i})$ is equal to $e_{\tilde{p}_i}$ for some $\tilde{p}_i \in \mathcal{B}(\mathbb{P}^2) \setminus \text{Base}(g)$ by (2). \square

Corollary 2.24. *For each $g \in W_\infty$ and each $q \in \mathcal{B}(\mathbb{P}^2)$, we have*

$$q \notin \text{Base}(g) \Leftrightarrow g(e_q) = e_{\tilde{q}} \text{ for some } \tilde{q} \in \mathcal{B}(\mathbb{P}^2).$$

Proof. Follows from Lemma 2.23. \square

Corollary 2.25. *Let $f, g \in W_\infty$ be such that $\text{Base}(f) \subseteq \text{Base}(g^{-1})$, then we have*

$$\text{Base}(f \circ g) \subseteq \text{Base}(g).$$

Proof. Take $q \in \mathcal{B}(\mathbb{P}^2) \setminus \text{Base}(g)$. Then, by Lemma 2.23, we have $g(e_q) = e_{\tilde{q}}$ for some $\tilde{q} \in \mathcal{B}(\mathbb{P}^2) \setminus \text{Base}(g^{-1})$. It follows that $\tilde{q} \in \mathcal{B}(\mathbb{P}^2) \setminus \text{Base}(f)$, so that $f(e_{\tilde{q}}) = e_{\tilde{\tilde{q}}}$ for some $\tilde{\tilde{q}} \in \mathcal{B}(\mathbb{P}^2)$, i.e. $f \circ g(e_q) = e_{\tilde{\tilde{q}}}$, proving that $q \notin \text{Base}(f \circ g)$. \square

As explained before, the infinite Weyl group W_∞ contains $\text{Bir}(\mathbb{P}^2)$. In some sense, this corresponds to forgetting the configuration of points. However, several properties of the action of $\text{Bir}(\mathbb{P}^2)$ on $\mathcal{Z}_{\mathbb{P}^2}$ extend to W_∞ . For instance, the Noether equalities (Lemma 2.20(2)) are fulfilled by any element of W_∞ . A priori, the degree and multiplicities could be negative, but we will show that it is not the case (Lemma 2.26). Also, there are some elements of $\mathcal{Z}_{\mathbb{P}^2}$ which satisfy the Noether equalities but which are not in the orbit of e_0 . However, there is an algorithm to decide if an element is in this orbit (Algorithm 3.7 below, corresponding to the classical Hudson test).

Lemma 2.26. *Let $f \in W_\infty \setminus \text{Sym}_{\mathbb{P}^2}$.*

(1) *For each finite set $\Delta = \{q_1, \dots, q_s\} \subseteq \mathcal{B}(\mathbb{P}^2)$ of $s \geq 1$ points there exists a dense open set $U \subseteq (\mathbb{P}^2)^s$ such that for each $(p_1, \dots, p_s) \in U$:*

- (i) *The points p_1, \dots, p_s are distinct;*
- (ii) *There exists an element $g \in \text{Bir}(\mathbb{P}^2)$ satisfying*

$$\deg g = \deg f \quad \text{and} \quad m_{p_i}(g) = m_{q_i}(f) \text{ for } i = 1, \dots, s.$$

(2) *The degree and multiplicities of f satisfy*

$$\deg f \geq 2 \quad \text{and} \quad m_q(f) \geq 0 \text{ for each } q \in \mathcal{B}(\mathbb{P}^2).$$

(3) *There exist $\alpha, \beta \in \text{Sym}_{\mathbb{P}^2}$ and $g \in \text{Bir}(\mathbb{P}^2)$, such that $f = \alpha \circ g \circ \beta$.*

Proof. Let us first observe that (2) and (3) follow from (1). Indeed, take for Δ the set $\text{Base}(f) = \{q_1, \dots, q_s\}$ and let U be the corresponding open subset of $(\mathbb{P}^2)^s$ given by (1). Choose $(p_1, \dots, p_s) \in U$ and choose $g \in \text{Bir}(\mathbb{P}^2)$ such that $\deg g = \deg f$ and $m_{p_i}(g) = m_{q_i}(f)$ for $i = 1, \dots, s$. Then choose $\beta \in \text{Sym}_{\mathbb{P}^2}$ that sends e_{q_i} onto e_{p_i} for each i , and note that $\beta \circ f^{-1}(e_0) = g^{-1}(e_0)$. Hence, the element $\alpha = f \circ \beta^{-1} \circ g^{-1}$ belongs to $\text{Sym}_{\mathbb{P}^2}$ by Lemma 2.20(4).

To prove (1), we write $f = \alpha_l \circ \sigma \circ \dots \circ \alpha_1 \circ \sigma \circ \alpha_0$ where $l \geq 1$, $\alpha_0, \dots, \alpha_l \in \text{Sym}_{\mathbb{P}^2}$ and prove the result by induction on l . As the result does not change under right or left-multiplication by elements of $\text{Sym}_{\mathbb{P}^2}$, we can moreover assume that α_0 and α_l are equal to the identity. We can also always enlarge the set Δ .

If $l = 1$, then $f = \sigma$, so that f has degree 2 and three base-points of multiplicity 1 (see Example 2.15). We may assume that q_1, q_2, q_3 are the base-points of f . Then, we can choose for U the open subset of points (p_1, \dots, p_s) in $(\mathbb{P}^2)^s$ where p_1, \dots, p_s are distinct and where p_1, p_2, p_3 are not collinear. For each $(p_1, \dots, p_s) \in U$, we choose an element $\alpha \in \text{Aut}(\mathbb{P}^2)$ sending p_i onto q_i for $i = 1, 2, 3$ and choose $g = f \circ \alpha$.

For $l \geq 2$, we write $f = f' \circ \sigma$ and apply the induction hypothesis to f' . Up to enlarging $\Delta = \{q_1, \dots, q_s\}$, we may assume that $q_1 = [1 : 0 : 0]$, $q_2 = [0 : 1 : 0]$, $q_3 = [0 : 0 : 1]$. For each $i \geq 4$, define q'_i as the unique point of $\mathcal{B}(\mathbb{P}^2)$ such that $\sigma(e_{q_i}) = e_{q'_i}$. For $i = 1, 2, 3$, set $q'_i = q_i$. One can assume that $\text{Base}(f') \subseteq \{q'_1, \dots, q'_s\}$. Let $U' \subseteq (\mathbb{P}^2)^s$ be an open subset associated to f' and $\Delta' = \{q'_1, \dots, q'_s\}$ via the induction hypothesis.

Let $T \subseteq (\mathbb{P}^2)^3$ be the open subset given by

$$T = \left\{ ([1 : a_1 : a_2], [1 : a_3 : a_4], [1 : a_5 : a_6]) \mid a_1, \dots, a_6 \in \mathbb{k}, \det \begin{pmatrix} 1 & 1 & 1 \\ a_1 & a_3 & a_5 \\ a_2 & a_4 & a_6 \end{pmatrix} \neq 0 \right\}$$

and let $\rho: T \rightarrow \text{Aut}(\mathbb{P}^2) = \text{PGL}_3(\mathbb{k})$ be the morphism defined by

$$\rho([1 : a_1 : a_2], [1 : a_3 : a_4], [1 : a_5 : a_6]) = \begin{pmatrix} 1 & 1 & 1 \\ a_1 & a_3 & a_5 \\ a_2 & a_4 & a_6 \end{pmatrix}.$$

If $\mathcal{T} = (p_1, p_2, p_3)$ belongs to T , we set

$$\sigma_{\mathcal{T}} := \rho(\mathcal{T}) \circ \sigma \circ \rho(\mathcal{T})^{-1} \in \text{Bir}(\mathbb{P}^2).$$

Note that $\sigma_{\mathcal{T}}$ is a quadratic involution having base-points at p_1, p_2, p_3 .

We then denote by $U \subseteq (\mathbb{P}^2)^s$ the dense open subset of s -uples $p = (p_1, \dots, p_s)$ such that:

- (1) No three of the points p_i are collinear (so that in particular the points p_i are distinct) ;
- (2) The triple $\mathcal{T} = (p_1, p_2, p_3)$ belongs to T ;
- (3) The s -uple $p' = (p'_1, \dots, p'_s)$ belongs to U' , where the elements p'_i are defined by $p'_i := p_i$ for $i \leq 3$ and by $p'_i := \sigma_{\mathcal{T}}(p_i) \in \mathbb{P}^2$ for $i \geq 4$.

For each $p \in U$, the corresponding $p' \in U'$ yields an element $g' \in \text{Bir}(\mathbb{P}^2)$ satisfying $\deg g' = \deg f'$ and $m_{q'_i}(f') = m_{p'_i}(g')$ for each i . Taking $\beta' \in \text{Sym}_{\mathbb{P}^2}$ that sends $e_{q'_i}$ onto $e_{p'_i}$ for each i , the fact that $\text{Base}(f') \subseteq \{q'_1, \dots, q'_s\}$ implies as before that $\beta'(f'^{-1}(e_0)) = g'^{-1}(e_0)$, so $f' = \alpha \circ g' \circ \beta'$, for some $\alpha \in \text{Sym}_{\mathbb{P}^2}$.

We write $\nu = \rho(\mathcal{T}) \in \text{Aut}(\mathbb{P}^2)$, $\sigma_{\mathcal{T}} = \nu \circ \sigma \circ \nu^{-1} \in \text{Bir}(\mathbb{P}^2)$ as before and obtain

$$f = f' \circ \sigma = \alpha \circ g' \circ \beta' \circ \sigma = \alpha \circ g \circ \beta,$$

where $g = g' \circ \sigma_{\mathcal{T}} \in \text{Bir}(\mathbb{P}^2)$ and $\beta = \sigma_{\mathcal{T}} \circ \beta' \circ \sigma = \nu \circ \sigma \circ \nu^{-1} \circ \beta' \circ \sigma \in W_{\infty}$. For $i \in \{1, 2, 3\}$, both ν and β' send e_{q_i} to e_{p_i} , hence $\nu^{-1} \circ \beta'$ fixes e_{q_i} . This shows that $\beta \in \text{Sym}_{\mathbb{P}^2}$ (Lemma 2.20(5)), and thus that $\deg g = \deg f$.

It remains to observe that β sends e_{q_i} to e_{p_i} for each i , to obtain $m_{p_i}(g) = m_{q_i}(f)$ for each i . The fact that $\nu^{-1} \circ \beta'$ fixes $e_0 - e_{q_1} - e_{q_2}$ implies that $\sigma \circ \nu^{-1} \circ \beta' \circ \sigma$ fixes $\sigma(e_0 - e_{q_1} - e_{q_2}) = e_{q_3}$ (see Example 2.15) and thus that β sends e_{q_3} to e_{p_3} . The same works for e_{q_1}, e_{q_2} . For $i \geq 4$, we have

$$\beta(e_{q_i}) = \sigma_{\mathcal{T}} \circ \beta' \circ \sigma(e_{q_i}) = \sigma_{\mathcal{T}} \circ \beta'(e_{q'_i}) = \sigma_{\mathcal{T}}(e_{p'_i}) = e_{p_i}. \quad \square$$

The first two corollaries are stated for an easier reading. The first one is [BlaCal2016, Proposition 2.4]:

Corollary 2.27. *For each homaloidal type $(d; m_1, \dots, m_s)$, there exists a dense open subset $U \subseteq (\mathbb{P}^2)^s$ such that for each $(p_1, \dots, p_s) \in U$ the following holds:*

- (1) *The points p_1, \dots, p_s are distinct;*
- (2) *There exists a Cremona transformation $f \in \text{Bir}(\mathbb{P}^2)$ such that*

$$f^{-1}(e_0) = de_0 - \sum_{i=1}^s m_i e_{p_i}.$$

Corollary 2.28. *We have $W_\infty = \text{Sym}_{\mathbb{P}^2} \text{Bir}(\mathbb{P}^2) \text{Sym}_{\mathbb{P}^2}$.*

In the next two corollaries, we give information on the orbits of e_0, e_q and $e_0 - e_q$, where q is a point of $\mathcal{B}(\mathbb{P}^2)$. The first one is the following positivity result on the degree and multiplicities of elements in the orbit of e_0 , which also follows from [BlaCan2016, Lemma 5.3] (with another proof).

Corollary 2.29. *Each element $a \in W_\infty(e_0)$ can be written as*

$$a = (\deg a) \cdot e_0 - \sum_{q \in \text{Base}(a)} m_q(a) \cdot e_q,$$

where $\deg a \geq 1$, $m_q(a) \geq 1$ for each $q \in \text{Base}(a)$ and

$$\sum_{q \in \text{Base}(a)} m_q(a) = 3(\deg a - 1), \quad \sum_{q \in \text{Base}(a)} (m_q(a))^2 = \deg(a)^2 - 1.$$

Moreover, for any two distinct $q, q' \in \mathcal{B}(\mathbb{P}^2)$ we have $m_q(a) + m_{q'}(a) \leq \deg a$.

Proof. We write $a \in W_\infty(e_0)$ as $a = f(e_0)$ for some $f \in W_\infty$, and write $f = \alpha \circ g \circ \beta$, where $\alpha, \beta \in \text{Sym}_{\mathbb{P}^2}$ and $g \in \text{Bir}(\mathbb{P}^2)$, using Lemma 2.26(3). Hence, $a = \alpha \circ g(e_0)$. The description above follows then from Lemmas 2.14 and 2.20. The inequality $m_q(a) + m_{q'}(a) \leq \deg(a)$ can be checked for $g(e_0)$, since $\alpha \in \text{Sym}_{\mathbb{P}^2}$. We can moreover assume that $q \in \mathbb{P}^2$ and q' is either in \mathbb{P}^2 or in the first blow-up of q (since $m_q(g(e_0)) \leq m_{q'}(g(e_0))$ if $q \geq q'$). The result follows then from Bézout theorem, by intersecting the line through q and q' with the linear system corresponding to $g(e_0)$. \square

The following result is obvious for orbits of $\text{Bir}(\mathbb{P}^2)$ and here is generalised to orbits of W_∞ . This allows to say that elements of W_∞ have a behaviour “not too far” from elements of $\text{Bir}(\mathbb{P}^2)$. See also Lemma 2.26(3) for another result in this direction.

Corollary 2.30. *Let $q \in \mathcal{B}(\mathbb{P}^2)$ and let $a \in W_\infty(e_0)$.*

- (1) *For each $b \in W_\infty(e_0)$, we have $a \cdot b \geq 1$.*
- (2) *For each $b \in W_\infty(e_0 - e_q)$, we have $a \cdot b \geq 1$.*
- (3) *For each $b \in W_\infty(e_q)$, we have $a \cdot b \geq 0$.*

Proof. We apply an element of W_∞ and assume that b is equal to $e_0, e_0 - e_q, e_q$ respectively. By Corollary 2.29, we have $a = (\deg a) \cdot e_0 - \sum_{p \in \text{Base}(a)} m_p(a) \cdot e_p$, where $\deg a \geq 1$, $m_p(a) \geq 1$ for each $p \in \text{Base}(a)$ and $\sum_{p \in \text{Base}(a)} (m_p(a))^2 = \deg(a)^2 - 1$. We then find that $a \cdot b$ is equal to $\deg a, \deg a - m_q(a), m_q(a)$ respectively. Assertions (1), (2), (3) are then given by $\deg a \geq 1, (m_q(a))^2 \leq \deg(a)^2 - 1$ and $m_q(a) \geq 0$. \square

2.3. Jonquières elements viewed in the Weyl group. We now define the analogue of the groups $\text{Jonq}_p \subseteq \text{Bir}(\mathbb{P}^2)$ in the Weyl group:

Definition 2.31. For each $q \in \mathcal{B}(\mathbb{P}^2)$ we define $J_q \subseteq W_\infty$ as the subgroup

$$J_q = \{\varphi \in W_\infty \mid \varphi(e_0 - e_q) = e_0 - e_q\}.$$

Lemma 2.32. *For each $q \in \mathbb{P}^2$, we have $\text{Jonq}_q = J_q \cap \text{Bir}(\mathbb{P}^2)$.*

Proof. Let $\pi: \mathbb{P}^2 \dashrightarrow \mathbb{P}^1$ be the projection from q , and let $\eta: X \rightarrow \mathbb{P}^2$ be the blow-up of q . The result follows from the fact that $e_0 - e_q \in \mathcal{Z}_{\mathbb{P}^2}$ corresponds to the divisor of $\text{Pic}(X)$ corresponding to the fibres of the morphism $\pi \circ \eta: X \rightarrow \mathbb{P}^1$. \square

Definition 2.33. For each $q \in \mathcal{B}(\mathbb{P}^2)$ and for each finite set $\Delta \subseteq \mathcal{B}(\mathbb{P}^2) \setminus \{q\}$ of even order $2n$, we define $\iota_{q,\Delta} \in J_q$ to be the involution given by

$$\begin{aligned} \iota_{q,\Delta}(e_0) &= (n+1)e_0 - ne_q - \sum_{r \in \Delta} e_r, & \iota_{q,\Delta}(e_r) &= e_0 - e_q - e_r, r \in \Delta, \\ \iota_{q,\Delta}(e_q) &= ne_0 - (n-1)e_q - \sum_{r \in \Delta} e_r, & \iota_{q,\Delta}(e_r) &= e_r, r \in \mathcal{B}(\mathbb{P}^2) \setminus (\Delta \cup \{q\}). \end{aligned}$$

Remark 2.34. In order to see that the elements $\iota_{q,\Delta}$ belong to $J_q \subseteq W_\infty$, we can observe that $\iota_{q,\emptyset}$ is the identity, that $\iota_{q,\Delta}$ is equal to σ , up to left and right-multiplication by elements of $\text{Sym}_{\mathbb{P}^2}$ when Δ contains 2 elements, and that $\iota_{q,\Delta} \circ \iota_{q,\Delta'} = \iota_{q,(\Delta \cup \Delta') \setminus (\Delta \cap \Delta')}$.

Definition 2.35. Let $p_1, p_2, p_3 \in \mathcal{B}(\mathbb{P}^2)$ be 3 distinct points. We define $\sigma_{p_1,p_2,p_3} \in W_\infty$ as the involution given by

$$\begin{aligned} \sigma_{p_1,p_2,p_3}(e_0) &= 2e_0 - e_{p_1} - e_{p_2} - e_{p_3}, & \sigma_{p_1,p_2,p_3}(e_{p_1}) &= e_0 - e_{p_2} - e_{p_3} \\ \sigma_{p_1,p_2,p_3}(e_{p_2}) &= e_0 - e_{p_1} - e_{p_3}, & \sigma_{p_1,p_2,p_3}(e_{p_3}) &= e_0 - e_{p_1} - e_{p_2}, \\ \sigma_{p_1,p_2,p_3}(e_r) &= e_r, r \in \mathcal{B}(\mathbb{P}^2) \setminus \{p_1, p_2, p_3\}, \end{aligned}$$

We observe that $\sigma_{p_1,p_2,p_3} \in J_{p_i}$ for $i = 1, 2, 3$, and that $\sigma_{p_1,p_2,p_3} = \tau_{p_2,p_3} \circ \iota_{p_1,\{p_2,p_3\}}$, where $\tau_{p_2,p_3} \in \text{Sym}_{\mathbb{P}^2}$ is the transposition permuting p_2 and p_3 .

Remark 2.36. When $p_1 = [1 : 0 : 0]$, $p_2 = [0 : 1 : 0]$, $p_3 = [0 : 0 : 1]$, we observe that σ_{p_1,p_2,p_3} is similar to the standard quadratic involution $\sigma: [x : y : z] \dashrightarrow [yz : xz : xy]$. It is however not realised by an element of $\text{Bir}(\mathbb{P}^2)$ as it fixes all points of $\mathbb{P}^2 \setminus \{p_1, p_2, p_3\}$. Moreover, σ_{p_1,p_2,p_3} and $\iota_{p_1,\{p_2,p_3\}}$ both belong to $\text{Sym}_{\mathbb{P}^2}$ $\sigma = \{\alpha \circ \sigma \mid \alpha \in \text{Sym}_{\mathbb{P}^2}\} \subseteq W_\infty$.

Lemma 2.37.

- (1) For each $q \in \mathcal{B}(\mathbb{P}^2)$ and each $\varphi \in J_q$, we have $m_q(\varphi) = \deg(\varphi) - 1$.
- (2) For each $q \in \mathcal{B}(\mathbb{P}^2)$ and each $\varphi \in W_\infty$ with $m_q(\varphi) = \deg(\varphi) - 1$, the set $\Delta = \text{Base}(\varphi) \setminus \{q\}$ has even cardinality $2n \geq 0$ and

$$\varphi^{-1}(e_0) = (\iota_{q,\Delta})^{-1}(e_0) = (n+1)e_0 - ne_q - \sum_{r \in \Delta} e_r = e_0 + \sum_{r \in \Delta} \left(\frac{e_0 - e_q}{2} - e_r \right).$$

This yields the existence of $\alpha \in \text{Sym}_{\mathbb{P}^2}$ such that

$$\varphi = \alpha \circ \iota_{q,\Delta}.$$

Moreover, $\alpha \in J_q$ if and only if $\varphi \in J_q$.

Proof. (1) The fact that $\varphi \in J_q$ implies that

$$\deg(\varphi) - m_q(\varphi) = (e_0 - e_q) \cdot \varphi^{-1}(e_0) = (e_0 - e_q) \cdot e_0 = 1.$$

(2) Since $\Delta = \text{Base}(\varphi) \setminus \{q\}$ and $m_q(\varphi) = \deg(\varphi) - 1$, we can write

$$\varphi^{-1}(e_0) = (n+1)e_0 - ne_q - \sum_{r \in \Delta} m_r e_r$$

where $n \geq 0$ and $m_r \geq 1$ for each $r \in \Delta$ (Corollary 2.29). The Noether equalities (Lemma 2.20(2)) yield $\sum_{r \in \Delta} m_r = \sum_{r \in \Delta} (m_r)^2 = 2n$, so $m_r = 1$ for each $r \in \Delta$, and thus Δ contains $2n$ elements.

Since $\varphi^{-1}(e_0) = (\iota_{q,\Delta})^{-1}(e_0)$, we have $\varphi = \alpha \circ \iota_{q,\Delta}$ for some $\alpha \in \text{Sym}_{\mathbb{P}^2}$ (Corollary 2.21). Since $\iota_{q,\Delta} \in J_q$, we have $\alpha \in J_q$ if and only if $\varphi \in J_q$. \square

Corollary 2.38. *Any element $\varphi \in J_q$ admits an expression*

$$\varphi = \alpha \circ \iota_{q,\Delta},$$

where $\Delta := \text{Base}(\varphi) \setminus \{q\}$ has even order and $\alpha \in \text{Sym}_{\mathbb{P}^2} \cap J_q = \{\beta \in \text{Sym}_{\mathbb{P}^2}, \beta(q) = q\}$.

Proof. Directly follows from Lemma 2.37. \square

Corollary 2.39. *If $q, q' \in \mathcal{B}(\mathbb{P}^2)$ are two distinct points, then $J_q \cap J_{q'}$ consists of elements of degree 1 or 2.*

Proof. It follows from Lemma 2.37 that if $\varphi \in J_q$ is a Jonquière's element with $\deg \varphi \geq 3$, the multiplicity at q is $\deg \varphi - 1 \geq 2$ and q is the unique point having this multiplicity. \square

We now give the following definition, which generalise the one of Jonquière's elements of $\text{Bir}(\mathbb{P}^2)$, as Lemma 2.42 explains.

Definition 2.40. An element $\varphi \in W_\infty$ is said to be a *Jonquière's element* if there exists a point $q \in \mathcal{B}(\mathbb{P}^2)$ such that $m_q(\varphi) = \deg(\varphi) - 1$.

Lemma 2.41. *Let $\psi \in W_\infty$. The following conditions are equivalent:*

- (1) ψ is a Jonquière's element of W_∞ ;
- (2) There exist $\alpha, \beta \in \text{Sym}_{\mathbb{P}^2}$, $q \in \mathcal{B}(\mathbb{P}^2)$ and $\varphi \in J_q$ such that $\psi = \alpha \circ \varphi \circ \beta$;
- (3) There exist $\alpha \in \text{Sym}_{\mathbb{P}^2}$, $q \in \mathcal{B}(\mathbb{P}^2)$ and $\varphi \in J_q$ such that $\psi = \alpha \circ \varphi$;
- (4) There exist $\alpha \in \text{Sym}_{\mathbb{P}^2}$, $q \in \mathcal{B}(\mathbb{P}^2)$ and $\varphi \in J_q$ such that $\psi = \varphi \circ \alpha$;
- (5) There exist $\alpha \in \text{Sym}_{\mathbb{P}^2}$, $q \in \mathcal{B}(\mathbb{P}^2)$ and a finite set of even order $\Delta \subseteq \mathcal{B}(\mathbb{P}^2) \setminus \{q\}$ such that $\psi = \alpha \circ \iota_{q,\Delta}$.

Proof. (1) \Rightarrow (5) is given by Lemma 2.37(2); (5) \Rightarrow (3) is given by the fact that $\iota_{q,\Delta} \in J_q$ and (3) \Rightarrow (2) follows by taking $\beta = \text{id}$.

(2) \Rightarrow (4): Writing $\varphi' = \alpha \circ \varphi \circ \alpha^{-1}$, we have $\varphi'(e_0 - e_p) = e_0 - e_p$ where $p \in \mathcal{B}(\mathbb{P}^2)$ is the element such that $\alpha(e_q) = e_p$. Hence $\psi = \varphi' \circ \alpha'$ where $\varphi' \in J_p$ and $\alpha' = \alpha \circ \beta \in \text{Sym}_{\mathbb{P}^2}$.

(4) \Rightarrow (1): Taking $p \in \mathcal{B}(\mathbb{P}^2)$ such that $\alpha(e_p) = e_q$ we get

$$m_p(\psi) = e_p \cdot \psi^{-1}(e_0) = \alpha(e_p) \cdot \varphi^{-1}(e_0) = e_q \cdot \varphi^{-1}(e_0) = m_q(\varphi) \stackrel{\text{Lemma 2.37(1)}}{=} \deg(\varphi) - 1.$$

It remains to observe that

$$\deg(\psi) = e_0 \cdot \alpha^{-1}(\varphi^{-1}(e_0)) = \alpha(e_0) \cdot \varphi^{-1}(e_0) = e_0 \cdot \varphi^{-1}(e_0) = \deg(\varphi). \quad \square$$

Lemma 2.42. *Let $f \in \text{Bir}(\mathbb{P}^2)$. The following conditions are equivalent:*

- (1) f is a Jonquière's element of $\text{Bir}(\mathbb{P}^2)$;
- (2) f is a Jonquière's element of W_∞ ;
- (3) There exist $\alpha, \beta \in \text{Aut}(\mathbb{P}^2)$, $q \in \mathbb{P}^2$, and $\varphi \in \text{Jonq}_q \subseteq \text{Bir}(\mathbb{P}^2)$ such that $f = \alpha \circ \varphi \circ \beta$;
- (4) There exist $\alpha \in \text{Aut}(\mathbb{P}^2)$, $q \in \mathbb{P}^2$, and $\varphi \in \text{Jonq}_q \subseteq \text{Bir}(\mathbb{P}^2)$ such that $f = \alpha \circ \varphi$;
- (5) There exist $\alpha \in \text{Aut}(\mathbb{P}^2)$, $q \in \mathbb{P}^2$, and $\varphi \in \text{Jonq}_q \subseteq \text{Bir}(\mathbb{P}^2)$ such that $f = \varphi \circ \alpha$.

Proof. By definition, f is a Jonquières element of $\text{Bir}(\mathbb{P}^2)$ if and only if there exist two points $p, q \in \mathbb{P}^2$ such that the pencil of lines through p is sent to the pencil of lines through q . Composing at the source or the target with a linear automorphism exchanging p and q yields then an element of Jonq_p or Jonq_q . This yields the equivalence of (1),(3),(4),(5). As $\text{Aut}(\mathbb{P}^2) \subseteq \text{Sym}_{\mathbb{P}^2}$ and every Jonquières element of $\text{Bir}(\mathbb{P}^2)$ is a Jonquières element of W_∞ , we have (3) \Rightarrow (2) (Lemma 2.41). It remains then to prove (2) \Rightarrow (1). Assertion (2) implies that f has a base-point p of multiplicity $\deg(f) - 1$. We can moreover assume that p is a proper point of \mathbb{P}^2 (replacing p with the proper point $p' \in \mathbb{P}^2$ above which p lies only increases the multiplicity). The image of the pencil of lines through p is then a pencil of lines, passing thus through a point $q \in \mathbb{P}^2$. This achieves the proof. \square

Definition 2.43. Two elements a, a' of $\mathcal{Z}_{\mathbb{P}^2}$ are said to be *equal modulo* $\text{Sym}_{\mathbb{P}^2}$ if there exists an element $\alpha \in \text{Sym}_{\mathbb{P}^2}$ such that $a' = \alpha(a)$. This is written $a \equiv_{\text{Sym}_{\mathbb{P}^2}} a'$.

Remark 2.44. Two elements a, a' of $\mathcal{Z}_{\mathbb{P}^2}$ are equal modulo $\text{Sym}_{\mathbb{P}^2}$ if and only if they have the same degree and if there exists a bijection $t: \text{Base}(a) \rightarrow \text{Base}(a')$ such that $m_{t(p)}(a') = m_p(a)$ for each $p \in \text{Base}(a)$.

In particular, the set $W_\infty(e_0)/\text{Sym}_{\mathbb{P}^2}$ of equivalence classes modulo $\text{Sym}_{\mathbb{P}^2}$ in $W_\infty(e_0)$ corresponds to the set of homaloidal types of birational transformations of \mathbb{P}^2 (called *proper homaloidal types* in [Alb2002, BlaCal2016]).

Lemma 2.45. For any element $a \in \mathcal{Z}_{\mathbb{P}^2}$ and any Jonquières element $\varphi \in W_\infty$, the element $\varphi(a)$ is equal to some element $\iota_{q,\Delta}(a)$ modulo $\text{Sym}_{\mathbb{P}^2}$.

Proof. This is a direct consequence of Corollary 2.38 and Definition 2.43. \square

We will use the following easy observation twice in the sequel.

Lemma 2.46. Let $\chi = (d; m_0, \dots, m_r)$ be the homaloidal type of a birational transformation of \mathbb{P}^2 , and let us assume that $d \geq 2$ and that $m_0 \geq \dots \geq m_r \geq 1$. If $m_0 + m_r = d$, then $\chi = (d; d-1, \underbrace{1, \dots, 1}_{2d-2})$ is the homaloidal type of a Jonquières element. In particular, we remark for later use that $r = 2d - 2$ is even.

Proof. As $m_0 + m_i \leq d$ for each $i \geq 1$ (Corollary 2.29), we have $m_1 = m_2 = \dots = m_r = d - m_0$. The second Noether equality (Lemma 2.20(2)) then gives $d^2 - 1 = (d - m_1)^2 + r m_1^2$, whence $m_1(2d - m_1(r + 1)) = 1$, so $m_1 = 1$ and $r = 2d - 2$. \square

3. THE ALGORITHM THAT COMPUTES THE LENGTH AND THE PROOF OF THEOREM 1

In this section, we give the proof of Theorem 1, by first working in the infinite Weyl group introduced in Section 2 (in particular in §2.2) and get the analogue of Theorem 1 in W_∞ , namely Proposition 3.19. We then show (in Section 3.3) that the algorithm given in W_∞ can actually be applied in $\text{Bir}(\mathbb{P}^2)$.

3.1. Degree, maximal multiplicity and comultiplicity.

Definition 3.1. Let $a \in W_\infty$. We define the *maximal multiplicity* of a to be

$$m_{\max}(a) = \max\{m_q(a) \mid q \in \mathcal{B}(\mathbb{P}^2)\}$$

and say that a has *maximal multiplicity* at $q \in \mathcal{B}(\mathbb{P}^2)$ if $m_q(a) = m_{\max}(a)$.

We define the *comultiplicity* of a to be $\text{comult } a = \deg a - m_{\max}(a)$.

Lemma 3.2. *Let $a \in W_{\infty}(e_0)$.*

- (1) *If $\deg a = 1$, then $a = e_0$, $m_{\max}(a) = 0$ and $\text{comult } a = 1$.*
- (2) *If $\deg a > 1$, then $1 \leq m_{\max}(a) \leq \deg a - 1$ and $1 \leq \text{comult } a \leq \deg a - 1$.*
- (3) *$\deg a = 2 \Leftrightarrow m_{\max}(a) = 1$.*
- (4) *$\text{comult } a = 1 \Leftrightarrow a = \varphi(e_0)$ for some Jonquières $\varphi \in W_{\infty}$.*

Proof. If $\deg a = 1$, then $a = e_0$ (Corollary 2.29), so $m_{\max}(a) = 0$ and $\text{comult } a = 1$.

If $\deg a > 1$, then $1 \leq m_{\max}(a) \leq \deg a - 1$ follows from Noether equalities and positivity of multiplicities, see Corollary 2.29. This yields $1 \leq \text{comult } a \leq \deg a - 1$. Moreover, we have $m_{\max}(a) = 1$ if and only if $\deg a = 2$ (again by Corollary 2.29), and $\text{comult } a = 1$ if and only if $a = \varphi(e_0)$, where φ is Jonquières (Lemma 2.37(2)). \square

We will often apply quadratic maps in the sequel, and need the following basic lemma.

Lemma 3.3. *Let $p_1, p_2, p_3 \in \mathcal{B}(\mathbb{P}^2)$ be three distinct points, let $\sigma_{p_1, p_2, p_3} \in W_{\infty}$ be as in Definition 2.35 and let $a \in \mathcal{Z}_{\mathbb{P}^2}$. The following hold:*

- (1) *$\sigma_{p_1, p_2, p_3}(a) = a \Leftrightarrow \deg \sigma_{p_1, p_2, p_3}(a) = \deg a \Leftrightarrow \deg(a) = m_{p_1}(a) + m_{p_2}(a) + m_{p_3}(a)$;*
- (2) *$\deg(\sigma_{p_1, p_2, p_3}(a)) < \deg(a) \Leftrightarrow \deg(a) < m_{p_1}(a) + m_{p_2}(a) + m_{p_3}(a)$;*
- (3) *$\deg(\sigma_{p_1, p_2, p_3}(a)) < \deg(a) \Rightarrow \text{comult}(\sigma_{p_1, p_2, p_3}(a)) \leq \text{comult}(a)$.*

Proof. Writing $\xi = e_0 - e_{p_1} - e_{p_2} - e_{p_3}$, we prove that $\sigma_{p_1, p_2, p_3}(v) = v + (\xi \cdot v) \cdot \xi$ for each $v \in \mathcal{Z}_{\mathbb{P}^2}$. As $v \mapsto (\xi \cdot v) \cdot \xi$ is \mathbb{Z} -linear, it suffices to check this for $v = e_0$ and $v = e_q$, $q \in \mathcal{B}(\mathbb{P}^2)$, and this follows directly from the definition given in Definition 2.35. We find

$$\deg \sigma_{p_1, p_2, p_3}(a) = e_0 \cdot (a + (\xi \cdot a) \cdot \xi) = \deg(a) + \xi \cdot a = 2 \deg(a) - m_{p_1}(a) - m_{p_2}(a) - m_{p_3}(a).$$

Hence, $\deg \sigma_{p_1, p_2, p_3}(a) = \deg a$ if and only if $a \cdot \xi = 0$, which is equivalent to $\sigma_{p_1, p_2, p_3}(a) = a$. This yields (1). Assertion (2) also follows from the above equalities. To prove (3), we write $b = \sigma_{p_1, p_2, p_3}(a) = a - n\xi$ where $n = -(\xi \cdot a) > 0$ and choose a point $q \in \mathcal{B}(\mathbb{P}^2)$ where a has maximal multiplicity. We have $\text{comult}(a) = \deg(a) - m_q(a) = a \cdot (e_0 - e_q) = b \cdot (e_0 - e_q) + \lambda(\xi \cdot (e_0 - e_q))$. As $b \cdot (e_0 - e_q) = \deg(b) - m_q(b) \geq \text{comult}(b)$, it suffices to observe that $\xi \cdot (e_0 - e_q) \in \{0, 1\}$. \square

Corollary 3.4. *If $p_1, p_2, p_3 \in \mathcal{B}(\mathbb{P}^2)$ are three distinct points and $a \in \mathcal{Z}_{\mathbb{P}^2}$ satisfies $\deg(a) = m_{p_1}(a) + m_{p_2}(a) + m_{p_3}(a)$, then $\iota_{p_1, \{p_2, p_3\}}(a) = \tau(a)$, where $\tau \in \text{Sym}_{\mathbb{P}^2}$ is the permutation of p_2 and p_3 .*

Proof. Follows from Lemma 3.3 and from the equality $\iota_{p_1, \{p_2, p_3\}} = \tau \circ \sigma_{p_1, p_2, p_3}$. \square

The following result is quite old, and was first showed by Max Noether. We give here a proof inspired by [Alb2002, Proposition 2.6.4].

Lemma 3.5. *Let $a \in \mathcal{Z}_{\mathbb{P}^2}$ be such that $a^2 = 1$, $\omega(a) = -3$, $\deg(a) \geq 2$ and $m_q(a) \geq 0$ for each $q \in \mathcal{B}(\mathbb{P}^2)$. Then, there exist three distinct points $p_1, p_2, p_3 \in \text{Base}(a)$ such that*

$$\sum_{i=1}^3 m_{p_i}(a) > \deg(a) \quad (\text{Noether inequality}).$$

Moreover, for all p_1, p_2, p_3 as above, we have $\deg \sigma_{p_1, p_2, p_3}(a) < \deg a$.

Proof. We write $a = de_0 - \sum_{i=1}^r m_i q_i$ where $d = \deg a \geq 2$, $p_1, \dots, p_r \in \mathcal{B}(\mathbb{P}^2)$ are distinct points and $m_i = m_{q_i}(a)$ for each i , and $m_1 \geq m_2 \geq \dots \geq m_r \geq 0$. The fact that $a^2 = 1$ and $\omega(a) = -3$ yield $\sum m_i^2 = d^2 - 1$, $\sum m_i = 3(d - 1)$. This implies that $m_i < d$ for each i , and thus that $r \geq 3$ and $m_3 > 0$. We then compute

$$(d - 1)(3m_3 - (d + 1)) = m_3(\sum m_i) - \sum m_i^2 = \sum m_i(m_3 - m_i) \geq \sum_{i=1}^2 m_i(m_3 - m_i).$$

Adding $(d - 1)(m_1 + m_2 - 2m_3)$ to both sides, we get

$$(d - 1)(m_1 + m_2 + m_3 - (d + 1)) \geq (m_1 - m_3)(d - 1 - m_1) + (m_2 - m_3)(d - 1 - m_2).$$

The right hand side being non-negative, we obtain $m_1 + m_2 + m_3 > d$, as expected. The last part follows from Lemma 3.3(2). \square

Corollary 3.6. *Let $a \in W_\infty(e_0)$ be such that $\deg(a) \geq 2$ and let $p \in \mathcal{B}(\mathbb{P}^2)$ be a point of maximal multiplicity of a . Then, there exists $\varphi \in J_p$ such that $\deg(\varphi(a)) < \deg(a)$.*

Proof. By Lemma 3.5, there exist three distinct points $p_1, p_2, p_3 \in \text{Base}(a)$ such that $\sum_{i=1}^3 m_{p_i}(a) > \deg(a)$. As p is a point of maximal multiplicity, we can assume $p = p_1$. We then choose $\varphi = \sigma_{p, p_2, p_3} \in J_p$, which satisfies $\deg(\varphi(a)) < \deg(a)$ (Lemma 3.5). \square

Algorithm 3.7 (Hudson Test). Lemma 3.5 yields the following algorithm, that decides whether an element of $\mathcal{Z}_{\mathbb{P}^2}$ belongs to $W_\infty(e_0)$ or not. If a belongs to $W_\infty(e_0)$, one first needs to have $a^2 = 1$, $\omega(a) = -3$ (Noether equalities). If $\deg a = 1$, then $a = e_0 \in W_\infty(e_0)$. Otherwise, one needs to have $\deg a \geq 2$ and $m_q(a) \geq 0$ for each $q \in \mathcal{B}(\mathbb{P}^2)$ (Corollary 2.29). Then one can apply Lemma 3.5 to obtain an element $a' \in W_\infty(a)$ of smaller degree, satisfying again the Noether equalities. If $\deg a' \geq 2$ and the multiplicities are again non-negative, one again applies the corollary and decreases the degree. At some moment, either we obtain e_0 , and then $a \in W_\infty(e_0)$, or we get some negative degree or multiplicity, and then $a \notin W_\infty(e_0)$ (by Corollary 2.29).

Example 3.8. Taking 12 different points $q_1, \dots, q_{12} \in \mathcal{B}(\mathbb{P}^2)$.

The element $a = -7e_0 + \sum_{i=1}^{12} 2e_{q_i} \in \mathcal{Z}_{\mathbb{P}^2}$ satisfies the Noether equalities, but has negative degree (and negative multiplicities), hence does not belong to $W_\infty(e_0)$.

The element $a' = 3e_0 + e_{q_1} - \sum_{i=4}^{11} e_{q_i}$ satisfies the Noether equalities, has positive degree but has one negative multiplicity, hence does not belong to $W_\infty(e_0)$.

The element $a'' = \sigma_{q_1, q_2, q_3}(a') = 7e_0 - 3e_{q_1} - 4e_{q_2} - 4e_{q_3} - \sum_{i=4}^{11} e_{q_i}$ satisfies the Noether equalities, has positive degree and non-negative multiplicities but does not belong to $W_\infty(e_0)$, as a' does not.

3.2. Predecessors.

Lemma 3.9. *Let $a \in W_\infty(e_0)$, $g \in W_\infty$, and $p_1, p_2 \in \mathcal{B}(\mathbb{P}^2)$ be two distinct points. Denote by $\tau \in \text{Sym}_{\mathbb{P}^2}$ the transposition that exchanges p_1 and p_2 . The comparison of the two elements $b = g(a)$ and $c = \tau \circ g \circ \tau(a)$ of $W_\infty(e_0)$ is given as follows:*

$$(\clubsuit) \quad \deg(c) - \deg(b) = (m_{p_1}(a) - m_{p_2}(a))(m_{p_1}(g) - m_{p_2}(g)).$$

Moreover, the following hold:

- (1) $\deg(b) = \deg(c) \Leftrightarrow b \equiv_{\text{Sym}_{\mathbb{P}^2}} c$;
- (2) $\deg(b) > \deg(c) \Rightarrow \text{comult}(b) \geq \text{comult}(c)$;
- (3) $\deg(b) < \deg(c) \Rightarrow \text{comult}(b) \leq \text{comult}(c)$.

Proof. For all $\alpha \in \mathcal{Z}_{\mathbb{P}^2}$ we have $\tau(\alpha) - \alpha = (m_{p_1}(\alpha) - m_{p_2}(\alpha))(e_{p_1} - e_{p_2})$. This yields

$$(\spadesuit) \quad (\tau(\alpha) - \alpha) \cdot \beta = (m_{p_1}(\alpha) - m_{p_2}(\alpha))(m_{p_1}(\beta) - m_{p_2}(\beta)) \text{ for all } \alpha, \beta \in \mathcal{Z}_{\mathbb{P}^2}.$$

We then write $\Lambda = g^{-1}(e_0)$, and obtain $m_{p_i}(\Lambda) = m_{p_i}(g)$ for $i = 1, 2$. As $\deg(c) = \deg(g \circ \tau(a))$, we get

$$\begin{aligned} \deg(c) - \deg(b) &= (g \circ \tau(a) - g(a)) \cdot e_0 = (\tau(a) - a) \cdot g^{-1}(e_0) = (\tau(a) - a) \cdot \Lambda \\ &\stackrel{\heartsuit}{=} (m_{p_1}(a) - m_{p_2}(a))(m_{p_1}(\Lambda) - m_{p_2}(\Lambda)) \\ &= (m_{p_1}(a) - m_{p_2}(a))(m_{p_1}(g) - m_{p_2}(g)), \end{aligned}$$

which achieves the proof of (\clubsuit) .

(1): If $b \equiv_{\text{Sym}_{\mathbb{P}^2}} c$, then $c \cdot e_0 = b \cdot e_0$, as e_0 is fixed by $\text{Sym}_{\mathbb{P}^2}$, i.e. $\deg(c) = \deg(b)$. Conversely, we suppose that $\deg(c) = \deg(b)$, which implies that $m_{p_1}(a) = m_{p_2}(a)$ or $m_{p_1}(g) = m_{p_2}(g)$ (by (\clubsuit)), and we want to prove that $b \equiv_{\text{Sym}_{\mathbb{P}^2}} c$. If $m_{p_1}(a) = m_{p_2}(a)$, then $\tau(a) = a$, which yields $c = \tau \circ g \circ \tau(a) = \tau(g(a)) = \tau(b)$. If $m_{p_1}(g) = m_{p_2}(g)$, then $\tau(\Lambda) = \Lambda$, i.e. $(g \circ \tau)^{-1}(e_0) = g^{-1}(e_0)$. There exists thus $\beta \in \text{Sym}_{\mathbb{P}^2}$ such that $\beta \circ g = g \circ \tau$ (Corollary 2.21). This yields $c = \tau \circ g \circ \tau(a) = \tau \circ \beta \circ g(a) = (\tau \circ \beta)(b) \in W_\infty(c)$.

(2): Assume that $\deg(b) > \deg(c)$. Up to exchanging p_1 and p_2 , we can assume that $m_{p_1}(a) > m_{p_2}(a)$ and $m_{p_1}(g) < m_{p_2}(g)$ (by (\clubsuit)). To show that $\text{comult } b = \text{comult } g(a) \geq \text{comult } \tau \circ g \circ \tau(a) = \text{comult } c$, we denote by $q \in \mathcal{B}(\mathbb{P}^2)$ a point of maximal multiplicity of b and write $\Gamma = g^{-1}(e_0 - e_q)$. This yields

$$\begin{aligned} \text{comult } b &= (e_0 - e_q) \cdot g(a) = g^{-1}(e_0 - e_q) \cdot a = \Gamma \cdot a, \\ \text{comult } c &\leq (e_0 - \tau(e_q)) \cdot c = (\tau \circ g \circ \tau)^{-1}(e_0 - \tau(e_q)) \cdot a = \tau(\Gamma) \cdot a. \end{aligned}$$

We then have $\text{comult } c - \text{comult } b \leq (\tau(\Gamma) - \Gamma) \cdot a \stackrel{\heartsuit}{=} (m_{p_1}(\Gamma) - m_{p_2}(\Gamma)) \cdot (m_{p_1}(a) - m_{p_2}(a))$, hence to prove the inequality $\text{comult } b \geq \text{comult } c$, it remains to see that $m_{p_1}(\Gamma) - m_{p_2}(\Gamma) > 0$ is impossible (as $m_{p_1}(a) > m_{p_2}(a)$). Indeed, this would yield (as $m_{p_1}(\Lambda) < m_{p_2}(\Lambda)$)

$$(\tau(\Gamma) - \Gamma) \cdot \Lambda \stackrel{\heartsuit}{=} (m_{p_1}(\Gamma) - m_{p_2}(\Gamma)) \cdot (m_{p_1}(\Lambda) - m_{p_2}(\Lambda)) < 0,$$

which implies that $\tau(\Gamma) \cdot \Lambda < \Gamma \cdot \Lambda = g^{-1}(e_0 - e_q) \cdot g^{-1}(e_0) = (e_0 - e_q) \cdot e_0 = 1$. This is impossible as $\tau(\Gamma) \in W_\infty(e_0 - e_q)$ and $\Lambda \in W_\infty(e_0)$ (see Corollary 2.30(2)).

Assertion (3) follows from (2) by replacing g with $\tau \circ g \circ \tau$, which exchanges b and c . \square

Definition 3.10. Let $a \in W_\infty(e_0)$. A *predecessor* of a is an element of

$$\{\varphi(a) \mid \varphi \in W_\infty \text{ is a Jonquières element}\}$$

of minimal degree.

If $a \in W_\infty(e_0)$ has degree at least 2 (i.e. $a \neq e_0$, see Corollary 2.29), it follows from Corollary 3.6 that the predecessors of a have degree smaller than a . The following fundamental lemma establishes (among others) the uniqueness of a predecessor modulo $\text{Sym}_{\mathbb{P}^2}$, and gives an explicit way to compute a predecessor.

Lemma 3.11. *Let $a \in W_\infty(e_0)$ be an element of degree $d > 1$. Denote by $(d; m_0, \dots, m_r)$ its homaloidal type, where we may assume that $m_0 \geq \dots \geq m_r \geq 1$. Setting $m_i = 0$ for $i > r$, we obtain an infinite non-increasing sequence $(m_i)_{i \geq 0}$. We will say that an ordering p_0, \dots, p_r of the base-points of a is non-increasing if the (finite) sequence of multiplicities $i \mapsto m_{p_i}(a)$ is non-increasing. Equivalently, this means that $m_{p_i}(a) = m_i$ for $0 \leq i \leq r$. Then, the following assertions are satisfied:*

- (1) The set $\mathcal{S} = \{s \geq 1 \mid m_0 + m_{2s-1} + m_{2s} \geq d \geq m_0 + m_{2s+1} + m_{2s+2}\}$ is a non-empty subset of consecutive integers of the interval $[1; \frac{r}{2}] \subseteq \mathbb{R}$ (whence $r \geq 2$).
- (2) All predecessors of a are equal modulo $\text{Sym}_{\mathbb{P}^2}$.
- (3) Choose any non-increasing ordering p_0, \dots, p_r of the base-points of a . Then, for each integer $s \in [1; \frac{r}{2}]$ and each $\alpha \in \text{Sym}_{\mathbb{P}^2}$, the element $\alpha \circ \iota_{p_0, \{p_1, \dots, p_{2s}\}}(a)$ is a predecessor of a if and only if $s \in \mathcal{S}$.
- (4) If $\varphi \in W_\infty$ is a Jonquière's element such that $\varphi(a)$ is a predecessor of a , then φ is equal to $\alpha \circ \iota_{p_0, \{p_1, \dots, p_{2s}\}}$ for some choice of a non-increasing ordering p_0, \dots, p_r of the base-points of a , and for some $\alpha \in \text{Sym}_{\mathbb{P}^2}$, $s \in \mathcal{S}$. In particular, we have $\text{Base}(\varphi) \subseteq \text{Base}(a)$.
- (5) If $\varphi \in W_\infty$ is a Jonquière's element, and c is a predecessor of a , then $\text{comult}(\varphi(a)) \geq \text{comult}(c)$. In particular, we have $\text{comult}(a) \geq \text{comult}(c)$.

Proof. We prove three assertions:

(I): Proof of (1). The inequality $r \geq 2$ follows from Lemma 3.5. We now show that the non-increasing sequence $i \mapsto u_i$ defined by $u_i := m_0 + m_{2i-1} + m_{2i}$ for $i \geq 1$ satisfies

$$\text{(Ia)} : u_1 > d, \text{ and } \text{(Ib)} : u_i < d \text{ for } i > r/2.$$

(Ia) : The inequality $u_1 = m_0 + m_1 + m_2 > d$ is Noether inequality (Lemma 3.5).

(Ib) : The inequality $2i > r$ yields $m_{2i} = 0$, which gives $u_i = m_0 + m_{2i-1} \leq m_0 + m_{2i-2} \leq \dots \leq m_0 + m_1 \leq d$ (Corollary 2.29). It remains to observe that $m_1 = m_2 = \dots = m_{2i-1} = d - m_0$ and $r = 2i - 1$ is impossible (Lemma 2.46). Therefore, (Ia) and (Ib) are proven. These assertions imply (1) because of the equality $\mathcal{S} = \{s \geq 1 \mid u_s \geq d \geq u_{s+1}\}$.

(II): For any non-increasing ordering p_0, \dots, p_r of the base-points of a , for any $\alpha \in \text{Sym}_{\mathbb{P}^2}$, and for any $s \in \mathcal{S}$, the elements $\alpha \circ \iota_{p_0, \{p_1, \dots, p_{2s}\}}(a)$ are all equal modulo $\text{Sym}_{\mathbb{P}^2}$.

Firstly, we fix a non-increasing ordering p_0, \dots, p_r of the base-points of a . Define $\iota_s = \iota_{p_0, \{p_1, \dots, p_{2s}\}}$ and $c_s = \iota_s(a)$ for each integer $s \in [1; \frac{r}{2}]$, and show that $c_s \equiv_{\text{Sym}_{\mathbb{P}^2}} c_{s'}$ for all $s, s' \in \mathcal{S}$. By (1), it suffices to prove this in the case where $s' = s + 1$. The fact that $s, s + 1 \in \mathcal{S}$ implies that $d = u_{s+1} = m_0 + m_{2s+1} + m_{2s+2}$, which means that $(e_0 - e_{p_0} - e_{p_{2s+1}} - e_{p_{2s+2}}) \cdot a = 0$ and implies that $\iota_{p_0, \{p_{2s+1}, p_{2s+2}\}}(a) = \tau(a)$ where $\tau \in \text{Sym}_{\mathbb{P}^2}$ is the permutation of p_{2s+1} and p_{2s+2} (Corollary 3.4). We moreover observe that $\iota_s = \iota_{p_0, \{p_1, \dots, p_{2s}\}}$ fixes p_{2s+1} and p_{2s+2} , and thus commutes with τ . This yields $c_{s'} = \iota_{s+1}(a) = \iota_s(\iota_{p_0, \{p_{2s+1}, p_{2s+2}\}}(a)) = \iota_s(\tau(a)) = \tau(\iota_s(a)) = \tau(c_s)$ as desired.

Secondly, we observe that the class of $\iota_s(a)$ does not depend on the non-increasing ordering p_0, \dots, p_r of the base-points of a . Indeed, two different orderings only differ by some product of transpositions of two points having the same multiplicity. The result then follows from Lemma 3.9.

(III): For each Jonquière's element $\varphi \in W_\infty$, one of the following holds:

- (A)** $\varphi = \alpha \circ \iota_{p_0, \{p_1, \dots, p_{2s}\}}$ for some non-increasing ordering p_0, \dots, p_r of the base-points of a , for some $\alpha \in \text{Sym}_{\mathbb{P}^2}$, and some $s \in \mathcal{S}$.
- (B)** There exists a Jonquière's element $\varphi' \in W_\infty$ such that $\deg(\varphi'(a)) < \deg(\varphi(a))$ and $\text{comult}(\varphi'(a)) \leq \text{comult}(\varphi(a))$.

Moreover, for any non-increasing ordering p_0, \dots, p_r of the base-points of a , for any $\alpha \in \text{Sym}_{\mathbb{P}^2}$, and any integer $s \in [1; \frac{r}{2}] \setminus \mathcal{S}$, the assertion (B) is satisfied.

We fix a Jonquière element $\varphi \in W_\infty$ and choose $l+1$ distinct points $p_0, \dots, p_l \in \mathcal{B}(\mathbb{P}^2)$ such that $\{p_0, \dots, p_l\} = \text{Base}(\varphi) \cup \text{Base}(a)$ (whence $l \geq r$).

Suppose that there exist $i, j \in \{0, \dots, l\}$ such that $(m_{p_i}(a) - m_{p_j}(a))(m_{p_i}(\varphi) - m_{p_j}(\varphi)) < 0$. We denote by $\tau \in \text{Sym}_{\mathbb{P}^2}$ the permutation of p_i and p_j , write $\varphi' = \tau \circ \varphi \circ \tau$ (which is again a Jonquière element, by Lemma 2.41), and get $\deg(\varphi'(a)) - \deg(\varphi(a)) \stackrel{\text{Lemma 3.9.}\clubsuit}{=} (m_{p_i}(a) - m_{p_j}(a))(m_{p_i}(\varphi) - m_{p_j}(\varphi)) < 0$. Moreover, Lemma 3.9(3) yields $\text{comult}(\varphi'(a)) \leq \text{comult}(\varphi(a))$. We are then in case (B).

We can therefore assume, after reordering the points p_i , that

$$m_{p_0}(a) \geq m_{p_1}(a) \geq \dots \geq m_{p_l}(a), \quad m_{p_0}(\varphi) \geq m_{p_1}(\varphi) \geq \dots \geq m_{p_l}(\varphi).$$

In particular, $\text{Base}(a) = \{p_k \mid k \in \{1, \dots, l\} \text{ and } m_{p_k}(a) > 0\} = \{p_0, \dots, p_r\}$, and the base-points p_0, \dots, p_r of a are given in non-increasing order. Moreover, φ has maximal multiplicity at p_0 . By Corollary 3.2(2), this implies that $1 \leq m_{p_0}(\varphi) \leq \deg(\varphi) - 1$. Since φ is a Jonquière element of W_∞ , it has a point of multiplicity $\deg(\varphi) - 1$, so that we have $m_{p_0}(\varphi) = \deg(\varphi) - 1$. There exists thus $\alpha \in \text{Sym}_{\mathbb{P}^2}$ such that $\varphi = \alpha \circ \iota_{p_0, \Delta}$, where $\Delta = \text{Base}(\varphi) \setminus \{p_0\} = \{p_k \mid k \in \{1, \dots, l\} \text{ and } m_{p_k}(\varphi) > 0\}$ has even cardinality (Lemma 2.37(2)). It follows that the set Δ is equal to $\{p_1, \dots, p_{2s}\}$ for some $s \geq 1$. If $s \in \mathcal{S}$, we are in case (A).

We now assume that $s \notin \mathcal{S}$, and show that we are in case (B), with $\varphi' = \varphi_{p_0, \{p_1, \dots, p_{2s'}\}}$, for some $s' \in \{s \pm 1\}$. We then only need to show that $\deg(c_{s'}) < \deg(c_s)$ and $\text{comult}(c_{s'}) \leq \text{comult}(c_s)$, where $c_s, c_{s'}$ are defined as in the proof of (II). As $s \notin \mathcal{S}$, we have either $u_s < d$ or $u_{s+1} > d$ (where u is the sequence defined above, in the proof of (I)).

If $u_s < d$, then $s > 1$ by (Ia), and we choose $s' = s - 1 \geq 1$. As $c_{s'} = \iota_{p_0, \{p_{2s-1}, p_{2s}\}}(c_s)$ is equal to $\sigma_{p_0, p_{2s-1}, p_{2s}}(c_s)$ modulo $\text{Sym}_{\mathbb{P}^2}$ (Definition 2.35), it suffices to prove that $(e_0 - e_{p_0} - e_{p_{2s-1}} - e_{p_{2s}}) \cdot c_s < 0$ (Lemma 3.3), which follows from

$$\begin{aligned} (e_0 - e_{p_0} - e_{p_{2s-1}} - e_{p_{2s}}) \cdot c_s &= \iota_s(e_0 - e_{p_0} - e_{p_{2s-1}} - e_{p_{2s}}) \cdot a \\ &= -(e_0 - e_{p_0} - e_{p_{2s-1}} - e_{p_{2s}}) \cdot a = u_s - d. \end{aligned}$$

If $u_{s+1} > d$, we choose $s' = s + 1$, which belongs to $[1; \frac{r}{2}] \subseteq [1; \frac{l}{2}]$ by (Ib). As before, it suffices to check that $(e_0 - e_{p_0} - e_{p_{2s+1}} - e_{p_{2s+2}}) \cdot c_s < 0$, which follows from

$$\begin{aligned} (e_0 - e_{p_0} - e_{p_{2s+1}} - e_{p_{2s+2}}) \cdot c_s &= \iota_s(e_0 - e_{p_0} - e_{p_{2s+1}} - e_{p_{2s+2}}) \cdot a \\ &= (e_0 - e_{p_0} - e_{p_{2s-1}} - e_{p_{2s}}) \cdot a = d - u_{s+1}. \end{aligned}$$

This achieves the proof of (III), which gives (4). Together with (II) and since a admits at least one predecessor, this also gives (2) and (3). It remains to prove (5). We do it by induction on $\deg(\varphi(a))$. The minimal case is when $\varphi(a)$ is a predecessor of a , so $\varphi(a) \equiv_{\text{Sym}_{\mathbb{P}^2}} c$ by (2), whence $\text{comult}(\varphi(a)) = \text{comult}(c)$. If $\deg(\varphi(a)) > \deg(c)$, then, by (III), there exists a Jonquière element $\varphi' \in W_\infty$ such that $\deg(\varphi(a)) > \deg(\varphi'(a))$ and $\text{comult}(\varphi(a)) \geq \text{comult}(\varphi'(a))$. Since we have $\text{comult}(\varphi'(a)) \geq \text{comult}(c)$ by the induction hypothesis, the result follows. \square

Corollary 3.12. *All predecessors of an element $a \in W_\infty(e_0)$ are equal modulo $\text{Sym}_{\mathbb{P}^2}$ and only depend on the class of a modulo $\text{Sym}_{\mathbb{P}^2}$.*

Proof. We first observe that the predecessors of $a \in W_\infty(e_0)$ are equal modulo $\text{Sym}_{\mathbb{P}^2}$. If $a = e_0$, this is because e_0 is the only predecessor of a ; otherwise, it follows from

Lemma 3.11(2). We then observe that the sets of predecessors of a and of $\alpha(a)$ are equal, for each $\alpha \in \text{Sym}_{\mathbb{P}^2}$. \square

Remark 3.13. By Remark 2.44, a homaloidal type may be identified with an element of $W_\infty(e_0)/\text{Sym}_{\mathbb{P}^2}$. It follows from Corollary 3.12 that for each homaloidal type $\chi \in W_\infty(e_0)/\text{Sym}_{\mathbb{P}^2}$, one can define its (unique) predecessor $\chi_1 \in W_\infty(e_0)/\text{Sym}_{\mathbb{P}^2}$ in the following way: Choose any representative $a \in W_\infty(e_0)$ of the coset χ , then χ_1 is defined as the equivalence class modulo $\text{Sym}_{\mathbb{P}^2}$ in $W_\infty(e_0)$ of any predecessor $a_1 \in W_\infty(e_0)$ of a .

Assume that $\chi = (d; m_0, \dots, m_r)$ with $d \geq 2$ and $m_0 \geq \dots \geq m_r \geq 1$. Choose an integer $s \in [1, \frac{r}{2}]$ such that $m_0 + m_{2s-1} + m_{2s} \geq d \geq m_0 + m_{2s+1} + m_{2s+2}$ (this is doable thanks to Lemma 3.11(1)). Then

$$\chi_1 = (d - \varepsilon; m_0 - \varepsilon, d - m_0 - m_1, \dots, d - m_0 - m_{2s}, m_{2s+1}, \dots, m_r)$$

where $\varepsilon = \sum_{i=1}^s (m_0 + m_{2i-1} + m_{2i} - d)$ is positive (the form of χ_1 follows from Lemma 3.11(3)). Of course, as usual, we can remove the multiplicities which are zero and order the remaining ones in a non-decreasing manner.

Example 3.14. To simplify the notation, a sequence of s multiplicities m is written m^s . In the following list, the notation $\chi \rightarrow \chi_1$ means that χ_1 is the predecessor of the homaloidal type χ .

$$\begin{aligned} (d; d-1, 1^{2d-2}) \rightarrow (1); & \quad (4; 2^3, 1^3) \rightarrow (2; 1^3); & (38; 18, 13^3, 12^4, 6) \rightarrow (23; 12, 8^3, 7^3, 6, 3); \\ (16; 6^5; 5^3) \rightarrow (12; 5^3, 4^4, 2); & (5; 2^6) \rightarrow (3; 2, 1^4); & (74; 28, 27^5, 19^2, 18) \rightarrow (58; 27, 19^6, 18, 12). \end{aligned}$$

Lemma 3.15. *Let $f, g \in W_\infty$ be elements such that $\text{Base}(f) \cap \text{Base}(g^{-1}) = \emptyset$ and $\deg(g) = d > 1$. Defining $h = f \circ g$, the following hold:*

- (1) $\deg(h) = \deg(f) \cdot \deg(g)$ and $|\text{Base}(h)| = |\text{Base}(f)| + |\text{Base}(g)|$.
- (2) *If $\varphi \in W_\infty$ is a Jonquières element such that $\varphi(g^{-1}(e_0))$ is a predecessor of $g^{-1}(e_0)$, then $\varphi(h^{-1}(e_0))$ is a predecessor of $h^{-1}(e_0)$.*

Proof. If $\deg f = 1$, we get $\text{Base}(f) = \emptyset$, $f^{-1}(e_0) = e_0$ and $h^{-1}(e_0) = g^{-1}(e_0)$, so that there is nothing to check. We can then assume that $\deg(f) = D > 1$. If $(d; m_0, \dots, m_r)$ and $(D; \mu_{r+1}, \dots, \mu_l)$ are the homaloidal types of g and f with $m_0 \geq \dots \geq m_r \geq 1$ and $\mu_{r+1} \geq \dots \geq \mu_l \geq 1$, choose orderings p_0, \dots, p_r and q_{r+1}, \dots, q_l of the base-points of g and f such that

$$g^{-1}(e_0) = de_0 - \sum_{i=0}^r m_i e_{p_i} \quad \text{and} \quad f^{-1}(e_0) = De_0 - \sum_{i=r+1}^l \mu_i e_{q_i}.$$

Since $\text{Base}(f) \cap \text{Base}(g^{-1}) = \emptyset$, there exist points $p_{r+1}, \dots, p_l \in \mathcal{B}(\mathbb{P}^2) \setminus \text{Base}(g) = \mathcal{B}(\mathbb{P}^2) \setminus \{p_0, \dots, p_r\}$ such that $g^{-1}(e_{q_i}) = e_{p_i}$ for $i = r+1, \dots, l$ (Lemma 2.23). We then obtain

$$\begin{aligned} h^{-1}(e_0) &= g^{-1}(De_0 - \sum_{i=r+1}^l \mu_i e_{q_i}) = Dg^{-1}(e_0) - \sum_{i=r+1}^l \mu_i e_{p_i} \\ &= Dde_0 - \sum_{i=0}^r Dm_i e_{p_i} - \sum_{i=r+1}^l \mu_i e_{p_i}. \end{aligned}$$

This already proves (1). It then remains to show (2). As $\varphi(g^{-1}(e_0))$ is a predecessor of $g^{-1}(e_0)$, Lemma 3.11(4) provides (up to a re-ordering of the points p_0, \dots, p_r) an element $\alpha \in \text{Sym}_{\mathbb{P}^2}$ and an integer $s \in [1, \frac{r}{2}]$ satisfying

$$(\heartsuit) \quad m_0 + m_{2s-1} + m_{2s} \geq d \geq m_0 + m_{2s+1} + m_{2s+2}$$

(where $m_i = 0$ if $i > r$) such that $\varphi = \alpha \circ \iota_{p_0, \{p_1, \dots, p_{2s}\}}$. Since $\mu_{r+1} \leq D-1$ (Corollary 2.29), we have

$$m_{p_0}(h) = m_0 D \geq \dots \geq m_{p_r}(h) = m_r D > m_{p_{r+1}}(h) = \mu_{r+1} \geq \dots \geq m_{p_l}(h) = \mu.$$

According to Lemma 3.11(3) for showing that $\varphi(h^{-1}(e_0))$ is a predecessor of $h^{-1}(e_0)$ it is sufficient (and necessary) to prove that

$$m_{p_0}(h) + m_{p_{2s-1}}(h) + m_{p_{2s}}(h) \geq Dd \geq m_{p_0}(h) + m_{p_{2s+1}}(h) + m_{p_{2s+2}}(h).$$

If $2s+2 \leq r$, this is just (♥) multiplied by D . Let us therefore now assume that $2s+2 > r$, so that we have $2s+2 \in \{r+1, r+2\}$. If $2s+2 = r+2$, it suffices to prove that $Dd \geq Dm_0 + \mu_{r+1} + \mu_{r+2}$. This follows from the inequalities $\mu_{r+1} + \mu_{r+2} \leq D$ and $m_0 \leq d-1$ (Corollary 2.29). If $2s+2 = r+1$, it suffices to prove that $Dd \geq Dm_0 + Dm_r + \mu_{r+1}$. If $m_0 + m_r \leq d-1$, this follows from $\mu_{r+1} \leq D-1$. It remains to show that the case $m_0 + m_r \geq d$ can not occur. Indeed, otherwise we would have $m_0 + m_r = d$ (Corollary 2.29) and then r should be even by Lemma 2.46. A contradiction. \square

Algorithm 3.16 (Computing the length in W_∞). To each $a_0 \in W_\infty(e_0)$ we can associate a sequence a_1, a_2, \dots , of elements of $W_\infty(e_0)$ such that a_i is a predecessor of a_{i-1} for each $i \geq 1$. We then say that a_i is a i -th predecessor of a_0 .

At some step $n \geq 0$, we have $a_n = e_0$ and then $a_j = e_0$ for all $j \geq n$. This provides then a finite sequence (a_0, \dots, a_n) ending with e_0 . Corollary 3.12 shows that this sequence is unique modulo $\text{Sym}_{\mathbb{P}^2}$, and Lemma 3.11 provides an explicit way to compute it.

The fact that the number n of steps is really the length in W_∞ will be proven in Proposition 3.19 below.

Example 3.17. As before, a sequence of s multiplicities m is written m^s . We apply Algorithm 3.16 to a list of homaloidal types:

- $(11; 6, 5, 4^2, 3^2, 2^2, 1) \rightarrow (5; 3, 2^3, 1^3) \rightarrow (2; 1^3) \rightarrow (1)$;
- $(19; 7^7, 4, 1) \rightarrow (13; 5^6, 4, 1^2) \rightarrow (8; 4, 3^5, 1^2) \rightarrow (4; 3, 1^6) \rightarrow (1)$;
- $(40; 18, 17, 14^2, 12^4, 3^2) \rightarrow (25; 12, 10^3, 8^2, 5, 3^3) \rightarrow (13; 8, 5^2, 3^6) \rightarrow (5; 2^6) \rightarrow (3; 2, 1^4) \rightarrow (1)$;
- $(38; 14^6, 11^2, 5) \rightarrow (29; 13, 11, 10^5, 5^2) \rightarrow (16; 6^5, 5^3) \rightarrow (12; 5^3, 4^4, 2) \rightarrow (7; 3^4, 2^3) \rightarrow (4; 2^3, 1^3) \rightarrow (2; 1^3) \rightarrow (1)$;
- $(184; 75, 61^6, 60, 48) \rightarrow (145; 60, 48^7, 36) \rightarrow (112; 48, 37^6, 36, 27) \rightarrow (82; 36, 27^7, 18) \rightarrow (58; 27, 19^6, 18, 12) \rightarrow (37; 18, 12^7, 6) \rightarrow (22; 12, 7^6, 6, 3) \rightarrow (10; 6, 3^7) \rightarrow (4; 3, 1^6) \rightarrow (1)$.

Example 3.18. Applying Algorithm 3.16 to $(17; 6^8)$ yields a sequence of homaloidal types of particular interest:

$$(17; 6^8) \rightarrow (14; 6, 5^6, 3) \rightarrow (8; 3^7) \rightarrow (5; 2^6) \rightarrow (3; 2, 1^4) \rightarrow (1)$$

It provides all the symmetric homaloidal types (symmetric means here that all multiplicities are the same) except the simplest one, namely $(2; 1^3)$ [Alb2002, Lemma 2.5.5]. Note that $(17; 6^8)$ and $(8; 3^7)$ are the homaloidal types of classical Bertini and Geiser involutions, associated to the blow-ups of 8, respectively 7 general points of \mathbb{P}^2 .

We can now give the following result, which is the analogue of Theorem 1 for W_∞ .

Proposition 3.19. *Let $n \geq 1$, let $a_0 \in W_\infty(e_0)$ and let $a_n \in W_\infty(e_0)$ be a n -th predecessor of a_0 . For all Jonquières elements $\psi_1, \dots, \psi_n \in W_\infty$, the element $b_n = \psi_n \circ \dots \circ \psi_1(a_0)$ satisfies*

- (1) $\deg(a_n) \leq \deg(b_n)$;
- (2) $\text{comult}(a_n) \leq \text{comult}(b_n)$;
- (3) If $\deg(a_n) = \deg(b_n)$, then a_n and b_n are equal modulo $\text{Sym}_{\mathbb{P}^2}$.

Proof. We prove the result by induction on the triples $(n, \text{lgth}(a_0), \deg(b_1))$, ordered lexicographically, where $b_1 = \psi_1(a_0)$.

If $n = 1$, (1) is given by the definition of a predecessor, and (2) and (3) follow respectively from Lemma 3.11(5) and Lemma 3.11(2). We will then assume that $n > 1$ in the sequel.

We write $\psi = \psi_n \circ \dots \circ \psi_1 \in W_\infty$, and write $\text{Base}(a_0) \cup \text{Base}(\psi) = \{p_1, \dots, p_l\}$, for some distinct points p_i that we can assume to be such that $m_{p_1}(a) \geq \dots \geq m_{p_l}(a) \geq 0$.

If there exist $i, j \in \{1, \dots, l\}$ such that $i < j$ and $m_{p_i}(\psi) < m_{p_j}(\psi)$, we denote by $\tau \in \text{Sym}_{\mathbb{P}^2}$ the permutation of p_i and p_j , write $b'_n = \tau \circ \psi \circ \tau(a_0)$ and replace b_n with b'_n and ψ_i with $\tau \circ \psi_i \circ \tau$ for $i = 1, \dots, n$, which are Jonquières elements of W_∞ (Lemma 2.41). To see that this is possible, we use

$$\deg(b'_n) - \deg(b_n) \stackrel{\text{Lemma 3.9} \spadesuit}{=} (m_{p_i}(a) - m_{p_j}(a))(m_{p_i}(\psi) - m_{p_j}(\psi)) \leq 0.$$

If $\deg(b'_n) = \deg(b_n)$ then b_n and b'_n are equal modulo $\text{Sym}_{\mathbb{P}^2}$ (Lemma 3.9(1)). And if $\deg(b'_n) < \deg(b_n)$, then $\text{comult}(b'_n) \leq \text{comult}(b_n)$ (Lemma 3.9(2)). In both cases, proving the result for b'_n gives the result for b_n . After finitely many steps, we then reduce to the case where

$$m_{p_1}(a) \geq \dots \geq m_{p_l}(a) \geq 0 \text{ and } m_{p_1}(\psi) \geq \dots \geq m_{p_l}(\psi) \geq 0.$$

In particular, both a and ψ have maximal multiplicity at p_1 .

For $i = 1, \dots, n-1$, we define a_i to be a i -th predecessor of a_0 , and write $b_i = \psi_i \circ \dots \circ \psi_1(a_0)$. If $a_n = e_0$, all assertions hold, so we can assume that $\deg(a_n) \geq 2$, which implies that $\text{comult}(a_{n-1}) \geq 2$ (by Lemma 3.2(4) the equality $\text{comult}(a_{n-1}) = 1$ would give $a_n = e_0$). By induction hypothesis, we have $\text{comult}(b_{n-1}) \geq \text{comult}(a_{n-1}) \geq 2$, so $b_n \neq e_0$. As this is true for any choice of Jonquières elements ψ_1, \dots, ψ_n , we find that

$$\text{lgth}(a_0) \geq n + 1.$$

We now apply the algorithm to $\Lambda_0 = \psi^{-1}(e_0)$. As p_1 is a point of Λ_0 of maximal multiplicity, there exists $\theta_1 \in J_{p_1}$ such that $\Lambda_1 = \theta_1(\Lambda_0)$ is a predecessor of Λ_0 (Lemma 3.11(3)). We then define a sequence $(\Lambda_i)_{i \geq 2}$, and Jonquières elements $(\theta_i)_{i \geq 2}$ such that $\Lambda_i = \theta_i(\Lambda_{i-1})$ is a predecessor of Λ_{i-1} for each $i \geq 1$. Since ψ is a product of n Jonquières elements, we have $\text{lgth}(\Lambda_0) \leq n < \text{lgth}(a_0)$. The pair $(n, \text{lgth}(a_0))$ is thus bigger than $(n, \text{lgth}(\Lambda_0))$, so we find that $\deg \Lambda_n \leq \deg \psi(\Lambda_0) = \deg e_0 = 1$ by induction hypothesis. This yields $\Lambda_n = e_0$, and thus $\theta = \theta_n \circ \dots \circ \theta_1$ satisfies $e_0 = \theta(\Lambda_0) = \theta(\psi^{-1}(e_0))$, which yields $\theta \circ \psi^{-1} \in \text{Sym}_{\mathbb{P}^2}$ (Lemma 2.20(4)). Hence, we can replace ψ_i with θ_i for $i = 1, \dots, n$, since $b_n = \psi(a_0)$ is equal to $\theta(a_0)$ modulo $\text{Sym}_{\mathbb{P}^2}$. This reduces to the case where $\psi_1 \in J_{p_1}$.

Applying induction hypothesis to b_1 and the $n-1$ Jonquières ψ_2, \dots, ψ_n , we can reduce to the case where b_i is a predecessor of b_{i-1} for $i = 2, \dots, n$. We can moreover assume that $\psi_2 \in J_q$ where $q \in \mathcal{B}(\mathbb{P}^2)$ is a base-point of b_1 of maximal multiplicity (Lemma 3.11).

If $\deg(a_1) = \deg(b_1)$, then a_1 and b_1 are equal modulo $\text{Sym}_{\mathbb{P}^2}$ (Lemma 3.11), so the result follows, as in this case b_i is a predecessor of b_{i-1} for $i = 1, \dots, n$, hence b_i is equal to a_i modulo $\text{Sym}_{\mathbb{P}^2}$ for $i = 1, \dots, n$. We can then assume that $\deg(a_1) < \deg(b_1)$.

If b_1 has maximal multiplicity at p_1 , then we can assume that $\psi_2 \in J_{p_1}$ (since $b_2 = \psi_2(b_1)$ is a predecessor of b_1), and apply the induction hypothesis to $\psi_2 \circ \psi_1, \psi_3, \dots, \psi_n$ to obtain that $\deg(a_{n-1}) \leq \deg(b_n)$ and $\text{comult}(a_{n-1}) \leq \text{comult}(b_n)$. The result follows in this case from $\deg(a_n) \leq \deg(a_{n-1})$ and $\text{comult}(a_n) \leq \text{comult}(a_{n-1})$ (Lemma 3.11(5)).

We can then find a point $q \in \mathcal{B}(\mathbb{P}^2) \setminus \{p_1\}$ such that b_1 has maximal multiplicity at q and $\psi_2 \in J_q$. We claim that there exists $r \in \mathcal{B}(\mathbb{P}^2) \setminus \{p_1, q\}$ such that $\deg(b_1) < m_{p_1}(b_1) + m_q(b_1) + m_r(b_1)$. Let us first show why this claim achieves the proof, before proving the claim. We choose the involution $\sigma_{p_1, q, r} \in J_{p_1} \cap J_q$ as in Definition 2.35, write $\psi'_1 = \sigma_{p_1, q, r} \circ \psi_1 \in J_{p_1}$ and $\psi'_2 = \psi_2 \circ \sigma_{p_1, q, r} \in J_q$. We can thus replace ψ_1 and ψ_2 with the Jonquières elements ψ'_1 and ψ'_2 respectively, without changing b_i for $i = 2, \dots, n$. This replaces b_1 with $b'_1 = \sigma_{p_1, q, r}(b_1)$, which satisfies $\deg(b'_1) < \deg(b_1)$ (Lemma 3.3(2)). The result then follows by induction hypothesis.

It remains to prove the claim. As p_1 is a point of maximal multiplicity of a_0 , there exists $\varphi_1 \in J_{p_1}$ such that $\varphi_1(a_0)$ is a predecessor of a_0 , so that $\varphi_1(a_0)$ is equal to a_1 mod $\text{Sym}_{\mathbb{P}^2}$ (Lemma 3.11). We set $\nu = \varphi_1 \circ (\psi_1)^{-1} \in J_{p_1}$ and find a set $\Delta \subseteq \mathcal{B}(\mathbb{P}^2) \setminus \{p_1\}$ of even order, such that $\nu^{-1}(e_0) = e_0 + \sum_{r \in \Delta} \frac{e_0 - e_{p_1}}{2} - e_r$ (Lemma 2.37). In particular,

$$\deg \varphi_1(a_0) = \deg \nu \circ \psi_1(a) = \nu^{-1}(e_0) \cdot b_1 = \deg b_1 + \sum_{r \in \Delta} \left(\frac{\deg b_1 - m_{p_1}(b_1)}{2} - m_r(b_1) \right).$$

Since $\deg \varphi_1(a_0) < \deg b_1$, there exist two distinct points $r_1, r_2 \in \Delta$ such that

$$\begin{aligned} 0 &> \left(\frac{\deg b_1 - m_{p_1}(b_1)}{2} - m_{r_1}(b_1) \right) + \left(\frac{\deg b_1 - m_{p_1}(b_1)}{2} - m_{r_2}(b_1) \right) \\ &= \deg b_1 - m_{p_1}(b_1) - m_{r_1}(b_1) - m_{r_2}(b_1). \end{aligned}$$

Since $m_q(b_1) \geq \max\{m_{r_1}(b_1), m_{r_2}(b_1)\}$ and $q \neq p_1$, we can replace one of the two points r_1, r_2 with q and denote by r the other one. This achieves the proof of the claim. \square

3.3. From the Weyl group to the Cremona group. Starting from an element of $\text{Bir}(\mathbb{P}^2)$, Algorithm 3.16 yields a way to decompose it into a product of Jonquières elements of W_∞ , and the optimality of the algorithm in W_∞ is given by Proposition 3.19. We now show that the algorithm also works in $\text{Bir}(\mathbb{P}^2)$. To do this, we first make the following easy observation, which relates the two notions of predecessors already defined for elements of $\text{Bir}(\mathbb{P}^2)$ and for elements of $W_\infty(e_0)$ (Definitions 1.2 and 3.10 respectively). We will then prove that the hypothesis of Lemma 3.20 is in fact always satisfied (Proposition 3.26), and this will allow us to give a stronger version of Lemma 3.20 (Corollary 3.27 below).

Lemma 3.20. *Let $f \in \text{Bir}(\mathbb{P}^2)$. If there exists a Jonquières element $\varphi \in \text{Bir}(\mathbb{P}^2)$ such that $\varphi(f^{-1}(e_0)) \in \mathcal{Z}_{\mathbb{P}^2}$ is a predecessor of $f^{-1}(e_0)$ (in the sense of Definition 3.10), then:*

- (1) $f \circ \varphi^{-1} \in \text{Bir}(\mathbb{P}^2)$ is a predecessor of f (in the sense of Definition 1.2).
- (2) $g^{-1}(e_0)$ is a predecessor of $f^{-1}(e_0)$, for each predecessor $g \in \text{Bir}(\mathbb{P}^2)$ of f .

Proof. (1): To prove that $f \circ \varphi^{-1}$ is a predecessor of f , we only need to show that $\deg(f \circ \psi^{-1}) \geq \deg(f \circ \varphi^{-1})$ (which is equivalent to $\deg(\varphi(f^{-1}(e_0))) \leq \deg(\psi(f^{-1}(e_0)))$) for each Jonquières element $\psi \in \text{Bir}(\mathbb{P}^2)$. As $\varphi(f^{-1}(e_0))$ is a predecessor of $f^{-1}(e_0)$ it satisfies $\deg(\varphi(f^{-1}(e_0))) \leq \deg(\psi(f^{-1}(e_0)))$ for each Jonquières element $\psi \in W_\infty$, and thus in particular for each Jonquières element $\psi \in \text{Bir}(\mathbb{P}^2)$ (Lemma 2.42).

(2): If $g \in \text{Bir}(\mathbb{P}^2)$ is a predecessor of f , then $g = f \circ \kappa$ for some Jonquières transformation $\kappa \in \text{Bir}(\mathbb{P}^2)$, and $\deg(g) = \deg(f \circ \varphi^{-1})$ by (1). The element $g^{-1}(e_0) = \kappa^{-1}(f^{-1}(e_0))$ has then the same degree as the predecessor $\varphi(f^{-1}(e_0))$ of $f^{-1}(e_0)$, and is thus also a predecessor of $f^{-1}(e_0)$ (by Definition 3.10). \square

We first recall the following famous result, corresponding to the algorithm defined in [Alb2002, Chapter 8], adapting the proof of Castelnuovo [Cas1901].

Proposition 3.21 (Castelnuovo reduction). *Let $f \in \text{Bir}(\mathbb{P}^2)$ be of degree $d > 1$, let $p \in \mathbb{P}^2$ be a base-point of f of maximal multiplicity. We define $M = \{q \in \text{Base}(f) \setminus \{p\} \mid m_p + 2m_q > d\}$. Then, M contains at least two elements, and the following hold:*

- (1) *If $|M|$ is even, there is an element $\varphi \in \text{Jonq}_p$ such that $\text{Base}(\varphi) = \{p\} \cup M$.*
- (2) *If $|M|$ is odd, there is $q \in M$ of minimal multiplicity and an element $\varphi \in \text{Jonq}_p$ such that $\text{Base}(\varphi) = \{p\} \cup (M \setminus \{q\})$.*

For each φ as above, we have $\deg(f \circ \varphi^{-1}) < \deg(f)$.

Remark 3.22. In [Alb2002], the elements of M are called the *major base-points* of f and their number $|M|$ is written h (see [Alb2002, Definition 8.2.1]).

Proof. The proof of the proposition lies in Chapter 8 and especially in §8.3 of [Alb2002]. The fact that $h = |M| \geq 2$ is [Alb2002, Lemma 8.2.6]. The existence of φ and the fact that $\deg(f \circ \varphi^{-1}) < \deg(f)$ is given at page 242, in the proof of [Alb2002, Theorem 8.3.4]. \square

Definition 3.23. Let $f \in \text{Bir}(\mathbb{P}^2)$. If $\deg(f) > 1$, we define a *Castelnuovo-predecessor* of f to be an element of the form $f \circ \varphi^{-1}$ where the Jonquières transformation $\varphi \in \text{Bir}(\mathbb{P}^2)$ has been described in Proposition 3.21. If $\deg(f) = 1$, we define f to be its own Castelnuovo-predecessor.

Lemma 3.24. *Let $f \in \text{Bir}(\mathbb{P}^2)$ be a Jonquières element of degree $d > 1$. Then every Castelnuovo-predecessor of f has degree 1.*

Proof. Let g be a Castelnuovo-predecessor of f , equal to $g = f \circ \varphi^{-1}$ for some Jonquières transformation $\varphi \in \text{Jonq}_p$ as in Proposition 3.21, where $p \in \mathbb{P}^2$ is a base-point of f of maximal multiplicity. We want to show that $\deg(g) = 1$. As $\deg(g) < \deg(f)$, we can assume that $d = \deg(f) > 2$. The element f is a Jonquières element of W_∞ (Lemma 2.42), and is thus equal to $\alpha \circ \iota_{q,\Delta}$, where $\alpha \in \text{Sym}_{\mathbb{P}^2}$, $q \in \mathcal{B}(\mathbb{P}^2)$ and $\Delta \subseteq \mathcal{B}(\mathbb{P}^2) \setminus \{q\}$ is a finite set of even order $2n$ (Lemma 2.41). As $f^{-1}(e_0) = \iota_{q,\Delta}(e_0) = (n+1)e_0 - ne_q - \sum_{r \in \Delta} e_r$, we get $\deg(f) = \deg(\iota_{q,\Delta}) = n+1$, whence $n \geq 2$ and q is the unique base-point of f of maximal multiplicity. It follows that $p = q$. Moreover, the set $M := \{r \in \text{Base}(f) \setminus \{q\} \mid m_q + 2m_r > d\}$ which was defined in Proposition 3.21 is equal to Δ which has even order. This implies that $\varphi \in \text{Jonq}_q$ is such that $\text{Base}(\varphi) = \{q\} \cup M = \text{Base}(f)$. We then find $\varphi = \beta \circ \iota_{q,\Delta}$ for some $\beta \in \text{Sym}_{\mathbb{P}^2}$ (Lemma 2.37). This yields $g = \alpha \circ \beta^{-1} \in \text{Sym}_{\mathbb{P}^2}$ and $\deg(g) = 1$ as desired. \square

Algorithm 3.25 (Algorithm of Castelnuovo). Taking $f_0 \in \text{Bir}(\mathbb{P}^2) \setminus \text{Aut}(\mathbb{P}^2)$, Proposition 3.21 yields a Jonquières element $\varphi_1 \in \text{Bir}(\mathbb{P}^2)$ such that the Castelnuovo predecessor $f_1 = f_0 \circ \varphi_1^{-1}$ satisfies $\deg(f_1) < \deg(f_0)$. Applying again the result finitely many times, we find a sequence of Castelnuovo predecessors f_0, f_1, \dots, f_n and a sequence of Jonquières

elements $\varphi_1, \varphi_2, \dots, \varphi_n$, which lead to a decomposition of f_0 into $f_n \circ \varphi_n \circ \dots \circ \varphi_1$, where $f_n \in \text{Aut}(\mathbb{P}^2)$. Since $\varphi'_n := f_n \circ \varphi_n$ is Jonquière, this algorithm actually provides a decomposition of f_0 into the product of n Jonquière elements

$$f_0 = \varphi'_n \circ \varphi_{n-1} \circ \dots \circ \varphi_1.$$

As we will show in Corollary 3.29, this integer n (which is the integer n for which the algorithm stops, i.e. for which $\deg f_n = 1$) is the length of f_0 .

Recall that a predecessor of an element $a \in W_\infty(e_0)$ is an element of minimal degree among all the elements of the form $\varphi(a)$ where φ is a Jonquière element of W_∞ . The next fundamental result shows that we can choose φ to be in $\text{Bir}(\mathbb{P}^2)$ if $a = f^{-1}(e_0)$ for some $f \in \text{Bir}(\mathbb{P}^2)$.

Proposition 3.26. *Let $f \in \text{Bir}(\mathbb{P}^2) \setminus \text{Aut}(\mathbb{P}^2)$ and let $p_0 \in \mathbb{P}^2$ be a base-point of f of maximal multiplicity.*

- (1) *There exists a Jonquière element $\psi \in \text{Jonq}_{p_0}$ such that $\psi(f^{-1}(e_0)) \in \mathcal{Z}_{\mathbb{P}^2}$ is a predecessor of $f^{-1}(e_0)$. For each such ψ , the element $f \circ \psi^{-1} \in \text{Bir}(\mathbb{P}^2)$ is a predecessor of f (Lemma 3.20).*
- (2) *If $\varphi \in \text{Jonq}_{p_0}$ is a Jonquière element provided by the Castelnuovo reduction of f (Proposition 3.21), one can choose ψ as above such that $\psi \circ \varphi^{-1} \in \text{Jonq}_{p'}$ for some point $p' \in \mathbb{P}^2$ of maximal multiplicity of $\varphi(f^{-1}(e_0))$.*

Proof. Let $\varphi \in \text{Jonq}_{p_0}$ be a Jonquière element provided by the Castelnuovo reduction. We choose r distinct points $p_1, \dots, p_r \in \mathcal{B}(\mathbb{P}^2)$ such that $\text{Base}(f) = \{p_0, \dots, p_r\}$, write $m_i = m_{p_i}(f)$ for each i , and choose the order such that $m_0 \geq m_1 \geq \dots \geq m_r$, and such that for any $i \geq 1$, either p_i is a proper point of \mathbb{P}^2 or p_i is in the first neighbourhood of some p_j with $j < i$. We then define $m_i = 0$ for each integer $i > r$, and write $M = \{p_i \mid i \geq 1, m_0 + 2m_i > d\}$ as in Proposition 3.21.

Suppose first that $|M|$ is even. In this case, $\varphi \in \text{Jonq}_{p_0}$ satisfies $\text{Base}(\varphi) = \{p_0\} \cup M$ (Proposition 3.21(1)) and is then equal to $\varphi = \alpha \circ \iota_{p_0, M}$ for some $\alpha \in \text{Sym}_{\mathbb{P}^2}$ (Lemma 2.37(2)). Writing $2s = |M|$, we find $m_0 + m_{2s-1} + m_{2s} = ((m_0 + 2m_{2s-1}) + (m_0 + m_{2s}))/2 > d \geq ((m_0 + 2m_{2s+1}) + (m_0 + m_{2s+2}))/2 = m_0 + m_{2s+1} + m_{2s+2}$, which implies that $\iota_{p_0, \{p_1, \dots, p_{2s}\}}(f^{-1}(e_0))$ is a predecessor of $f^{-1}(e_0)$ Lemma 3.11(3), so the same holds for $\varphi(f^{-1}(e_0))$. This achieves the proof, by choosing $\psi = \varphi$, whence $\rho = \text{id}$.

Suppose now that $|M|$ is odd. In this case, $\varphi \in \text{Jonq}_{p_0}$ satisfies $\text{Base}(\varphi) = \{p_0\} \cup (M \setminus \{q\})$ for some $q \in M$ of minimal multiplicity (Proposition 3.21(2)) and is equal to $\varphi = \alpha \circ \iota_{p_0, M \setminus \{q\}}$ for some $\alpha \in \text{Sym}_{\mathbb{P}^2}$ (Lemma 2.37(2)). Writing $2s + 1 = |M|$, we can assume that $p_{2s+1} = q$, which yields $\iota_{p_0, M \setminus \{q\}} = \iota_{p_0, \{p_1, \dots, p_{2s}\}}$, and find $m_0 + m_{2s-1} + m_{2s} = ((m_0 + 2m_{2s-1}) + (m_0 + m_{2s}))/2 > d$. We then obtain two cases:

If $d \geq m_0 + m_{2s+1} + m_{2s+2}$, then $\iota_{p_0, \{p_1, \dots, p_{2s}\}}(f^{-1}(e_0))$ is a predecessor of $f^{-1}(e_0)$ (Lemma 3.11(3)), so the same holds for $\varphi(f^{-1}(e_0))$. We then choose $\psi = \varphi$ as before.

The last case is when $d < m_0 + m_{2s+1} + m_{2s+2}$, which implies that $m_{2s+2} > 0$ (Corollary 2.29) and thus that $2s + 2 \leq r$. We can assume that p_{2s+2} is not infinitely near to a point p_i with $i > 2s + 2$ (otherwise we have $m_i = m_{2s+2}$, so we exchange p_{2s+2} with p_i). Since $d \geq m_0 + m_{2s+3} + m_{2s+4}$, the element $a_{s+1} = \iota_{p_0, \{p_1, \dots, p_{2s+2}\}}(f^{-1}(e_0))$ is a predecessor of $f^{-1}(e_0)$ (Lemma 3.11(3)). Moreover, $a_s = \iota_{p_0, \{p_1, \dots, p_{2s}\}}(f^{-1}(e_0))$ is not

a predecessor of $f^{-1}(e_0)$ (Lemma 3.11(3)), which implies that $\deg(a_s) > \deg(a_{s+1})$. We will prove the following numerical assertions:

- (I) The point p_{2s+2} is a base-point of a_{s+1} of maximal multiplicity;
- (II) $m_{p_{2s+2}}(a_s) > m_{p_i}(a_s)$ for $i = 1, \dots, 2s$;
- (III) $(e_0 - e_{p_0} - e_{p_{2s+1}} - e_{p_{2s+2}}) \cdot a_s = \deg(a_s) - m_{p_0}(a_s) - m_{p_{2s+1}}(a_s) - m_{p_{2s+2}}(a_s) < 0$;
- (IV) The point p_{2s+1} is the unique base-point of a_s of maximal multiplicity.

Before proving these assertions, let us show how they imply the result.

We write $\Lambda = \varphi(f^{-1}(e_0))$, which corresponds to the linear system of $f \circ \varphi^{-1}$. For $i = 1, \dots, r$, we then denote by $q_i \in \mathcal{B}(\mathbb{P}^2)$ the point such that $\alpha(e_{p_i}) = e_{q_i}$. As $\varphi = \alpha \circ \iota_{p_0, \{p_1, \dots, p_{2s}\}}$, we find $\text{Base}(\Lambda) \subseteq \{q_0, \dots, q_r\}$. Moreover, φ and $\iota_{p_0, \{p_1, \dots, p_{2s}\}}$ belong to J_{p_0} , so that α also belongs to J_{p_0} . This gives us $\alpha(e_0 - e_{p_0}) = e_0 - e_{p_0}$ and finally $q_0 = p_0$ is a proper point of \mathbb{P}^2 . Assertion (IV) implies that q_{2s+1} is the unique base-point of Λ of maximal multiplicity, so $p' := q_{2s+1}$ is a proper point of \mathbb{P}^2 . We then observe that q_{2s+2} is either a proper point of \mathbb{P}^2 or a point infinitely near p_0 or p' . Indeed, it cannot be infinitely near q_i if $i \in \{1, \dots, 2s\}$ by (II) and if $i > 2s + 1$, because p_{2s+1} is not infinitely near p_i . Moreover, p_0, p', q_{2s+2} are not collinear because of (III) and Bézout Theorem. Up to change of coordinates, we can thus assume that $\{p_0, p'\} = \{[1 : 0 : 0], [0 : 1 : 0]\}$ and that p_0, p', q_{2s+2} are the three base-points of a quadratic involution $\rho \in \text{Jonq}_{p_0} \cap \text{Jonq}_{p'} \subseteq \text{Bir}(\mathbb{P}^2)$ which is one of the two following

$$[x : y : z] \dashrightarrow [yz : xz : xy] \text{ or } [x : y : z] \dashrightarrow [z^2 : xy : xz],$$

and satisfies then $\rho = \beta \circ \sigma_{p_0, p', q_{2s+2}}$ for some $\beta \in \text{Sym}_{\mathbb{P}^2}$ (see Definition 2.35). The result then follows by setting $\psi := \rho \circ \varphi \in \text{Jonq}_{p_0}$. Indeed, as $\text{Base}(\beta \circ \sigma_{p_0, p', q_{2s+2}} \circ \alpha) = \{p_0, p_{2s+1}, p_{2s+2}\}$, we have $\beta \circ \sigma_{p_0, p', q_{2s+2}} \circ \alpha = \gamma \circ \iota_{p_0, \{p_{2s+1}, p_{2s+2}\}}$ for some $\gamma \in \text{Sym}_{\mathbb{P}^2}$, which yields $\psi = \beta \circ \sigma_{p_0, p', q_{2s+2}} \circ \alpha \circ \iota_{p_0, \{p_1, \dots, p_{2s}\}} = \gamma \circ \iota_{p_0, \{p_1, \dots, p_{2s+2}\}}$, and implies that $\psi(f^{-1}(e_0)) = \gamma(a_{s+1})$ is a predecessor of $f^{-1}(e_0)$. Moreover, $p' = q_{2s+1} \in \mathbb{P}^2$ is such that $\rho \in \text{Jonq}_{p'}$ and is a point of maximal multiplicity of $\psi(f^{-1}(e_0))$; this follows from (I) and from $\gamma(e_{p_{2s+2}}) = e_{p'}$, which is given by $e_0 - e_{p'} = \rho(e_0 - e_{p'}) = \rho \circ \alpha(e_0 - e_{p_{2s+1}}) = \gamma \circ \iota_{p_0, \{p_{2s+1}, p_{2s+2}\}}(e_0 - e_{p_{2s+1}}) = \gamma(e_0 - e_{p_{2s+2}}) = e_0 - \gamma(e_{p_{2s+2}})$.

It remains to prove the assertions (I)–(IV).

(I): Writing $\mu = \frac{d-m_0}{2}$, we have $m_0 \geq \dots \geq m_{2s} \geq m_{2s+1} > \mu \geq m_{2s+2} \geq \dots \geq m_r$ and $2\mu = d - m_0 < m_{2s+1} + m_{2s+2}$. This yields $m_{p_{2s+2}}(a_{s+1}) = d - m_0 - m_{2s+2} = 2\mu - m_{2s+2} \geq \mu > 2\mu - m_i = d - m_0 - m_i = m_{p_i}(a_{s+1})$ for each $i \in \{1, \dots, 2s+1\}$. We moreover have $m_{p_{2s+2}}(a_{s+1}) \geq \mu \geq m_i = m_{p_i}(a_{s+1})$ for each $i \in \{2s+3, \dots, r\}$. It then remains to show that $m_{p_{2s+2}}(a_{s+1}) > m_{p_0}(a_{s+1})$. This holds, because otherwise p_0 would be a point of maximal multiplicity of a_{s+1} , which would yield the existence of $\kappa \in J_{p_0}$ such that $\deg(\kappa(a)) < \deg(a)$ (Corollary 3.6), contradicting the fact that a_{s+1} is a predecessor of $f^{-1}(e_0)$.

(II): Follows from $m_{p_{2s+2}}(a_s) = m_{2s+2} > d - m_0 - m_{2s+1} \geq d - m_0 - m_i = m_{p_i}(a_s)$, for $i \in \{1, \dots, 2s\}$.

(III)–(IV): Set $\nu = m_{p_0}(a_s) + m_{p_{2s+1}}(a_s) + m_{p_{2s+2}}(a_s) - \deg(a_s)$. The equality $a_{s+1} = \iota_{p_0, \{p_{2s+1}, p_{2s+2}\}}(a_s)$ gives $\deg(a_{s+1}) = \deg(a_s) - \nu$, whence $\nu > 0$, i.e. (III). It also provides

$$m_{p_0}(a_s) = m_{p_0}(a_{s+1}) + \nu, m_{p_{2s+1}}(a_s) = m_{p_{2s+2}}(a_{s+1}) + \nu, m_{p_{2s+2}}(a_s) = m_{p_{2s+1}}(a_{s+1}) + \nu,$$

and $m_{p_i}(a_s) = m_{p_i}(a_{s+1})$ for $i \neq 0, 2s+1, 2s+2$. Since p_{2s+2} was a base-point of maximal multiplicity of a_s (by (I)), it follows that p_{2s+1} is a base-point of maximal multiplicity of a_{s+1} and that such base-points of maximal multiplicity of a_{s+1} are included into the set $\{p_0, p_{2s+1}, p_{2s+2}\}$. However, we have already seen in the proof of (I) that $m_{p_{2s+2}}(a_{s+1}) > m_{p_i}(a_{s+1})$ for $i = 0$ or $i = 2s+1$. This proves (IV). \square

Corollary 3.27. *Let $f, g \in \text{Bir}(\mathbb{P}^2)$. Then, the two following assertions are equivalent:*

- (1) g is a predecessor of f (Definition 1.2);
- (2) $g^{-1}(e_0)$ is a predecessor of $f^{-1}(e_0)$ (Definition 3.10) and $f^{-1} \circ g \in \text{Jonq} \subseteq \text{Bir}(\mathbb{P}^2)$.

Proof. By Proposition 3.26(1), the assumptions of Lemma 3.20 are always fulfilled. Therefore, the implication (1) \Rightarrow (2) follows from Lemma 3.20(2) and the implication (2) \Rightarrow (1) follows from Lemma 3.20(1) applied with $\varphi = g^{-1} \circ f \in \text{Jonq}$. \square

We are now ready to give the proof of Lemma 1.3 and Theorem 1.

Proof of Lemma 1.3. Let $f \in \text{Bir}(\mathbb{P}^2)$, and let $g \in \text{Bir}(\mathbb{P}^2)$ be a predecessor of f , which is then equal to $g = f \circ \varphi$ for some Jonquières element $\varphi \in \text{Bir}(\mathbb{P}^2)$.

By Corollary 3.27, $g^{-1}(e_0) = \varphi^{-1}(f^{-1}(e_0))$ is a predecessor of $f^{-1}(e_0)$. This implies that $\text{Base}(\varphi^{-1}) \subseteq \text{Base}(f^{-1}(e_0)) = \text{Base}(f)$ (Lemma 3.11(4)), and gives (3). Moreover, the homaloidal type of g , which is the class of $g^{-1}(e_0)$ modulo $\text{Sym}_{\mathbb{P}^2}$, is uniquely determined by the homaloidal type of f (Corollary 3.12). This proves (1).

It remains to prove (2). The set of predecessors being invariant under right multiplication by elements of $\text{Aut}(\mathbb{P}^2)$, it is infinite. It remains to see that the number of classes modulo $\text{Aut}(\mathbb{P}^2)$ is finite. This corresponds to saying that the number of possibilities for $\text{Base}(\varphi^{-1})$ is finite, and is thus given by (3). \square

Proof of Theorem 1. For each $i \geq 0$, we set $a_i := (f_i)^{-1}(e_0) \in W_\infty(e_0)$. By Corollary 3.27, a_i is a predecessor of a_{i-1} for each $i \geq 1$, so a_i is a i -th predecessor of a_0 for each $i \geq 1$.

We then write $b_n = g_n^{-1}(e_0) = \varphi_n^{-1} \circ \dots \circ \varphi_1^{-1}(a_0)$. As $\varphi_1, \dots, \varphi_n$ are Jonquières elements of $\text{Bir}(\mathbb{P}^2)$, they are all the more Jonquières elements of W_∞ (Lemma 2.42). Then, the three assertions (1)-(2)-(3) of Theorem 1 directly follow from the three corresponding assertions (1)-(2)-(3) of Proposition 3.19 that we now recall: (1) $\deg(a_n) \leq \deg(b_n)$; (2) $\text{comult}(a_n) \leq \text{comult}(b_n)$; (3) if $\deg(a_n) = \deg(b_n)$, then a_n and b_n are equal modulo $\text{Sym}_{\mathbb{P}^2}$.

It remains to observe that $\text{lgth}(f_0) = \min\{n \mid \deg(f_n) = 1\}$. To do this, we write $\ell = \text{lgth}(f_0)$, $m = \min\{n \mid \deg(f_n) = 1\}$, and prove $\ell \geq m$ and $m \geq \ell$.

The fact that $\ell = \text{lgth}(f)$ yields the existence of Jonquières elements $\varphi_1, \dots, \varphi_\ell \in \text{Bir}(\mathbb{P}^2)$ such that $\deg(f \circ \varphi_1 \circ \dots \circ \varphi_\ell) = 1$. By (1), we find $\deg(f_\ell) \leq 1$, which yields $\ell \geq m$.

Writing for each i a Jonquières element $\psi_i \in \text{Jonq}$ such that $f_{i-1} = f_i \circ \psi_i$, we obtain $f_0 = f_m \circ \psi_m \circ \dots \circ \psi_1$. As $\deg(f_m) = 1$, this implies that $\ell = \text{lgth}(f_0) \leq m$. \square

Corollary 3.28 (Length of predecessors, associated to any point of maximal multiplicity). *Let $f \in \text{Bir}(\mathbb{P}^2) \setminus \text{Aut}(\mathbb{P}^2)$. For each point $q \in \mathbb{P}^2$ of maximal multiplicity of f , there exists $\varphi \in \text{Jonq}_q \subseteq \text{Bir}(\mathbb{P}^2)$ such that $f \circ \varphi$ is a predecessor of f . Moreover, every predecessor g of f satisfies $\text{lgth}(g) = \text{lgth}(f) - 1$ and $\deg(g) < \deg(f)$.*

Proof. The existence of φ is given by Proposition 3.26. Any predecessor g of f satisfies $\deg(g) < \deg(f)$ (Corollary 3.6). Finally, we have $\text{lgth}(g) = \text{lgth}(f) - 1$ by Theorem 1. \square

Corollary 3.29 (The Algorithm of Castelnuovo also provides the length). *Let $f \in \text{Bir}(\mathbb{P}^2) \setminus \text{Aut}(\mathbb{P}^2)$, let $g \in \text{Bir}(\mathbb{P}^2)$ be a predecessor of f , and let $h \in \text{Bir}(\mathbb{P}^2)$ be a Castelnuovo predecessor of f . Then, the following hold:*

- (1) *The predecessors of g and h have the same homaloidal type;*
- (2) $\text{lgth}(g) = \text{lgth}(h) = \text{lgth}(f) - 1$.

Hence, writing $f = f_0$ and denoting by f_1, f_2, \dots elements of $\text{Bir}(\mathbb{P}^2)$ such that f_i is a Castelnuovo-predecessor of f_{i-1} for $i \geq 1$, we find $\text{lgth}(f) = \min\{n \mid \deg(f_n) = 1\}$.


Proof. (1): By definition of a Castelnuovo predecessor there exists a proper base-point p_0 of maximal multiplicity of f and a Jonquière's transformation $\varphi \in \text{Jonq}_{p_0}$ such that $h = f \circ \varphi^{-1}$. By Proposition 3.26 there exists a Jonquière's transformation $\psi \in \text{Jonq}_{p_0}$ such that the elements $f_1 := f \circ \psi^{-1}$ and $\rho := \psi \circ \varphi^{-1}$ satisfy the following assertions:

- (1) f_1 is a predecessor of f ;
- (2) $\rho \in \text{Jonq}_{p'}$ for some point $p' \in \mathbb{P}^2$ of maximal multiplicity of $\psi(f^{-1}(e_0))$.

By Lemma 1.3, f_1 and g have the same homaloidal type. Applying Proposition 3.26 to f_1 and p' , we find $\kappa \in \text{Jonq}_{p'}$ such that $f_2 := f_1 \circ \kappa \in \text{Bir}(\mathbb{P}^2)$ is a predecessor of f_2 .

We now prove that f_2 is a predecessor of h . Firstly, $f_2 = f_1 \circ \kappa = f \circ \psi^{-1} \circ \kappa = h \circ \rho^{-1} \circ \kappa = h \circ \kappa'$, where $\kappa' = \rho^{-1} \circ \kappa \in \text{Jonq}_{p'}$. Secondly, if $\nu \in \text{Bir}(\mathbb{P}^2)$ is any Jonquière's transformation, we prove that $\deg(f_2) \leq \deg(h \circ \nu)$. This follows from Theorem 1, because $h \circ \nu = f \circ \varphi^{-1} \circ \nu$.

As f_2 is a predecessor of h and a predecessor of f_1 , which has the same homaloidal type as g , all predecessors of g and h have the same homaloidal type (Lemma 1.3). This achieves the proof of (1). The situation is as follows:



(2): As $f \notin \text{Aut}(\mathbb{P}^2)$, we have $\deg(f) > 1$. Corollary 3.28 yields $\text{lgth}(f_1) = \text{lgth}(g) = \text{lgth}(f) - 1 \geq 0$, because g and f_1 are predecessors of f . It remains to prove that $\text{lgth}(h) = \text{lgth}(f_1)$.

If f is a Jonquière's element, then $\text{lgth}(f) = 1$, whence $\text{lgth}(f_1) = 0$. As $\deg(g) = 1$ in this case (Lemma 3.24), we also have $\text{lgth}(g) = 0$.

If f is not a Jonquière's element, then $\deg(h) > 1$ and $\deg(f_1) > 1$. We find $\text{lgth}(f_2) = \text{lgth}(h) - 1 = \text{lgth}(f_1) - 1$, as f_2 is a predecessor of f_1 and h (Corollary 3.28).

The last sentence is a direct consequence of (2). \square

4. EXAMPLES AND APPLICATIONS

4.1. Length of birational maps of small degree. The next table gives all homaloidal types of degree ≤ 6 . The homaloidal types are given in the second column (as already

said before, a sequence of s multiplicities m is written m^s). In the first column ($\#$) a label is associated to each homaloidal type: This is the degree “ d ” if the type is Jonquières, or “ $d.i$ ” for the others (the order, for each degree, being the anti-lexicographic order according to the multiplicities). Then, the third column (ℓ) gives the length and the fourth (pr.) gives the predecessor (designated by its label). If the Castelnuovo-predecessor is different from the predecessor, it is also given, but in parenthesis. We see that the lengths are not directly related to the ordering of homaloidal types that we use.

#	h. type	ℓ	pr.	#	h. type	ℓ	pr.	#	h. type	ℓ	pr.
1	1	0		4.1	$2^3, 1^3$	2	2	6	$5, 1^{10}$	1	1
2	1^3	1	1	5	$4, 1^8$	1	1	6.1	$4, 2^4, 1^3$	2	2
3	$2, 1^4$	1	1	5.1	$3, 2^3, 1^3$	2	2(3)	6.2	$3^3, 2, 1^4$	2	3
4	$3, 1^6$	1	1	5.2	2^6	2	3	6.3	$3^2, 2^4, 1$	2	3

The types of degree 7, 8, 9 are given below, and provide the first types of length 3:

#	hom. type	ℓ	pr.	#	hom. type	ℓ	pr.	#	hom. type	ℓ	pr.
7	$6, 1^{12}$	1	1	8.3	$5, 3^2, 2^5$	2	3	9.2	$6, 3^4, 2, 1^4$	2	3
7.1	$5, 2^5, 1^3$	2	2(3)	8.4	$4^3, 3, 1^6$	2	4	9.3	$6, 3^3, 2^4, 1$	2	3
7.2	$4, 3^3, 1^5$	2	3(4)	8.5	$4^3, 2^3, 1^3$	3	4.1	9.4	$5, 4^3, 1^7$	2	4(5)
7.3	$4, 3^2, 2^3, 1^2$	2	3	8.6	$4^2, 3^2, 2^3, 1$	3	4.1(5.1)	9.5	$5, 4^2, 3, 2^3, 1^2$	3	4.1(5.1)
7.4	$3^4, 2^3$	3	4.1(5.1)	8.7	$4, 3^5, 1^2$	2	4	9.6	$5, 4, 3^4, 1^3$	2	4
8	$7, 1^{14}$	1	1	8.8	3^7	3	5.2	9.7	$5, 4, 3^3, 2^3$	3	4.1
8.1	$6, 2^6, 1^3$	2	2	9	$8, 1^{16}$	1	1	9.8	$4^4, 2^4$	3	5.1(6.1)
8.2	$5, 3^3, 2^2, 1^3$	2	3	9.1	$7, 2^7, 1^3$	2	2(3)	9.9	$4^3, 3^3, 2, 1$	3	5.1

There are then 17 types of degree 10 and 19 types of degree 11, each of length ≤ 3 .

#	hom. type	ℓ	pr.	#	hom. type	ℓ	pr.	#	hom. type	ℓ	pr.
10	$9, 1^{18}$	1	1	10.6	$6, 4^2, 3^2, 2^3, 1$	3	4.1	10.12	$5^2, 4, 3^3, 2, 1^2$	3	5.1
10.1	$8, 2^8, 1^3$	2	2	10.7	$6, 3^7$	2	4	10.13	$5^2, 3^5, 2$	3	5.2
10.2	$7, 3^5, 1^5$	2	3(4)	10.8	$5^3, 4, 1^8$	2	5	10.14	$5, 4^3, 3^2, 2^2$	3	5.1
10.3	$7, 3^4, 2^3, 1^2$	2	3	10.9	$5^3, 3, 2^3, 1^3$	3	5.1	10.15	$4^6, 1^3$	3	6.1
10.4	$6, 4^3, 2^3, 1^3$	3	4.1(6.1)	10.10	$5^3, 2^6$	3	5.2	10.16	$4^5, 3^2, 1$	3	6.3
10.5	$6, 4^2, 3^3, 1^4$	2	4	10.11	$5^2, 4^2, 2^4, 1$	3	5.1(6.1)				

#	hom. type	ℓ	pr.	#	hom. type	ℓ	pr.	#	hom. type	ℓ	pr.
11	$10, 1^{20}$	1	1	11.7	$7, 4, 3^6, 1$	2	4	11.14	$6, 4^3, 3^4$	3	5.1
11.1	$9, 2^9, 1^3$	2	2(3)	11.8	$6, 5^3, 1^9$	2	5(6)	11.15	$5^4, 2^5$	3	6.1(7.1)
11.2	$8, 3^5, 2^2, 1^3$	2	3	11.9	$6, 5^2, 4, 2^4, 1^2$	3	5.1(6.1)	11.16	$5^3, 4, 3^3, 1^2$	3	6.2(7.2)
11.3	$8, 3^4, 2^5$	2	3	11.10	$6, 5^2, 3^3, 2, 1^3$	3	5.1	11.17	$5^2, 4^4, 2, 1^2$	3	6.1
11.4	$7, 4^3, 3^2, 1^5$	2	4	11.11	$6, 5^2, 3^2, 2^4$	3	5.2	11.18	$5^2, 4^3, 3^2, 2$	3	6.3
11.5	$7, 4^3, 3, 2^3, 1^2$	3	4.1	11.12	$6, 5, 4^2, 3^2, 2^2, 1$	3	5.1				
11.6	$7, 4^2, 3^3, 2^3$	3	4.1(5.1)	11.13	$6, 4^5, 1^4$	2	5				

There are 29 types of degree 12, each of length ≤ 4 .

#	hom. type	ℓ	pr.	#	hom. type	ℓ	pr.	#	hom. type	ℓ	pr.
12	$11, 1^{22}$	1	1	12.10	$7, 5^2, 3^4, 2^2$	3	5.2	12.20	$6^2, 5, 4, 3^2, 2^3$	3	63(73)
12.1	$10, 2^{10}, 1^3$	2	2	12.11	$7, 5, 4^4, 1^5$	2	5	12.21	$6^2, 4^4, 2, 1^3$	3	6.1
12.2	$9, 3^6, 2, 1^4$	2	3	12.12	$7, 5, 4^3, 3, 2^3$	3	5.1	12.22	$6^2, 4^3, 3^2, 2, 1$	3	6.3
12.3	$9, 3^5, 2^4, 1$	2	3	12.13	$7, 5, 4^2, 3^4, 1$	3	5.1	12.23	$6, 5^3, 3^3, 2, 1$	3	62(82)
12.4	$8, 4^4, 3, 1^6$	2	4	12.14	$6^3, 5, 1^{10}$	2	6	12.24	$6, 5^2, 4^3, 2^2, 1$	3	6.1
12.5	$8, 4^4, 2^3, 1^3$	3	4.1	12.15	$6^3, 4, 2^4, 1^3$	3	6.1	12.25	$6, 5, 4^4, 3^2$	3	63(73)
12.6	$8, 4^3, 3^2, 2^3, 1$	3	4.1(5.1)	12.16	$6^3, 3^3, 2, 1^4$	3	6.2	12.26	$5^4, 4^2, 3, 1^2$	3	7.3
12.7	$8, 4^2, 3^5, 1^2$	2	4	12.17	$6^3, 3^2, 2^4, 1$	3	6.3	12.27	$5^4, 4, 3^3$	4	7.4
12.8	$7, 5^3, 2^4, 1^3$	3	5.1(7.1)	12.18	$6^2, 5^2, 2^5, 1$	3	6.1(7.1)	12.28	$5^3, 4^4, 2$	4	7.4
12.9	$7, 5^2, 4, 3^2, 2^2, 1^2$	3	5.1	12.19	$6^2, 5, 4, 3^3, 1^3$	3	6.2(7.2)				

We could of course continue like this but the number of homaloidal types grows very quickly. We give below the homaloidal types of length $\ell \in \{2, \dots, 7\}$ of smallest degree:

ℓ	d	mult.	ℓ	d	mult.	ℓ	d	mult.	ℓ	d	mult.
2	4	$2^3, 1^3$	4	12	$5^4, 4, 3^3$	5	16	$6^5, 5^3$	6	27	$10^4, 9^4, 2$
3	7	$3^4, 2^3$	4	12	$5^3, 4^4, 2$	6	27	$11^4, 10, 6^4$	7	38	$14^6, 11^2, 5$

4.2. Automorphisms of the affine plane. As explained before, there is a natural length in the group $\text{Aut}(\mathbb{A}^2)$, since this one is an amalgamated product of $\text{Aff}_2 = \text{Aut}(\mathbb{P}^2) \cap \text{Aut}(\mathbb{A}^2)$ and $\text{Jonq}_{p, \mathbb{A}^2} = \text{Jonq}_p \cap \text{Aut}(\mathbb{A}^2)$, where we fix a linear embedding $\mathbb{A}^2 \hookrightarrow \mathbb{P}^2$, and a point $p \in \mathbb{P}^2$ outside the image. By construction, the length of an element of $\text{Aut}(\mathbb{A}^2)$, viewed in the amalgamated product, is at least equal to its length in $\text{Bir}(\mathbb{P}^2)$. We show in Proposition 4.2 that the two lengths are in fact equal, using the following result.

Lemma 4.1. *Let $f, g \in \text{Bir}(\mathbb{P}^2)$ be elements such that $\text{Base}(f) \cap \text{Base}(g^{-1}) = \emptyset$. Then,*

$$\begin{aligned} \deg(f \circ g) &= \deg(f) \cdot \deg(g), & |\text{Base}(f \circ g)| &= |\text{Base}(f)| + |\text{Base}(g)| \quad \text{and} \\ \text{lgth}(f \circ g) &= \text{lgth}(f) + \text{lgth}(g). \end{aligned}$$

Proof. The two first equalities follow from Lemma 3.15(1). We prove the third one by induction on $\text{lgth}(g)$, the case where $\text{lgth}(g) = 0$ being obvious, since $g \in \text{Aut}(\mathbb{P}^2)$ in that case.

We then consider the case where $d = \deg(g) = \deg(g^{-1}) > 1$, and take $\varphi \in \text{Jonq} \subseteq \text{Bir}(\mathbb{P}^2)$ such that $g_1 = g \circ \varphi^{-1}$ is a predecessor of g . Then, $g_1^{-1}(e_0) = \varphi(g^{-1}(e_0))$ is a predecessor of $g^{-1}(e_0)$ (Corollary 3.27). This implies that $\varphi((f \circ g)^{-1}(e_0))$ is a predecessor of $(f \circ g)^{-1}(e_0)$ (Lemma 3.15) and thus that $f \circ g_1 = f \circ g \circ \varphi^{-1}$ is a predecessor of $f \circ g$ (Corollary 3.27). Hence, we obtain $\text{lgth}(g_1) = \text{lgth}(g) - 1$ and $\text{lgth}(f \circ g_1) = \text{lgth}(f \circ g) - 1$ (Theorem 1).

Since $\text{Base}(\varphi) \subseteq \text{Base}(g)$ (Lemma 1.3), we have $\text{Base}(g_1^{-1}) = \text{Base}(\varphi \circ g^{-1}) \subseteq \text{Base}(g^{-1})$ (Corollary 2.25), and thus $\text{Base}(f) \cap \text{Base}(g_1^{-1}) = \emptyset$. We can thus apply the induction hypothesis to get $\text{lgth}(f \circ g_1) = \text{lgth}(f) + \text{lgth}(g_1)$. This achieves the proof. \square

Proposition 4.2. *Let $f \in \text{Aut}(\mathbb{A}^2)$. Taking an inclusion $\text{Aut}(\mathbb{A}^2) \hookrightarrow \text{Bir}(\mathbb{P}^2)$ given by a linear embedding $\mathbb{A}^2 \hookrightarrow \mathbb{P}^2$, the length of f in $\text{Bir}(\mathbb{P}^2)$ is equal to its length in the amalgamated product of $\text{Aut}(\mathbb{A}^2)$.*

Proof. We write $f = a_n \circ j_n \circ \cdots \circ a_1 \circ j_1 \circ a_0$, where each a_i is an element of Aff_2 and each j_i is an element of $\text{Jonq}_{p, \mathbb{A}^2}$. If f belongs to A , then its length is 0 in $\text{Bir}(\mathbb{P}^2)$ and in the amalgamated product, so we can assume that $n \geq 1$, that $j_i \in \text{Jonq}_{p, \mathbb{A}^2} \setminus \text{Aff}_2$ for $i = 1, \dots, n$ and that $a_i \in \text{Aff}_2 \setminus \text{Jonq}_{p, \mathbb{A}^2}$ for $i = 1, \dots, n-1$. We then need to prove that $\text{lgth}(f) = n$. To do this, we first observe that j_i and $(j_i)^{-1}$ contract exactly one curve of \mathbb{P}^2 , namely the line $L_\infty = \mathbb{P}^2 \setminus \mathbb{A}^2$. It implies that $(j_i)^{-1}$ and j_i have only one proper base-point, and since both preserve lines through p , then p is the unique proper base-point of j_i and $(j_i)^{-1}$. Moreover, each a_i is an automorphism of \mathbb{P}^2 that does not fix p , for $i = 1, \dots, n-1$. Hence by induction the element $a_i \circ j_i \circ \cdots \circ a_1 \circ j_1 \circ a_0$ contracts the line L_∞ onto $a_i(p) \neq p$, which is the unique proper base-point of $(a_i \circ j_i \circ \cdots \circ a_1 \circ j_1 \circ a_0)^{-1}$. In particular, $(a_i \circ j_i \circ \cdots \circ a_1 \circ j_1 \circ a_0)^{-1}$ and j_{i+1} do not have any common base-point, so $\text{lgth}(a_{i+1} \circ j_{i+1} \circ \cdots \circ a_1 \circ j_1 \circ a_0) = \text{lgth}(a_i \circ j_i \circ \cdots \circ a_1 \circ j_1 \circ a_0) + 1$ for each i . This provides the result. \square

4.3. Decreasing the length and increasing the degree via a single Jonquières element. In this section, we mainly provide an example of a Cremona transformation f and a Jonquières element φ such that $\text{lgth}(f \circ \varphi^{-1}) = \text{lgth}(f) - 1$ and $\deg(f \circ \varphi^{-1}) > \deg(f)$:

Proposition 4.3. *Fixing 8 general points $p_0, \dots, p_7 \in \mathbb{P}^2$, the following hold:*

(1) *There exists a birational involution $f \in \text{Bir}(\mathbb{P}^2)$ such that*

$$f(e_0) = 17e_0 - \sum_{j=0}^7 6e_{p_j} \text{ and } f(e_i) = 6e_0 - e_{p_i} - 2 \sum_{j=0}^7 e_{p_j} \text{ for } i = 0, \dots, 7.$$

(2) *For each general point $q \in \mathbb{P}^2$, there exists a Jonquières element φ_q such that*

$$\varphi_q^{-1}(e_0) = 5e_0 - 4e_{p_0} - e_{p_1} - \cdots - e_{p_7} - e_q.$$

(3) *For all f and φ_q as in (1) and (2), the birational map $f_q = f \circ \varphi_q^{-1}$ satisfies:*

$$\text{lgth}(f_q) = 4 < 5 = \text{lgth}(f) \quad \text{and} \quad \deg(f_q) = 19 > 17 = \deg(f).$$

(4) *For any two distinct general points $q, q' \in \mathbb{P}^2$, and all choices of f_q, f'_q as in (3), we have*

$$f_{q'} \notin f_q \text{Aut}(\mathbb{P}^2).$$

Proof. (1)-(2): Let $U_D \subseteq (\mathbb{P}^2)^8$ the dense open subset such that for each $(p_0, \dots, p_7) \in U_D$ the points p_i are pairwise distinct and the blow-up $\pi: X \rightarrow \mathbb{P}^2$ of p_0, \dots, p_7 is a Del Pezzo surface (see Lemma 4.4 below). For each $(p_0, \dots, p_7) \in U_D$, we can then define $\hat{f} \in \text{Aut}(X)$ to be the *Bertini involution* (see [Dol2012, §8.8.2]) and obtain that $f = \pi \circ \hat{f} \circ \pi^{-1} \in \text{Bir}(\mathbb{P}^2)$ has degree 17 and satisfies the conditions given in (1). Since $(5; 4, 1^8)$ is a homaloidal type, Corollary 2.27 yields a dense open subset $V_9 \subseteq (\mathbb{P}^2)^9$ such that for all each $(p_0, \dots, p_7, q) \in V_9$ there exists $\varphi_q \in \text{Bir}(\mathbb{P}^2)$ satisfying the condition given in (2). The element φ_q is then Jonquières (Lemma 2.42). We then denote by V the open set $V = V_9 \cap (U_D \times \mathbb{P}^2) \subseteq (\mathbb{P}^2)^9$, by $\kappa: (\mathbb{P}^2)^9 \rightarrow (\mathbb{P}^2)^8$ the projection on

the first eight factors, and by $U \subseteq (\mathbb{P}^2)^8$ the open set given by $U = \kappa(V)$. For each $(p_0, \dots, p_7) \in U$, the assertions (1) and (2) are then satisfied.

(3): We take f and φ_q as in (1) and (2), write $f_q = f \circ \varphi_q^{-1}$, and observe that

$$f_q(e_0) = f(5e_0 - 4e_{p_0} - e_{p_1} - \dots - e_{p_7} - e_q) = 19e_0 - 4e_{p_0} - 7e_{p_1} - \dots - 7e_{p_7} - f(e_q),$$

whence the homaloidal type of f_q^{-1} is $(19; 7^7, 4, 1)$. In particular, we have $\deg(f_q) = \deg(f_q^{-1}) = 19$. Recall that we also have $\text{lgth}(f_q) = \text{lgth}(f_q^{-1})$. Therefore, to show that $\text{lgth}(f) = 5$ and $\text{lgth}(f_q) = 4$, it suffices to look at Examples 3.17 and 3.18, where we already observed that Algorithm 3.16 applied to $(19; 7^7, 4, 1)$ and $(17; 6^8)$ yields the following sequences of homaloidal types:

$$\begin{aligned} (17; 6^8) &\rightarrow (14; 6, 5^6, 3) \rightarrow (8; 3^7) \rightarrow (5; 2^6) \rightarrow (3; 2, 1^4) \rightarrow (1), \\ (19; 7^7, 4, 1) &\rightarrow (13; 5^6, 4, 1^2) \rightarrow (8; 4, 3^5, 1^2) \rightarrow (4; 3, 1^6) \rightarrow (1). \end{aligned}$$

One could also check that the homaloidal type of f_q is $(19; 11, 8, 5^7)$. However, it is useless for the proof we propose.

(4): Take two general points q, q' and suppose that $f_{q'} = f_q \circ \alpha$ for some $\alpha \in \text{Aut}(\mathbb{P}^2)$. Then, $f_q(e_0) = f_{q'}(e_0)$, which implies that $f(e_q) = f(e_{q'})$, whence $q = q'$. \square

The following result is classical:

Lemma 4.4. *Let $\pi: X \rightarrow \mathbb{P}^2$ be the blow-up of r distinct proper points, with $1 \leq r \leq 8$. Then, X is a Del Pezzo surface if and only if the following conditions are satisfied: no 3 of the points are collinear, no 6 lie on the same conic, and no 8 of the points lie on the same cubic singular at one of the points. Moreover, these conditions correspond to a dense open subset of $(\mathbb{P}^2)^r$, and are thus satisfied for sufficiently general points.*

Proof. Follows from [Dol2012, Proposition 8.1.25]. \square

4.4. The number of predecessors is not uniformly bounded. Let $f \in \text{Bir}(\mathbb{P}^2)$ be a Cremona transformation. If $\deg(f) \leq 4$, one can check that f admits a unique predecessor up to right multiplication by an element of $\text{Aut}(\mathbb{P}^2)$. If $\deg(f) \leq 5$, then f may admit more than one predecessor modulo $\text{Aut}(\mathbb{P}^2)$ (but all having the same homaloidal type, by Lemma 1.3). Example 4.5 gives an example of degree 5 with two predecessors having distinct configurations of base-points, and Lemma 4.6 shows that the number of predecessors modulo $\text{Aut}(\mathbb{P}^2)$ is not uniformly bounded.

Example 4.5. Let us consider the birational involution $g \in \text{Bir}(\mathbb{P}^2)$ given by

$$g: [x : y : z] \mapsto \left[\frac{x}{-\frac{1}{x} + \frac{1}{y} + \frac{1}{z}} : \frac{y}{\frac{1}{x} - \frac{1}{y} + \frac{1}{z}} : \frac{z}{\frac{1}{x} + \frac{1}{y} - \frac{1}{z}} \right] = [xvw : yuw : zuv]$$

where $u = -yz + xz + xy$, $v = yz - xz + xy$, and $w = yz + xz - xy$. We see that g is of degree 5. It has moreover 6 base-points of multiplicity 2, namely the 3 points $p_1 = [1 : 0 : 0]$, $p_2 = [0 : 1 : 0]$, $p_3 = [0 : 0 : 1]$, and 3 other points q_1, q_2, q_3 , where each q_i is infinitely near to p_i . The homaloidal type of g is therefore $(5; 2^6)$.

The algorithm consists of applying a cubic birational transformation whose linear system consists of cubics singular at one of the p_i and passing through 4 of the remaining 5 base-points. The predecessors of g are thus of degree 3. However, we get distinct classes up to automorphism of \mathbb{P}^2 , depending on our choice of the 4 points.

Denoting by $\rho_1, \rho_2 \in \text{Bir}(\mathbb{P}^2)$ the birational maps of degree 3 given by

$$\begin{aligned} \rho_1: [x : y : z] &\mapsto \left[\frac{z}{x(\frac{1}{x} + \frac{1}{y} - \frac{1}{z})} : y : z \right] = [yz^2 : yw : zw] \text{ and} \\ \rho_2: [x : y : z] &\mapsto \left[\frac{1}{-\frac{1}{x} + \frac{1}{y} + \frac{1}{z}} : y : z \right] = [xyz : yu : zu], \end{aligned}$$

we observe that $\alpha_1 = \rho_1 g \rho_1^{-1}$ and $\alpha_2 = \rho_2 g \rho_2^{-1}$ are the two linear involutions

$$\alpha_1: [x : y : z] \mapsto [x : y : 2x - z] \text{ and } \alpha_2: [x : y : z] \mapsto [x : 2x - y : 2x - z].$$

For each $i \in \{1, 2\}$ the map $\rho_i \in \text{Bir}(\mathbb{P}^2)$ is a Jonquière map, preserving a general line through $[1 : 0 : 0]$. This implies that $\psi_i = \rho_i^{-1} \circ \alpha_i = g \circ \rho_i^{-1}$ is a predecessor of g .

The linear system of ρ_1 consists of cubics singular at p_1 and passing through p_2, p_3, q_1, q_2 . The linear system of ρ_2 consists of cubics singular at p_1 and passing through p_2, p_3, q_2, q_3 . The configuration of the points being different (for ρ_2 , there is a tangent direction fixed at the singular point, contrary to ρ_1), this shows that $\psi_1 \notin \text{Aut}(\mathbb{P}^2) \circ \psi_2 \circ \text{Aut}(\mathbb{P}^2)$.

Lemma 4.6. *For each integer $i \geq 1$, there exists $f \in \text{Bir}(\mathbb{P}^2)$ which has at least i predecessors up to left and right composition with elements of $\text{Aut}(\mathbb{P}^2)$.*

Proof. We choose n such that $2n - 1 \geq i$ and $2n - 1 \geq 5$, and then observe that $\chi = (n^2 + 1; n^2 - n + 1, n^{2n-1}, 1^{2n-1})$ is a homaloidal type, whose predecessor is $\chi_1 = (n; n - 1, 1^{2n-2})$. We take $4n - 1$ general points $p_0, p_1, \dots, p_{2n-1}, q_1, \dots, q_{2n-1} \in \mathbb{P}^2$ and choose $f \in \text{Bir}(\mathbb{P}^2)$ such that $f^{-1}(e_0) = (n^2 + 1)e_0 - (n^2 - n + 1)e_{p_0} - \sum_{i=1}^{2n-1} n e_{p_i} - \sum_{i=1}^{2n-1} e_{q_i}$ (which exists by Lemma 2.26).

For each $j \in \{1, \dots, 2n - 1\}$, there exists a Jonquière element $\varphi_j \in \text{Bir}(\mathbb{P}^2)$ such that $(\varphi_j)^{-1}(e_0) = n e_0 - (n - 1)e_{p_0} - \sum_{i=1}^{2n-1} e_{p_i} - e_{q_j}$ (again by Lemma 2.26). Then, $f_j = f \circ \varphi_j^{-1}$ is a predecessor of f (follows from Lemma 3.11(3) and Lemma 3.20).

As $\text{Base}(\varphi_j) \subseteq \text{Base}(f)$, we get $\text{Base}(f_j^{-1}) = \text{Base}(\varphi_j \circ f^{-1}) \subseteq \text{Base}(f^{-1})$ (Corollary 2.25). Moreover, f_j is Jonquière of degree n (because of the type of χ_1), so the same holds for f_j^{-1} .

It remains to see that if $j, k \in \{1, \dots, 2n - 1\}$ are such that $j \neq k$, then there are no elements $\alpha, \beta \in \text{Aut}(\mathbb{P}^2)$ such that $f_j = \alpha \circ f_k \circ \beta$. Indeed, otherwise we would have $f_j(e_0) = \alpha(f_k(e_0))$, so α sends the $2n - 1$ base-points of f_k^{-1} onto those of f_j^{-1} , respecting the multiplicities. Note that $f_j(e_0) \neq f_k(e_0)$, because $\varphi_j^{-1}(e_0) \neq \varphi_k^{-1}(e_0)$. The map $\alpha \in \text{Aut}(\mathbb{P}^2)$ has then to send a sequence of $2n - 1$ points of $\text{Base}(f^{-1})$ onto another sequence of $2n - 1$ points of $\text{Base}(f^{-1})$. This is impossible, as the points are general points and $2n - 1 \geq 5$. \square

4.5. Reduced decompositions of arbitrary lengths. Recall that a reduced decomposition of an element $f \in \text{Bir}(\mathbb{P}^2)$ is a product $f = \varphi_n \circ \dots \circ \varphi_1$ of Jonquière elements such that $\varphi_{i+1} \circ \varphi_i$ is not Jonquière for $i = 1, \dots, n - 1$. One can of course always obtain a reduced decomposition by starting with any decomposition, simply replacing φ_i and φ_{i+1} with their product if this one is Jonquière. Proposition 4.10 shows that the length of reduced decompositions is unbounded.

We begin with the following classical result:


Lemma 4.7. *Let p_0, p_1, p_2 be three non-collinear points of \mathbb{P}^2 . There is a quadratic birational involution $\nu \in \text{Bir}(\mathbb{P}^2)$ preserving the pencil of lines through any of the three base-points and satisfying $\text{Base}(\nu) = \{p_0, p_1, p_2\}$.*

Proof. It suffices to change coordinates to have $\{p_0, p_1, p_2\} = \{[1 : 0 : 0], [0 : 1 : 0], [0 : 0 : 1]\}$ and to choose $\nu : [x : y : z] \dashrightarrow [yz : xz : xy]$. \square

Lemma 4.8. *Let $p_0, \dots, p_5 \in \mathbb{P}^2$ be six distinct points such that no 3 of them are collinear and not all lie on the same conic. Then, there exist three quadratic birational involutions $\psi_1, \psi_2, \psi_3 \in \text{Bir}(\mathbb{P}^2)$, each having (three) proper base-points, such that the following hold:*

- (1) $\text{Base}(\psi_1) = \{p_0, p_1, p_2\}$;
- (2) $\text{Base}(\psi_2) \cap \text{Base}(\psi_1) = \text{Base}(\psi_3) \cap \text{Base}(\psi_2) = \emptyset$;
- (3) $(\psi_3 \circ \psi_2 \circ \psi_1)^{-1}(e_0) = 5e_0 - 2 \sum_{i=0}^5 e_{p_i}$.

Proof. The blow-up $\pi_1 : X \rightarrow \mathbb{P}^2$ of the six points p_0, \dots, p_5 is a Del Pezzo surface (Lemma 4.4) and there exists a quadratic birational involution $\psi_1 \in \text{Bir}(\mathbb{P}^2)$ with $\text{Base}(\psi_1) = \{p_0, p_1, p_2\}$ (Lemma 4.7). We then write $q_i = \psi_1(p_i) \in \mathbb{P}^2$ for $i = 3, 4, 5$. We now prove that $\pi_2 = \psi_1 \circ \pi_1 : X \rightarrow \mathbb{P}^2$ is the blow-up of $p_0, p_1, p_2, q_3, q_4, q_5$: we have a commutative diagram



where η is the blow-up of p_0, p_1, p_2 , $\hat{\psi}_1 \in \text{Aut}(Y)$ is an automorphism of order 2, τ_1 is the blow-up of $\{\eta^{-1}(p_i) \mid i = 3, 4, 5\}$, and τ_2 is the blow-up of $\{\eta^{-1}(q_i) = \hat{\psi}_1(\eta^{-1}(p_i)) \mid i = 3, 4, 5\}$.

Because X is a Del Pezzo surface, the points q_3, q_4, q_5 are not collinear (Lemma 4.4), so there is a quadratic birational involution $\psi_2 \in \text{Bir}(\mathbb{P}^2)$ with $\text{Base}(\psi_2) = \{q_3, q_4, q_5\}$ (Lemma 4.7). We then write $q_i = \psi_2(p_i) \in \mathbb{P}^2$ for $i = 0, 1, 2$ and obtain similarly that $q_0, \dots, q_5 \in \mathbb{P}^2$ are such that no 3 of them are collinear (since $\psi_2 \circ \psi_1 \circ \pi_1 : X \rightarrow \mathbb{P}^2$ is the blow-up of q_0, \dots, q_5).

We now choose a quadratic birational involution $\psi_3 \in \text{Bir}(\mathbb{P}^2)$ with $\text{Base}(\psi_3) = \{q_0, q_1, q_2\}$ (again by Lemma 4.7). It remains to calculate

$$\begin{aligned} (\psi_3 \circ \psi_2 \circ \psi_1)^{-1}(e_0) &= \psi_1(\psi_2(2e_0 - e_{q_0} - e_{q_1} - e_{q_2})) \\ &= \psi_1(4e_0 - e_{p_0} - e_{p_1} - e_{p_2} - 2e_{q_3} - 2e_{q_4} - 2e_{q_5}) \\ &= 5e_0 - 2 \sum_{i=0}^5 e_{p_i}, \end{aligned}$$

where we have used the fact that $\psi_1(e_{p_0}) = e_0 - e_{p_1} - e_{p_2}$, $\psi_1(e_{p_1}) = e_0 - e_{p_0} - e_{p_2}$, and $\psi_1(e_{p_2}) = e_0 - e_{p_0} - e_{p_1}$ (see Example 2.15). \square

Corollary 4.9. *There exist quadratic birational maps $\varphi_1, \dots, \varphi_6$, each having (three) proper base-points, such that*

- (1) $\varphi_6 \circ \varphi_5 \circ \varphi_4 \circ \varphi_3 \circ \varphi_2 \circ \varphi_1 = \text{id}$;
- (2) $\text{Base}(\varphi_i^{-1}) \cap \text{Base}(\varphi_{i+1}) = \emptyset$ for $i = 1, \dots, 5$.

Proof. Let $p_0, \dots, p_5 \in \mathbb{P}^2$ be six distinct points such that no 3 are collinear and not all lie on the same conic. Then, choose ψ_1, ψ_2, ψ_3 as in Lemma 4.8. Recall that we have in particular $(\psi_3 \circ \psi_2 \circ \psi_1)^{-1}(e_0) = 5e_0 - 2 \sum_{i=0}^5 e_{p_i}$. Applying Lemma 4.8 to the same points, but taken in the order $p_3, p_4, p_5, p_0, p_1, p_2$, we get quadratic birational involutions $\psi'_1, \psi'_2, \psi'_3 \in \text{Bir}(\mathbb{P}^2)$, each having three proper base-points, such that $\text{Base}(\psi'_1) = \{p_3, p_4, p_5\}$, $\text{Base}(\psi'_2) \cap \text{Base}(\psi'_1) = \text{Base}(\psi'_3) \cap \text{Base}(\psi'_2) = \emptyset$ and $(\psi'_3 \circ \psi'_2 \circ \psi'_1)^{-1}(e_0) = 5e_0 - 2 \sum_{i=0}^5 e_{p_i}$.

The birational map $\alpha = \psi'_3 \circ \psi'_2 \circ \psi'_1 \circ \psi_1 \circ \psi_2 \circ \psi_3$ satisfies then $\alpha^{-1}(e_0) = e_0$, so that it is an automorphism of \mathbb{P}^2 . It remains to choose $(\varphi_6, \dots, \varphi_1) := (\psi'_3, \psi'_2, \psi'_1, \psi_1, \psi_2, \psi_3 \circ \alpha^{-1})$ to obtain the result. \square

Proposition 4.10. *For each $f \in \text{Bir}(\mathbb{P}^2)$ the set of reduced decompositions of f has unbounded length.*

Proof. To prove the result, we start with a reduced decomposition $f = \varphi_n \circ \dots \circ \varphi_1$, and construct another one, with length $\geq n+5$. To do this, we take quadratic birational maps $\psi_1, \dots, \psi_6 \in \text{Bir}(\mathbb{P}^2)$, each having three proper base-points, such that $\psi_6 \circ \dots \circ \psi_1 = \text{id}$ and $\text{Base}(\psi_i^{-1}) \cap \text{Base}(\psi_{i+1}) = \emptyset$ for $i = 1, \dots, 5$ (which exist by Corollary 4.9). For $i = 1, \dots, 5$, we observe that $\text{lgth}(\psi_{i+1} \circ \psi_i) = 2$ (Lemma 4.1), so $\psi_{i+1} \circ \psi_i$ is not Jonquières. Replacing all ψ_i with $\alpha \circ \psi_i \circ \alpha^{-1}$ for some general $\alpha \in \text{Aut}(\mathbb{P}^2)$, we can assume that $\text{Base}(\varphi_n^{-1}) \cap \text{Base}(\psi_1) = \emptyset$, which implies that $\text{lgth}(\psi_1 \circ \varphi_n) = \text{lgth}(\varphi_n) + 1$ (Lemma 4.1) and is thus not Jonquières if $\varphi_n \notin \text{Aut}(\mathbb{P}^2)$. In this latter case, we obtain a reduced decomposition of f of length $n+6$ as $f = \psi_6 \circ \dots \circ \psi_1 \circ \varphi_n \circ \dots \circ \varphi_1$. The last case is when $\varphi_n \in \text{Aut}(\mathbb{P}^2)$. This implies that $n = 1$, as otherwise $\varphi_n \circ \varphi_{n-1}$ would be Jonquières. Hence, $f = \varphi_n \in \text{Aut}(\mathbb{P}^2)$. In this case, it suffices to write $f = \psi'_6 \circ \psi_5 \circ \dots \circ \psi_1$ with $\psi'_6 = f \circ \psi_6$, to get a reduced decomposition of length $6 = n+5$. \square

4.6. Examples of dynamical lengths.

Lemma 4.11. *The element $\kappa \in \text{Bir}(\mathbb{P}^2)$ given by $[x : y : z] \dashrightarrow [yz + x^2 : xz : z^2]$ satisfies*

$$\text{lgth}(\kappa^a) = a, \quad \deg(\kappa^a) = 2^a, \quad |\text{Base}(\kappa^a)| = 3a$$

for each integer $a \geq 1$. In particular, we have

$$\mathfrak{d}_{\text{lgth}}(\kappa^m) = m, \quad \lambda(\kappa^m) = 2^m, \quad \mu(\kappa^m) = 3m$$

for each $m \geq 1$.

Proof. As κ is a Jonquières element of degree 2, we have $\text{lgth}(\kappa) = 1$, $\deg(\kappa) = 2$ and $|\text{Base}(\kappa^a)| = 3$ (this last assertion follows from Noether equalities, see Lemma 2.20).

Denoting by $L \subset \mathbb{P}^2$ the line given by $z = 0$, the restriction of κ is automorphism of $\mathbb{P}^2 \setminus L \simeq \mathbb{A}^2$, so the same holds for κ^a , for each $a \in \mathbb{Z}$. There can then be at most one proper base-point of κ^a , namely the image by κ^{-a} of the line $z = 0$. We check that κ contracts L onto $q = [1 : 0 : 0]$, which is then the unique proper base-point of κ^{-1} , and that $p = [0 : 1 : 0]$ is the unique proper base-point of κ . This implies that κ^a contracts the line L onto q for each $a \geq 1$, and thus q is the unique proper base-point of κ^{-a} for each $a \geq 1$. We obtain, for each $a \geq 1$, that $\text{Base}(\kappa) \cap \text{Base}(\kappa^{-a}) = \emptyset$. Lemma 4.1 then yields $\deg(\kappa^{a+1}) = \deg(\kappa) \cdot \deg(\kappa^a)$, $|\text{Base}(\kappa^{a+1})| = |\text{Base}(\kappa)| + |\text{Base}(\kappa^a)|$ and $\text{lgth}(\kappa^{a+1}) = \text{lgth}(\kappa) + \text{lgth}(\kappa^a)$. This provides the result, by induction over a . \square

Lemma 4.12. *Choosing $\sigma, \alpha_2, \alpha_3 \in \text{Bir}(\mathbb{P}^2)$ as $\sigma: [x : y : z] \dashrightarrow [yz : xz : xy]$,*

$$\alpha_2: [x : y : z] \mapsto [z + y : x + z : y] \text{ and } \alpha_3: [x : y : z] \mapsto [y + z : x : y]$$

we obtain $\mathfrak{d}_{\text{lgth}}(\alpha_2\sigma) = \frac{1}{2}$ and $\mathfrak{d}_{\text{lgth}}(\alpha_3\sigma) = \frac{1}{3}$.

Proof. The birational involution $\sigma = \sigma^{-1}$ is quadratic with base-points $p_1 = [1 : 0 : 0]$, $p_2 = [0 : 1 : 0]$, $p_3 = [0 : 0 : 1]$ and its action on $\mathcal{Z}_{\mathbb{P}^2}$ satisfies

$$\sigma(e_0) = 2e_0 - e_{p_1} - e_{p_2} - e_{p_3}, \quad \sigma(e_{p_i}) = e_0 - e_{p_1} - e_{p_2} - e_{p_3} + e_{p_i}, \quad i = 1, 2, 3,$$

as in Example 2.15. Writing $p_4 = [1 : 0 : 1]$, $p_5 = [1 : 1 : 0]$, one gets

$$\alpha_2(p_1) = p_2, \quad \alpha_2(p_2) = p_4, \quad \alpha_2(p_3) = p_5 \text{ and } \alpha_3(p_1) = p_2, \quad \alpha_3(p_2) = p_4, \quad \alpha_3(p_3) = p_1.$$

Since p_4 and p_5 are general points of the lines contracted by σ onto respectively p_2 and p_3 , we find $\sigma(e_{p_4}) = e_{q_2}$ and $\sigma(e_{p_5}) = e_{q_3}$, where q_2, q_3 are points infinitely near to p_2 and p_3 respectively. In particular, writing

$$\begin{aligned} T_2 &= \left\{ \bigoplus \mathbb{Z}e_q \mid q \in \mathcal{B}(\mathbb{P}^2), q \text{ is infinitely near } p_4 \text{ or } p_5 \right\} \subseteq \mathcal{Z}(\mathbb{P}^2), \\ T_3 &= \left\{ \bigoplus \mathbb{Z}e_q \mid q \in \mathcal{B}(\mathbb{P}^2), q \text{ is infinitely near } p_4 \right\} \subseteq \mathcal{Z}(\mathbb{P}^2), \end{aligned}$$

and writing $V_i = \mathbb{Z}e_0 \oplus \mathbb{Z}e_{p_1} \oplus \mathbb{Z}e_{p_2} \oplus \mathbb{Z}e_{p_3} \oplus T_i$, one observes that

$$\alpha_i\sigma(T_i) \subseteq T_i \text{ and } \alpha_i\sigma(V_i) \subseteq V_i \text{ for } i = 2, 3.$$

We then get, for $i = 2, 3$, a linear map $V_i/T_i \rightarrow V_i/T_i$ given by a matrix M_i with respect to the basis $e_0, e_{p_1}, e_{p_2}, e_{p_3}$, as follows :

$$M_2 = \begin{pmatrix} 2 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 \\ -1 & 0 & -1 & -1 \\ 0 & 0 & 0 & 0 \end{pmatrix} \text{ and } M_3 = \begin{pmatrix} 2 & 1 & 1 & 1 \\ -1 & -1 & -1 & 0 \\ -1 & 0 & -1 & -1 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

We then compute

$$A := (M_2)^2 = \begin{pmatrix} 3 & 2 & 1 & 1 \\ 0 & 0 & 0 & 0 \\ -1 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \text{ and } B := (M_3)^3 = \begin{pmatrix} 3 & 1 & 1 & 2 \\ -1 & 0 & -1 & -1 \\ -1 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

Let us check that $f := (\alpha_2\sigma)^2$ and $g := (\alpha_3\sigma)^3$ have dynamical lengths equal to 1.

The expression of A shows us that $\deg f = 3$ and $\text{comult}(f) \leq 3 - 2 = 1$, so that f is a Jonquière's transformation. Set $d_{-1} = 0$, $d_0 = 1$ and $d_n = 3d_{n-1} - d_{n-2}$ for $n \geq 1$. A straightforward induction on n would show that for each nonnegative integer n , the coefficients $(1, 1)$ and $(1, 3)$ of A^n satisfy:

$$(A^n)_{1,1} = d_n, \quad (A^n)_{1,3} = d_n - d_{n-1}.$$

Since $2(d_n - d_{n-1}) + 1 > d_n$, it follows from Corollary 2.29 that $d_n - d_{n-1}$ is the highest multiplicity of f^n . Therefore, a predecessor g_n of f^n satisfies

$$\deg g_n \geq \text{comult}(f^n) = d_n - (d_n - d_{n-1}) = d_{n-1} = \deg f^{n-1}.$$

This proves that f^{n-1} is a predecessor of f^n , so that $\text{lgth}(f^n) = \text{lgth}(f^{n-1}) + 1$, proving that $\text{lgth}(f^n) = n$ and $\mathfrak{d}_{\text{lgth}}(f) = 1$.

One would prove analogously that g has dynamical degree 1, since the coefficients $(1, 1)$ and $(1, 4)$ of B^n satisfy:

$$(B^n)_{1,1} = d_n, \quad (B^n)_{1,4} = d_n - d_{n-1}. \quad \square$$

Corollary 4.13. *We have*

$$\frac{1}{2}\mathbb{Z}_{\geq 0} \cup \frac{1}{3}\mathbb{Z}_{\geq 0} \subseteq \mathfrak{d}_{\text{lgth}}(\text{Bir}(\mathbb{P}^2)) = \{\mathfrak{d}_{\text{lgth}}(f) \mid f \in \text{Bir}(\mathbb{P}^2)\}$$

Proof. Lemma 4.12 yields elements $f_2, f_3 \in \text{Bir}(\mathbb{P}^2)$ such that $\mathfrak{d}_{\text{lgth}}(f_2) = \frac{1}{2}$ and $\mathfrak{d}_{\text{lgth}}(f_3) = \frac{1}{3}$. We then get $\mathfrak{d}_{\text{lgth}}(f_2^m) = \frac{m}{2}$ and $\mathfrak{d}_{\text{lgth}}(f_3^m) = \frac{m}{3}$ for each $m \geq 0$. \square

4.7. Length of monomial transformations. Let us recall that the group $\text{GL}_2(\mathbb{Z})$ can be viewed as the subgroup of monomial transformations of $\text{Bir}(\mathbb{P}^2)$: a matrix $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ corresponds to the transformation $[x : y : 1] \dashrightarrow [x^a y^b : x^c y^d : 1]$. In this section, we give an easy algorithm to compute the length and the dynamical length of any such element in $\text{Bir}(\mathbb{P}^2)$. We first study *ordered* elements of $\text{SL}_2(\mathbb{Z})$ (see Definition 4.15 and Lemma 4.16 for description of these elements, and the relation with continued fractions). Their lengths are easy to compute (Proposition 4.19). We then give the length of every element of $\text{GL}_2(\mathbb{Z})$, by reducing to the case of ordered elements (Lemma 4.21). The dynamical length of every element of $\text{SL}_2(\mathbb{Z})$ is given in Proposition 4.25 at the end of the section. This provides the dynamical length of every element of $\text{SL}_2(\mathbb{Z})$, as $\mathfrak{d}_{\text{lgth}}(M) = \frac{1}{2}\mathfrak{d}_{\text{lgth}}(M^2)$ for each $M \in \text{GL}_2(\mathbb{Z})$.

We first recall the following :

Lemma 4.14. *Writing $\text{SL}_2(\mathbb{Z})_{\geq 0} = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}_2(\mathbb{Z}) \mid a, b, c, d \geq 0 \right\}$, we get:*

- (1) $\text{SL}_2(\mathbb{Z})_{\geq 0}$ is the free monoid generated by $L = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$ and $R = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$.
- (2) $\text{SL}_2(\mathbb{Z})_{\geq 0} = S_L \uplus S_R \uplus \{\text{id}\}$ (disjoint union), where

$$S_L = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}_2(\mathbb{Z}) \mid \begin{array}{l} a \geq b \geq 0 \\ c \geq d \geq 0 \end{array} \right\} = \text{SL}_2(\mathbb{Z})_{\geq 0} \cdot L,$$

$$S_R = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}_2(\mathbb{Z}) \mid \begin{array}{l} 0 \leq a \leq b \\ 0 \leq c \leq d \end{array} \right\} = \text{SL}_2(\mathbb{Z})_{\geq 0} \cdot R.$$

- (3) $L \cdot \text{SL}_2(\mathbb{Z})_{\geq 0} \cdot R = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}_2(\mathbb{Z}) \mid \begin{array}{l} 0 \leq a \leq b \leq d \\ 0 \leq a \leq c \leq d \end{array} \right\}$

Proof. We write

$$S_L = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}_2(\mathbb{Z}) \mid \begin{array}{l} a \geq b \geq 0 \\ c \geq d \geq 0 \end{array} \right\}, S_R = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}_2(\mathbb{Z}) \mid \begin{array}{l} 0 \leq a \leq b \\ 0 \leq c \leq d \end{array} \right\}$$

and obtain $S_L \cup S_R \subseteq \text{SL}_2(\mathbb{Z})_{\geq 0}$. The sets S_L , S_R and $\{\text{id}\}$ are pairwise disjoint. To show $\text{SL}_2(\mathbb{Z})_{\geq 0} = S_L \cup S_R \cup \{\text{id}\}$, we take $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}_2(\mathbb{Z})_{\geq 0} \setminus \{\text{id}\}$ and show that $M \in S_L \cup S_R$. As $a, b, c, d \geq 0$ and $ad - bc = 1$, we have $a, d > 0$. If $b = 0$, then $a = d = 1$ and $c > 0$, so $M = L^c \in S_L$. Similarly, if $c = 0$, then $M = R^b \in S_R$.

We can thus assume that $a, b, c, d > 0$ and write $1 = ad - bc = (a - b)d - b(c - d)$, so $(a - b) \cdot (c - d) \geq 0$, which yields $M \in S_L \cup S_R$.

We then observe that $\mathrm{SL}_2(\mathbb{Z})_{\geq 0} \cdot L \subseteq S_L$ and $S_L \cdot L^{-1} \subseteq \mathrm{SL}_2(\mathbb{Z})_{\geq 0}$, which yield $\mathrm{SL}_2(\mathbb{Z})_{\geq 0} \cdot L = S_L$. We similarly obtain $\mathrm{SL}_2(\mathbb{Z})_{\geq 0} \cdot R = S_R$. This yields (2), which implies (1). Assertion (3) follows from

$$L \cdot \mathrm{SL}_2(\mathbb{Z})_{\geq 0} \cdot R = (L \cdot \mathrm{SL}_2(\mathbb{Z})_{\geq 0}) \cap (\mathrm{SL}_2(\mathbb{Z})_{\geq 0} \cdot R) = {}^t(\mathrm{SL}_2(\mathbb{Z})_{\geq 0} \cdot R) \cap (\mathrm{SL}_2(\mathbb{Z})_{\geq 0} \cdot R). \quad \square$$

Definition 4.15. We say that an element $M \in \mathrm{SL}_2(\mathbb{Z})$ is *ordered* if $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ satisfies $0 \leq a \leq b \leq d$ and $0 \leq a \leq c \leq d$. As noted in the previous proof, this implies that a is positive, so that all coefficients of M are positive.

Lemma 4.16. A matrix $M \in \mathrm{SL}_2(\mathbb{Z}) = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ is ordered if and only if it may be written under the form $M = L^{s_n} R^{s_{n-1}} \dots R^{s_3} L^{s_2} R^{s_1}$ for some integers $s_1, \dots, s_n \geq 1$ with $n \geq 2$ even. In this case, the integers s_1, \dots, s_n and a, b, c, d are linked by the continued fractions

$$\frac{b}{a} = s_1 + \frac{1}{s_2 + \frac{1}{\dots + \frac{1}{s_{n-1}}}} \quad \text{and} \quad \frac{d}{c} = s_1 + \frac{1}{s_2 + \frac{1}{\dots + \frac{1}{s_n}}}.$$

Proof. The fact that a matrix $M \in \mathrm{SL}_2(\mathbb{Z})$ is ordered if and only if it can be written $M = L^{s_n} R^{s_{n-1}} \dots R^{s_3} L^{s_2} R^{s_1}$ with $n \geq 2$ even and $s_1, \dots, s_n \geq 1$ follows from Lemma 4.14. We then prove the equalities given by the continued fractions by induction on n .

If $n = 2$, then $L^{s_2} R^{s_1} = \begin{pmatrix} 1 & s_1 \\ s_2 & s_1 s_2 + 1 \end{pmatrix}$, so $\frac{b}{a} = s_1$ and $\frac{d}{c} = \frac{s_1 s_2 + 1}{s_2} = s_1 + \frac{1}{s_2}$.

If $n > 2$, then $\begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} a' & b' \\ c' & d' \end{pmatrix} L^{s_2} R^{s_1}$, where $\frac{b'}{a'} = s_3 + \frac{1}{\dots + \frac{1}{s_{n-1}}}$ and $\frac{d'}{c'} = s_3 + \frac{1}{\dots + \frac{1}{s_n}}$. We replace these in $\frac{c}{d} = \frac{c' s_1 + d' (s_1 s_2 + 1)}{c' d' s_2} = s_1 + \frac{1}{s_2 + \frac{c'}{d'}}$ and $\frac{b}{a} = s_1 + \frac{1}{s_2 + \frac{b'}{a'}}$. \square

As we now explain, the length of ordered matrices of $\mathrm{SL}_2(\mathbb{Z})$ is easy to calculate:

Definition 4.17. For each sequence (s_1, \dots, s_n) of positive integers, with $n \geq 1$, we denote by $M(s_1, \dots, s_n) \in \mathrm{SL}_2(\mathbb{Z})$ the element given by

$$M(s_1, \dots, s_n) = \begin{cases} R^{s_n} L^{s_{n-1}} \dots R^{s_3} L^{s_2} R^{s_1} & \text{if } n \text{ is odd,} \\ L^{s_n} R^{s_{n-1}} \dots R^{s_3} L^{s_2} R^{s_1} & \text{if } n \text{ is even,} \end{cases}$$

where $L = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$, $R = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z})$ and denote by $\ell(s_1, \dots, s_n)$ the length of $M(s_1, \dots, s_n)$, viewed in $\mathrm{Bir}(\mathbb{P}^2)$.

Remark 4.18. The conjugation by $\tau = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \in \mathrm{GL}_2(\mathbb{Z})$ exchanges L and R , so $\ell(s_1, \dots, s_n)$ is also the length of

$$\tau M(s_1, \dots, s_n) \tau = \begin{cases} L^{s_n} R^{s_{n-1}} \dots L^{s_3} R^{s_2} L^{s_1} & \text{if } n \text{ is odd,} \\ R^{s_n} L^{s_{n-1}} \dots L^{s_3} R^{s_2} L^{s_1} & \text{if } n \text{ is even.} \end{cases}$$

Hence, the following lemma gives the length of any element of $\mathrm{SL}_2(\mathbb{Z})_{\geq 0}$.

Proposition 4.19. *Let $n \geq 1$. For each sequence (s_1, \dots, s_n) of positive integers, $M(s_1, \dots, s_n)$ is a product of $\ell(s_1, \dots, s_n)$ elements of length 1 that are of the form R^s, L^s, LR^s or RL^s for some $s \geq 1$ and thus belong to $\mathrm{SL}_2(\mathbb{Z})_{\geq 0}$. Moreover, we have:*

$$\ell(s_1, \dots, s_n) = \begin{cases} 1 & \text{if } n = 1, \\ 1 & \text{if } n = 2 \text{ and } s_2 = 1, \\ 2 & \text{if } n = 2 \text{ and } s_2 \neq 1, \\ \ell(s_2 - 1, s_3, \dots, s_n) + 1 & \text{if } n \geq 3, s_2 \geq 2, \\ \ell(s_3, \dots, s_n) + 1 & \text{if } n \geq 3, s_2 = 1. \end{cases}$$

Proof. We write $M = M(s_1, \dots, s_n) = \dots L^{s_2} R^{s_1}$. For each $s \geq 1$, R^s, LR^s are

$$R^s: [x : y : z] \mapsto [xy^s : yz^s : z^{s+1}], \quad LR^s: [x : y : z] \mapsto [xy^s z : xy^{s+1} : z^{s+2}],$$

and thus have length equal to 1. The same holds for L^s, RL^s (by conjugating with τ , see Remark 4.18). This gives the proof when $n = 1$ or $(n, s_2) = (2, 1)$. We thus assume $n \geq 3$, or $n = 2$ and $s_2 \geq 2$. Since $\mathrm{lgth}(LR^{s_1}) = 1$, we have $\mathrm{lgth}(M) \leq \mathrm{lgth}(M') + 1$, where $M' = M(LR^{s_1})^{-1}$. It remains to show that equality holds to obtain the result (using Remark 4.18). Applying Lemma 4.14, we can write

$$M = \begin{pmatrix} a & b \\ c & d \end{pmatrix}, M' = \begin{pmatrix} a' & b' \\ c' & d' \end{pmatrix} = \begin{pmatrix} a + (as_1 - b) & b - as_1 \\ c + (cs_1 - d) & d - cs_1 \end{pmatrix}$$

with $b \geq a \geq 0, d \geq c \geq 0$ and $a', b', c', d' \geq 0$. The degrees of M and M' , as birational maps of \mathbb{P}^2 , are respectively $D = \max\{a + b, c + d\} \geq 2$ and $D' = \max\{a' + b', c' + d'\} = \max\{a, c\}$. The element M corresponds to the birational map

$$M: [x : y : z] \mapsto [x^a y^b z^{D-a-b} : x^c y^d z^{D-c-d} : z^D],$$

which has degree D and exactly two proper base-points, namely $p_1 = [1 : 0 : 0]$ and $p_2 = [0 : 1 : 0]$, having multiplicity $m_1 = D - \max\{a, c\}$ and $m_2 = D - \max\{b, d\}$ respectively. Hence, p_1 is a base-point of maximal multiplicity and every predecessor of M has degree at least $D - m_1 = D'$. \square

Corollary 4.20. *Let $n \geq 1$. If s_1, \dots, s_n and s'_1 are positive integers, we have*

$$\ell(s_1, \dots, s_n) = \ell(s'_1, s_2, \dots, s_n).$$

Proof. Directly follows from Proposition 4.19. \square

We now give a way to compute the length of any element of $\mathrm{GL}_2(\mathbb{Z})$, by reducing to elements of $\mathrm{SL}_2(\mathbb{Z})$ of the form $M(s_1, \dots, s_n)$.


Lemma 4.21. *For each $M \in \mathrm{GL}_2(\mathbb{Z})$, the following hold:*

- (1) $\mathrm{lgth}(M) = 0 \Leftrightarrow M \in \mathrm{GL}_2(\mathbb{Z}) \cap \mathrm{Aut}(\mathbb{P}^2) = \left\langle \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 1 & -1 \\ 0 & -1 \end{pmatrix} \right\rangle \simeq \mathrm{Sym}_3$.
- (2) *There exist $A, B \in \mathrm{GL}_2(\mathbb{Z}) \cap \mathrm{Aut}(\mathbb{P}^2)$ such that either AMB or $-AMB$ belongs to $\mathrm{SL}_2(\mathbb{Z})_{\geq 0}$.*
- (3) *If $\mathrm{lgth}(M) \geq 1$, there exist $A, B \in \mathrm{GL}_2(\mathbb{Z}) \cap \mathrm{Aut}(\mathbb{P}^2)$ such that either AMB or $-AMB$ is equal to $M' = M(s_1, \dots, s_n)$ for some $n \geq 1$ and some positive integers $s_1, \dots, s_n \geq 1$. We then have $\mathrm{lgth}(M) = \mathrm{lgth}(M')$ (which can be computed directly by Proposition 4.19).*

Proof. (1): We observe that the group $\mathrm{GL}_2(\mathbb{Z}) \cap \mathrm{Aut}(\mathbb{P}^2)$ corresponds to the group Sym_3 of permutations of the coordinates, generated by $[x : y : z] \mapsto [y : x : z]$ and $[x : y : z] \mapsto [x : z : y]$, which correspond to $\tau = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ and $\nu = \begin{pmatrix} 1 & -1 \\ 0 & -1 \end{pmatrix}$.

(2): We consider the natural action of $\mathrm{GL}_2(\mathbb{Z})$ on the circle $\mathbb{P}^1(\mathbb{R}) \simeq \mathbb{S}^1$, via $\mathrm{GL}_2(\mathbb{Z}) \rightarrow \mathrm{PGL}_2(\mathbb{R})$. This action induces an isomorphism between $\mathrm{Sym}_3 = \mathrm{GL}_2(\mathbb{Z}) \cap \mathrm{Aut}(\mathbb{P}^2)$ and the group of permutations of the set $\Delta = \{[1 : 0], [0 : 1], [1 : 1]\}$. These three points delimit the three closed intervals of $\mathbb{P}^1(\mathbb{R})$ given by

$$I_1 = \{[\alpha : \beta] \mid \alpha \geq \beta \geq 0\}, I_2 = \{[\alpha : \beta] \mid 0 \leq \alpha \leq \beta\}, I_3 = \{[\alpha : \beta] \mid \alpha \geq 0, \beta \leq 0\}.$$



Suppose first that $M(\Delta)$ is contained in the union of two of these three intervals. Replacing M with AM where $A \in \mathrm{Sym}_3$, we can assume that $M(\Delta)$ is contained in the interval $I_1 \cup I_2 = \{[\alpha : \beta] \mid \alpha, \beta \in \mathbb{R}_{\geq 0}\}$.

The open interval I_3 being infinite, $M^{-1}(I_3)$ contains elements of $\mathbb{P}^1(\mathbb{R}) \setminus \Delta$. Replacing then M with MB , with $B \in \mathrm{Sym}_3$, we can assume that $M^{-1}(I_3) \cap I_3 \neq \emptyset$ or equivalently $M(I_3) \cap I_3 \neq \emptyset$.

We finish by replacing M with $\pm M$ or $\pm \tau M$, to assume moreover that $\det(M) = 1$ and that the first column of M has non-negative coefficients. It remains to observe that $M \in \mathrm{SL}_2(\mathbb{Z})_{\geq 0}$. Indeed, we have $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z})$ with $a, c \geq 0$ and $bd \geq 0$ since $[b : d] \in I_1 \cup I_2$. If $b, d \geq 0$, we are done. Otherwise $b, d \leq 0$, which yields $M(I_3) \subseteq I_1 \cup I_2$, contradicting $M(I_3) \cap I_3 \neq \emptyset$.

To finish the proof of *(2)*, we suppose that $M(\Delta)$ is not contained in the union of two of the three intervals I_1, I_2, I_3 and derive a contradiction. This implies that the three points of $M(\Delta)$ are in the interiors of three distinct intervals. Replacing M with $\pm AM$, with $A \in \mathrm{Sym}_3$, we can assume that $M([0 : 1]) \in I_1, M([1 : 0]) \in I_2, M([1 : 1]) \in I_3$, and that the coefficients of the first column of M are positive. The second column has then negative coefficients. We get $M = \begin{pmatrix} a & -b \\ c & -d \end{pmatrix}$ with $0 < a < c$ and $b > d > 0$. This yields $\det(M) = -ad + bc = a(b - d) + b(c - a) \geq 2$, a contradiction.

(3): Using *(2)*, we find $A, B \in \mathrm{GL}_2(\mathbb{Z}) \cap \mathrm{Aut}(\mathbb{P}^2)$ such that $M' = \pm AMB$ belongs to $\mathrm{SL}_2(\mathbb{Z})_{\geq 0}$. Since $\mathrm{lgth}(M) \geq 1$, then M' is not the identity. We can thus replace M' with $\tau M' \tau$ if needed and assume that $M' \in S_R = \mathrm{SL}_2(\mathbb{Z})_{\geq 0} \cdot R$ is an ordered matrix (follows from Lemma 4.14). This implies that M' has the desired form. It remains to prove that $\mathrm{lgth}(M) = \mathrm{lgth}(M')$. If $M' = ABM$, this is because $\mathrm{lgth}(A) = \mathrm{lgth}(B) = 0$. If $M' = -ABM$, we observe that $-M'$ is a product of $\mathrm{lgth}(-M')$ elements of length 1 of the form R^s, L^s, LR^s or $RL^s, s \geq 1$ (Proposition 4.19). Since

$$-R^s: [x : y : z] \mapsto [z^{s+1} : xy^{s-1}z : xy^s], \quad -LR^s: [x : y : z] \mapsto [z^{s+1} : yz^s : xy^s]$$

have length 1, the same hold for $-L^s, -RL^s$ (using conjugation by τ as in Remark 4.18). We thus get $\mathrm{lgth}(M') = \mathrm{lgth}(-M') = \mathrm{lgth}(M)$. \square

We now compute the dynamical length of elements of $\mathrm{GL}_2(\mathbb{Z})$. The case of ordered elements of $\mathrm{SL}_2(\mathbb{Z})$ is the easiest, as we now show:

Lemma 4.22. *Let $m, n \geq 1$ and let $(s_1, \dots, s_n), (t_1, \dots, t_m)$ be two sequences of positive integers, such that $t_1 \geq 2$ and $m \geq 2$. We then have*

$$\ell(s_1, \dots, s_n, t_1, \dots, t_m) = \ell(s_1, \dots, s_n) + \ell(t_1, \dots, t_m).$$

Proof. We prove the result by induction on n .

If $n = 1$, then Proposition 4.19 yields $\ell(s_1, t_1, \dots, t_m) = \ell(t_1 - 1, t_2, \dots, t_m) + 1 \stackrel{\text{Corollary 4.20}}{=} \ell(t_1, t_2, \dots, t_m) + 1 = \ell(s_1) + \ell(t_1, \dots, t_m)$.

If $n = 2$ and $s_2 = 1$, then Proposition 4.19 yields $\ell(s_1, s_2, t_1, \dots, t_m) = \ell(t_1, \dots, t_m) + 1 = \ell(t_1, \dots, t_m) + \ell(s_1, s_2)$. If $n = 2$ and $s_2 \geq 2$, then Proposition 4.19 yields $\ell(s_1, s_2, t_1, \dots, t_m) = \ell(s_2 - 1, t_1, \dots, t_m) + 1$, which is equal to $\ell(t_1, \dots, t_m) + 2$ by induction hypothesis. This achieves the proof since $\ell(s_1, s_2) = 2$ by Proposition 4.19.

If $n \geq 3$, then Proposition 4.19 yields

$$\begin{aligned} \ell(s_1, \dots, s_n, t_1, \dots, t_m) &= \begin{cases} \ell(s_2 - 1, s_3, \dots, s_n, t_1, \dots, t_m) + 1 & \text{if } s_2 \geq 2, \\ \ell(s_3, \dots, s_n, t_1, \dots, t_m) + 1 & \text{if } s_2 = 1. \end{cases} \\ \ell(s_1, \dots, s_n) &= \begin{cases} \ell(s_2 - 1, s_3, \dots, s_n) + 1 & \text{if } s_2 \geq 2, \\ \ell(s_3, \dots, s_n) + 1 & \text{if } s_2 = 1, \end{cases} \end{aligned}$$

so the result follows by induction. \square

Corollary 4.23. *Let $n \geq 2$ be an even integer and let (s_1, \dots, s_n) be a sequence of positive integers such that either $s_1 \geq 2$ or $s_1 = \dots = s_n = 1$. Then, the ordered element $M = M(s_1, \dots, s_n) \in \mathrm{SL}_2(\mathbb{Z})$ satisfies*

$$\mathfrak{d}_{\mathrm{lgth}}(M) = \mathrm{lgth}(M).$$

Proof. If $s_1 = 2$, then Lemma 4.22 yields $\mathrm{lgth}(M^m) = m \cdot \mathrm{lgth}(M)$ for each $m \geq 1$, which yields $\mathfrak{d}_{\mathrm{lgth}}(M) = \mathrm{lgth}(M)$.

If $s_1 = \dots = s_n = 1$, then $M^m = ((LR)^n)^m$. It then suffices to show that $\mathrm{lgth}((LR)^n) = n$ for each $n \geq 1$. For $n = 1$, this is directly given by Proposition 4.19. For $n \geq 1$, we also apply Proposition 4.19 and get $\mathrm{lgth}((LR)^n) = \mathrm{lgth}((LR)^{n-1}) + 1$, which yields the result by induction. \square

Remark 4.24. Note that $M(s_1, \dots, s_n)$ is conjugate in $\mathrm{SL}_2(\mathbb{Z})$ to $M(s_2, \dots, s_n, s_1)$. Hence, each element $M(s_1, \dots, s_n)$ admits a conjugate which satisfies the hypotheses of Corollary 4.23.

The general case is provided by the next result.

Proposition 4.25. *Let $M \in \mathrm{SL}_2(\mathbb{Z})$.*

- (1) *If $\mathrm{trace}(M) \in \{0, \pm 1\}$ then M has order $m \in \{3, 4, 6\}$ so $\mathfrak{d}_{\mathrm{lgth}}(M) = 0$.*
- (2) *If $\mathrm{trace}(M) \in \{\pm 2\}$ then M is conjugate to $\pm \begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix}$ for some $a \in \mathbb{Z}$, so $\mathfrak{d}_{\mathrm{lgth}}(M) = 0$.*
- (3) *If $|\mathrm{trace}(M)| \geq 3$ then $\pm M$ is conjugate to an ordered element M' , so $\mathfrak{d}_{\mathrm{lgth}}(M) = \mathfrak{d}_{\mathrm{lgth}}(M')$ is a positive integer.*

Proof. Writing $\lambda = \text{trace}(M)$, the characteristic polynomial of M is equal to $\chi_M = X^2 - \lambda X + 1$. If $\lambda \in \{0, \pm 1\}$, we obtain then orders 3, 4, 6, yielding (1). If $\lambda = \pm 2$, then $\chi_M = (X \pm 1)^2$, so there is an eigenvector of eigenvalue ± 1 , which can be chosen in \mathbb{Z}^2 with coprime coefficients. This yields (2).

In case (3) we can replace M with $-M$ if needed and assume that $\lambda \geq 3$. We will then show that M is conjugate to an ordered matrix. The fact that $\lambda = \text{trace}(M) \geq 3$ implies that M has distinct positive real eigenvalues μ, μ^{-1} with $\max\{\mu, \mu^{-1}\} = \frac{\lambda + \sqrt{\lambda^2 - 4}}{2} > 2$. Write $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$. The two eigenspaces are spanned by two vectors $(1, \xi_1)$ and $(1, \xi_2)$, where ξ_1, ξ_2 are nonzero reals (note that $(1, 0)$ and $(0, 1)$ are not eigenvectors of M since $bc \neq 0$).

(a) If $\xi_1 \xi_2 < 0$, then we may assume without loss of generality that $\xi_1 > 0$ and $\xi_2 < 0$ (by exchanging the names of ξ_1 and ξ_2). Up to replacing M with M^{-1} , we may furthermore assume that we have $\mu > 1$, where we have used the two following facts:

- (i) We have $\text{trace}(M^{-1}) = \text{trace}(M)$;
- (ii) The inverse of an ordered matrix is conjugate to an ordered matrix since the matrix $P = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ satisfies $PR^{-1}P^{-1} = L$ and $PL^{-1}P^{-1} = R$.

Then, we have $\begin{pmatrix} a + b\xi_1 = \mu \\ c + d\xi_1 = \mu\xi_1 \\ a + b\xi_2 = \mu^{-1} \\ c + d\xi_2 = \mu^{-1}\xi_2 \end{pmatrix}$ which yields $\begin{pmatrix} b(\xi_1 - \xi_2) = \mu - \mu^{-1} \\ c(\xi_1^{-1} - \xi_2^{-1}) = \mu - \mu^{-1} \\ a = \mu^{-1} - b\xi_2 \\ d = \mu^{-1} - \xi_2^{-1}c \end{pmatrix}$ which in

turn proves that $b, c > 0$, and then $a, d > 0$. Therefore, M belongs to the monoid generated by L and R (Lemma 4.14). As $\text{trace}(M) \neq 2$, M is not conjugate to a matrix of the form L^s or R^s for some $s \in \mathbb{Z}$ and is thus conjugate to an element which starts with L and ends with R , hence to an ordered element (Lemma 4.14). The result then follows from Corollary 4.23.

(b) If there exists $s \in \mathbb{Z}$ such that $(\xi_1 + s)(\xi_2 + s) < 0$, we conjugate M with L^s . This replaces ξ_i with $\xi_i + s$ and reduces to case (a).

(c) If (b) is not possible, there exists $s \in \mathbb{Z}$ such that $0 < \xi_i + s < 1$ for $i = 1, 2$. Replacing s with $s - 1$ if needed, we can rather assume that $|\xi_i + s| < 1$ for both i and that $|\xi_i + s| < \frac{1}{2}$ for at least one i , which we will assume to be 1 (by exchanging the names of ξ_1 and ξ_2). Therefore, by conjugating M with L^s we may assume that $|\xi_1| < \frac{1}{2}$ and $|\xi_2| < 1$. We then conjugate M with $\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$. This replaces ξ_i with $\xi'_i = -\frac{1}{\xi_i}$ for $i = 1, 2$ and we have $|\xi'_1 - \xi'_2| = |\xi_1 - \xi_2| \cdot \left|\frac{1}{\xi_1}\right| \cdot \left|\frac{1}{\xi_2}\right| > 2 \cdot |\xi_1 - \xi_2|$. After finitely many such steps we obtain $|\xi_1 - \xi_2| > 1$, which then gives case (b). \square

Corollary 4.26. *For each $M \in \text{GL}_2(\mathbb{Z})$, the following conditions are equivalent:*

- (1) We have $\mathfrak{d}_{\text{lgth}}(M) = \frac{1}{2}$.
- (2) The matrix M is conjugate in $\text{GL}_2(\mathbb{Z})$ to $\pm \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}$.
- (3) We have $\det(M) = -1$ and $\text{trace}(M) \in \{\pm 1\}$.

Proof. (2) \Rightarrow (3) is obvious.

(3) \Rightarrow (1). If M satisfies (3), we have $\text{trace}(M^2) = (\text{trace}(M))^2 - 2\det(M) = 3$, so that M^2 is conjugate to an ordered element of trace 3 by Proposition 4.25(3). The unique ordered element of trace 3 being the matrix $\begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} = LR = M(1, 1)$ of dynamical length 1, this proves (1).

(1) \Rightarrow (2). If M satisfies (1), we necessarily have $\det(M) = -1$ and $\mathfrak{d}_{\text{lgth}}(M^2) = 1$ (Proposition 4.25). By Proposition 4.25 and up to conjugation of M into $\text{GL}_2(\mathbb{Z})$, there exists an element $\varepsilon \in \{\pm 1\}$ such that $M' := \varepsilon M^2$ is an ordered matrix satisfying $\mathfrak{d}_{\text{lgth}}(M') = \text{lgth}(M') = 1$. This yields the existence of an integer $s \geq 1$ such that $M' = LR^s = \begin{pmatrix} 1 & s \\ 1 & s+1 \end{pmatrix}$. Writing $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$, we obtain

$$M^2 = \begin{pmatrix} a^2 + bc & b(a+d) \\ c(a+d) & d^2 + bc \end{pmatrix} = \varepsilon \begin{pmatrix} 1 & s \\ 1 & s+1 \end{pmatrix}.$$

The equality $c(a+d) = \varepsilon$ gives us $\text{trace}(M) = a+d = \pm 1$, so that (as above) $\text{trace}(M^2) = (\text{trace}(M))^2 - 2\det(M) = 3$. This implies $\varepsilon(s+2) = 3$, so that $\varepsilon = s = 1$. This proves that $b = c = a+d \in \{\pm 1\}$ and since $a^2 + bc = 1$, we finally obtain $a = 0$ and $b = c = d \in \{\pm 1\}$, proving that $M = \pm \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}$. \square

4.8. Dynamical length of regularisable elements and the proof of Theorem 2.

Recall the two following definitions:

Definition 4.27. An element $f \in \text{Bir}(\mathbb{P}^2)$ is said to be *regularisable* if there exists a birational map $\eta: X \dashrightarrow \mathbb{P}^2$, where X is a smooth projective surface, such that $\eta^{-1} \circ f \circ \eta \in \text{Aut}(X)$. By [BlaDés2015, Theorem B], this is equivalent to $\mu(f) = 0$ (where μ denotes the dynamical number of base-points, as explained in the introduction).

Definition 4.28. An element $f \in \text{Bir}(\mathbb{P}^2)$ is said to be *loxodromic* if $\log(\lambda(f)) > 0$ (where $\lambda(f) = \lim_{n \rightarrow \infty} (\deg(f^n))^{1/n}$ is the dynamical degree of f , as explained in the introduction).

It follows from [Giz1980, DF2001, BlaDés2015] that a Cremona transformation $f \in \text{Bir}(\mathbb{P}^2)$ belongs to exactly one of the five following categories:

- (1) **Algebraic elements** ;
- (2) **Jonquières twists:** $f \in \text{Bir}(\mathbb{P}^2)$ is a *Jonquières twist* if the sequence $n \mapsto \deg(f^n)$ grows linearly, i.e. if the sequence $n \mapsto \frac{\deg(f^n)}{n}$ admits a nonzero limit when n goes to infinity. Equivalently, f preserves a rational fibration $\mathbb{P}^2 \dashrightarrow \mathbb{P}^1$ and is not algebraic;
- (3) **Halphen twists:** $f \in \text{Bir}(\mathbb{P}^2)$ is a *Halphen twist* if the sequence $n \mapsto \deg(f^n)$ grows quadratically, i.e. if the sequence $n \mapsto \frac{\deg(f^n)}{n^2}$ admits a nonzero limit when n goes to infinity. Equivalently, f preserves an elliptic fibration $\mathbb{P}^2 \dashrightarrow \mathbb{P}^1$ and is not algebraic;
- (4) **Regularisable loxodromic elements:** In this case, $\lambda(f)$ is a Salem number or a reciprocal quadratic integer (see [DF2001, BlaCan2016]);
- (5) **Non-regularisable loxodromic elements:** In this case, $\lambda(f)$ is a Pisot number by [BlaCan2016].

If f is a Halphen twist or a regularisable loxodromic element, we will prove that the dynamical length $\mathfrak{d}_{\text{lgth}}(f)$ is positive in Corollary 4.32 and Proposition 4.34. In Lemma 4.11, an example of non-regularisable loxodromic element $f \in \text{Bir}(\mathbb{P}^2)$ whose dynamical length $\mathfrak{d}_{\text{lgth}}(f)$ is positive was given. These results are summarised in Figure 1 and achieve the proof of Theorem 2.

The following result follows from the proof of [BlaCan2016, Lemma5.10].

Lemma 4.29. *Let $f_1, f_2 \in W_\infty$ be two elements such that $\text{Base}(f_1) \cup \text{Base}(f_2)$ contains at most 9 points. Then,*

$$\sqrt{\deg(f_1 \circ f_2^{-1})} \leq \sqrt{\deg(f_1)} + \sqrt{\deg(f_2)}$$

Lemma 4.30. *Let $\psi \in \text{Bir}(\mathbb{P}^2) \setminus \text{Aut}(\mathbb{P}^2)$ be a birational map, and let r be its number of base-points. Recall that we have $r \geq 3$ by Lemma 3.5. Then, the following hold:*

- (1) $\text{lgth}(\psi) \geq \frac{\log(\deg(\psi))}{\log(\frac{r+1}{2})}$.
- (2) If $r \leq 9$, then $\text{lgth}(\psi) \geq \sqrt{\frac{\deg(\psi)}{5}}$.

Proof. We prove the result by induction on $\text{lgth}(\psi)$. When $\text{lgth}(\psi) = 1$, then ψ is a Jonquières transformation of degree $d = \deg(\psi) > 1$, which implies that $r = 2 \deg(\psi) - 1$. Hence, (1) is an equality and (2) holds, as $r \leq 9$ yields $\deg(\psi) \leq 5$.

Suppose now that $\text{lgth}(\psi) > 1$, and let $\varphi \in \text{Bir}(\mathbb{P}^2)$ be a Jonquières element such that $\psi' = \psi \circ \varphi$ is a predecessor of ψ (which implies in particular that $\text{lgth}(\psi') = \text{lgth}(\psi) - 1$). We then have $\text{Base}(\varphi^{-1}) \subseteq \text{Base}(\psi)$ (Lemma 1.3), which yields $\text{Base}(\psi'^{-1}) \subseteq \text{Base}(\psi^{-1})$ (Corollary 2.25). This proves that φ and ψ' have at most r base-points. In particular, we have $\deg(\varphi) \leq \frac{r+1}{2}$.

To prove (1) we start with $\deg(\psi) \leq \deg(\psi') \cdot \deg(\varphi) \leq \deg(\psi') \cdot \frac{r+1}{2}$, which yields $\log(\deg(\psi)) \leq \log(\deg(\psi')) + \log(\frac{r+1}{2})$. Applying induction to ψ' , we then get $\frac{\log(\deg(\psi))}{\log(\frac{r+1}{2})} \leq \frac{\log(\deg(\psi'))}{\log(\frac{r+1}{2})} + 1 \leq \text{lgth}(\psi') + 1 = \text{lgth}(\psi)$, as desired.

To prove (2), we denote by $\pi: Z \rightarrow \mathbb{P}^2$ the blow-up of the base-points of φ^{-1} . As $\text{Base}(\varphi^{-1}) \subseteq \text{Base}(\psi)$, there is a birational morphism $\epsilon: X \rightarrow Z$, such that $\pi \circ \epsilon$ is the blow-up of the base-points of ψ . We then get a commutative diagram

$$\begin{array}{ccccc}
 & & X & & \\
 & & \downarrow \epsilon & \searrow \eta' & \\
 & & Z & & \\
 \eta \swarrow & & \downarrow \pi & & \searrow \\
 \mathbb{P}^2 & \xleftarrow{\varphi} & \mathbb{P}^2 & \xrightarrow{\psi} & \mathbb{P}^2 \\
 & \dashleftarrow \psi' & & & \dashrightarrow
 \end{array}$$

where η, η' are the blow-ups of the base-points of φ and ψ^{-1} respectively. As η' blows-up r points, the same holds for $\eta \circ \epsilon$. Hence, we find that $\text{Base}(\psi') \cup \text{Base}(\varphi) \subseteq \text{Base}((\eta \circ \epsilon)^{-1})$ contains at most $r \leq 9$ points. We can then apply Lemma 4.29, which yields $\sqrt{\deg(\psi)} = \sqrt{\deg(\psi' \circ \varphi^{-1})} \leq \sqrt{\deg(\psi')} + \sqrt{\deg(\varphi)} \leq \sqrt{\deg(\psi')} + \sqrt{5}$. Applying induction to ψ' , we find

$$\sqrt{\frac{\deg(\psi)}{5}} \leq \sqrt{\frac{\deg(\psi')}{5}} + 1 \leq \text{lgth}(\psi') + 1 = \text{lgth}(\psi). \quad \square$$

Proposition 4.31. *Let $\pi: X \rightarrow \mathbb{P}^2$ be a birational morphism which is the blow-up of at most 9 points and let $\varphi \in \pi \operatorname{Aut}(X)\pi^{-1} \subseteq \operatorname{Bir}(\mathbb{P}^2)$ be a Cremona transformation. Then, the sequence $n \mapsto \frac{\deg(\varphi^n)}{n^2}$ admits a limit $L \in \mathbb{R}$ when n goes to infinity and we have*

$$\mathfrak{d}_{\text{lgth}}(\varphi) \geq \sqrt{\frac{L}{5}}.$$

Proof. Denote by $\Delta \subseteq \mathcal{B}(\mathbb{P}^2)$ the set of points blown-up by π . For each $n \in \mathbb{Z}$, we have $\operatorname{Base}(\varphi^n) \subseteq \Delta$. In particular we have $\sqrt{\deg(\varphi^{m+n})} = \sqrt{\deg(\varphi^m \circ (\varphi^{-n})^{-1})} \leq \sqrt{\deg(\varphi^m)} + \sqrt{\deg(\varphi^{-n})} = \sqrt{\deg(\varphi^m)} + \sqrt{\deg(\varphi^n)}$ for all $m, n \geq 1$ (Lemma 4.29). This means that the sequence $n \mapsto \sqrt{\deg(\varphi^n)}$ is subadditive, so $\lim_{n \rightarrow \infty} \frac{\sqrt{\deg(\varphi^n)}}{n}$ exists, which is equivalent to saying that $\lim_{n \rightarrow \infty} \frac{\deg(\varphi^n)}{n^2}$ exists.

For each $n \geq 1$, the number of base-points of φ^n is at most 9. This yields $\text{lgth}(\varphi^n) \geq \sqrt{\frac{\deg(\varphi^n)}{5}}$ (Lemma 4.30(2)), whence $\frac{\text{lgth}(\varphi^n)}{n} \geq \sqrt{\frac{\deg(\varphi^n)}{5n^2}}$ and the result follows by taking the limit when n goes to infinity. \square

Corollary 4.32. *If $\varphi \in \operatorname{Bir}(\mathbb{P}^2)$ is a birational transformation such that $(\deg(\varphi^n))_{n \geq 1}$ grows quadratically (i.e. φ is a Halphen twist), then $\mathfrak{d}_{\text{lgth}}(\varphi) \geq \sqrt{\frac{1}{5} \cdot \lim_{n \rightarrow \infty} \frac{\deg(\varphi^n)}{n^2}} > 0$.*

Proof. A birational transformation of \mathbb{P}^2 has a quadratic growth if and only if it is conjugate to an automorphism of a Halphen surface, obtained by blowing-up 9 points of \mathbb{P}^2 [Giz1980]. The result then follows from Proposition 4.31. \square

Remark 4.33. Using an analogue method as in [BlaDés2015, Proposition 5.1] we are able to give a uniform lower bound $C > 0$ such that $\mathfrak{d}_{\text{lgth}}(\varphi) \geq C$ for all Halphen twists. However, this bound is far from being reached from the known examples.

Proposition 4.34. *Let $\varphi \in \operatorname{Bir}(\mathbb{P}^2)$ be a loxodromic birational map which is regularisable, i.e. such that there exists a birational map $\kappa: \mathbb{P}^2 \dashrightarrow X$ that conjugates φ to an automorphism $g = \kappa \circ \varphi \circ \kappa^{-1} \in \operatorname{Aut}(X)$ where X is a smooth projective surface.*

Then, each X as above is isomorphic to the blow-up of finitely many points $p_1, \dots, p_r \in \mathcal{B}(\mathbb{P}^2)$ with $r \geq 10$, and the dynamical length of φ satisfies $\mathfrak{d}_{\text{lgth}}(\varphi) \geq \frac{\log(\lambda(\varphi))}{\log(\frac{r+1}{2})} > 0$.

Proof. We first show that there exists a birational morphism $\eta: X \rightarrow \mathbb{P}^2$. Suppose the converse, for contradiction. Then, [Har1987, Corollary 1.2] implies that the action of $\operatorname{Aut}(X)$ on $\operatorname{Pic}(X)$ is finite (i.e. factorises through the action of a finite group). This is impossible: the dynamical degree of g , equal to the one of φ as both are conjugate, is the spectral radius of the action of g on $\operatorname{Pic}(X) \otimes_{\mathbb{Z}} \mathbb{C}$ (see the introductions of [DF2001] and [BlaCan2016] for details on these two facts). This dynamical degree is therefore equal to 1, contradicting the fact that φ is loxodromic.

We then obtain the existence of a birational morphism $\eta: X \rightarrow \mathbb{P}^2$ which is the blow-up of finitely many points $p_1, \dots, p_r \in \mathcal{B}(\mathbb{P}^2)$. Observe moreover that $r \geq 10$ because φ is loxodromic. One way to see this classical fact is to use Proposition 4.31 which implies that $\{\deg(\tilde{\varphi}^n)\}_{n \geq 1}$ grows at most quadratically if $r \leq 9$, where $\tilde{\varphi} = \eta \circ g \circ \eta^{-1}$. This is impossible since $\{\deg(\varphi^n)\}_{n \geq 1}$ grows exponentially and because $\varphi, \tilde{\varphi} \in \operatorname{Bir}(\mathbb{P}^2)$ are conjugate (by $\eta \circ \kappa \in \operatorname{Bir}(\mathbb{P}^2)$).

We then replace φ with its conjugate $\eta \circ g \circ \eta^{-1} \in \text{Bir}(\mathbb{P}^2)$. After this, we get $\varphi^n = \eta \circ g^n \circ \eta^{-1}$ for each $n \geq 1$, so $\text{Base}(\varphi^n) \subseteq \{p_1, \dots, p_r\}$. By Lemma 4.30(1) we have $\text{lgth}(\varphi^n) \geq \frac{\log(\deg(\varphi^n))}{\log(\frac{r+1}{2})}$. From $\lim_{n \rightarrow \infty} (\deg(\varphi^n))^{1/n} = \lambda(\varphi)$, we deduce $\lim_{n \rightarrow \infty} \frac{\log(\deg(\varphi^n))}{n} = \log(\lambda(\varphi))$, which then yields

$$\mathfrak{d}_{\text{lgth}}(\varphi) = \lim_{n \rightarrow \infty} \frac{\text{lgth}(\varphi^n)}{n} \geq \lim_{n \rightarrow \infty} \frac{\log(\deg(\varphi^n))}{n \log(\frac{r+1}{2})} = \frac{\log(\lambda(\varphi))}{\log(\frac{r+1}{2})} > 0$$

as desired. □

5. LOWER SEMICONTINUITY OF THE LENGTH: THE PROOF OF THEOREM 3

Throughout this section, $\Pi: \mathbb{P}^2 \dashrightarrow \mathbb{P}^1$ will denote the standard linear projection

$$\begin{array}{ccc} \mathbb{P}^2 & \dashrightarrow & \mathbb{P}^1 \\ [x : y : z] & \mapsto & [x : y]. \end{array}$$

5.1. Variables. The proof of Theorem 3 uses the notion of variables, that we now define.

Definition 5.1. A rational map $v: \mathbb{P}^2 \dashrightarrow \mathbb{P}^1$ is called a *variable*, if there exists a birational map $f \in \text{Bir}(\mathbb{P}^2)$ such that $\Pi \circ f = v$.

$$\begin{array}{ccc} \mathbb{P}^2 & \xrightarrow{f} & \mathbb{P}^2 \\ & \searrow v & \downarrow \Pi \\ & & \mathbb{P}^1 \end{array}$$

Writing a variable $v: \mathbb{P}^2 \dashrightarrow \mathbb{P}^1$ as $[x : y : z] \dashrightarrow [v_0(x, y, z) : v_1(x, y, z)]$ where $v_0, v_1 \in k[x, y, z]$ are homogeneous of the same degree, without common factor, we define its degree as the common degree of v_0 and v_1 (which is also the degree of a general fibre of v).

Remark 5.2. Let us make the following observations:

- (1) A rational map $v: \mathbb{P}^2 \dashrightarrow \mathbb{P}^1$ is a variable if and only if there exists a rational map $w: \mathbb{P}^2 \dashrightarrow \mathbb{P}^1$ such that the rational map $(v, w): \mathbb{P}^2 \dashrightarrow \mathbb{P}^1 \times \mathbb{P}^1$ is birational.
- (2) Writing a rational map $v: \mathbb{P}^2 \dashrightarrow \mathbb{P}^1$ as $[x : y : z] \dashrightarrow [v_0(x, y, z) : v_1(x, y, z)]$ where $v_0, v_1 \in k[x, y, z]$ are homogeneous of the same degree d , then v is a variable if and only if there exist an integer $D \geq d$ and homogeneous polynomials $h, v_2 \in k[x, y, z]$ of degrees $D - d$ and D , such that $[x : y : z] \dashrightarrow [hv_0 : hv_1 : v_2]$ is an element of $\text{Bir}(\mathbb{P}^2)$.
- (3) For each $p \in \mathbb{P}^2$, every projection $\pi_p: \mathbb{P}^2 \dashrightarrow \mathbb{P}^1$ away from p is a variable of degree 1. Conversely, all variables of degree 1 are obtained like this.
- (4) The group $\text{Bir}(\mathbb{P}^2)$ acts transitively on the set of variables by right-composition. Similarly, $\text{Aut}(\mathbb{P}^1)$ acts on the set of variables by left-composition.

Definition 5.3. Let $v: \mathbb{P}^2 \dashrightarrow \mathbb{P}^1$ be a variable. We define the length of v , written $\text{lgth}(v)$, to be the minimum of the lengths of the birational maps $\varphi \in \text{Bir}(\mathbb{P}^2)$ such that $\Pi \circ \varphi = v$.

Lemma 5.4. *Let $v: \mathbb{P}^2 \dashrightarrow \mathbb{P}^1$ be a variable. For each $\varphi \in \text{Bir}(\mathbb{P}^2)$ such that $\Pi \circ \varphi = v$, we have*

$$\text{lgth}(\varphi) \in \{\text{lgth}(v), \text{lgth}(v) + 1\}.$$

Proof. By definition, there exists $\psi \in \text{Bir}(\mathbb{P}^2)$ such that $\Pi \circ \psi = v$ and $\text{lgth}(\psi) = \text{lgth}(v)$. Since $\Pi \circ \varphi \circ (\psi)^{-1} = \Pi$, the map $\varphi \circ (\psi)^{-1}$ is a Jonquières transformation, which implies that the lengths of φ and ψ differ at most by one. As $\text{lgth}(v) \leq \text{lgth}(\varphi)$ by definition, we get the result. \square

Lemma 5.5. *Let $v: \mathbb{P}^2 \dashrightarrow \mathbb{P}^1$ be a variable, and let $\theta: \mathbb{P}^1 \rightarrow \mathbb{P}^1$ be a morphism. Then, the following are equivalent:*

- (1) *The rational map $\theta \circ v: \mathbb{P}^2 \dashrightarrow \mathbb{P}^1$ is a variable.*
- (2) *The morphism $\theta: \mathbb{P}^1 \rightarrow \mathbb{P}^1$ is an automorphism.*

Proof. (1) \Rightarrow (2): If $\theta \circ v$ is a variable, its general scheme theoretic fibre is irreducible, so that the general scheme theoretic fibre of θ also. This proves that θ has degree 0 or 1. As $\theta \circ v$ is non-constant, so is θ , which is thus an automorphism.

(2) \Rightarrow (1): If θ is an automorphism of \mathbb{P}^1 , we have already noted in Remark 5.2(4) that $\theta \circ v$ is a variable. \square

Lemma 5.6. *Let $f: \mathbb{P}^2 \dashrightarrow \mathbb{P}^2$ be a non-dominant and non-constant rational map and let $v: \mathbb{P}^2 \dashrightarrow \mathbb{P}^1$ be a variable. Then, the following are equivalent:*

- (1) *There exists a morphism $\kappa: \mathbb{P}^1 \rightarrow \mathbb{P}^2$ such that $\kappa \circ v = f$.*
- (2) *For each linear projection $\pi: \mathbb{P}^2 \dashrightarrow \mathbb{P}^1$, there exists a morphism $\theta: \mathbb{P}^1 \rightarrow \mathbb{P}^1$ such that $\pi \circ f = \theta \circ v$.*
- (3) *There exists a linear projection $\pi: \mathbb{P}^2 \dashrightarrow \mathbb{P}^1$ and a non-constant morphism $\theta: \mathbb{P}^1 \rightarrow \mathbb{P}^1$ such that $\pi \circ f = \theta \circ v$.*

Proof. As f is non-constant, $\pi \circ f: \mathbb{P}^2 \dashrightarrow \mathbb{P}^1$ is a well-defined rational map for each linear projection $\pi: \mathbb{P}^2 \dashrightarrow \mathbb{P}^1$.

(1) \Rightarrow (2): For each linear projection $\pi: \mathbb{P}^2 \dashrightarrow \mathbb{P}^1$ we get $\pi \circ f = (\pi \circ \kappa) \circ v$.

(2) \Rightarrow (3): Since f is not constant, there is a linear projection $\pi: \mathbb{P}^2 \dashrightarrow \mathbb{P}^1$ such that $\pi \circ f$ is not constant.

(3) \Rightarrow (1): Let us choose $\alpha \in \text{Aut}(\mathbb{P}^2)$, $\beta \in \text{Aut}(\mathbb{P}^1)$, and $g \in \text{Bir}(\mathbb{P}^2)$ such that $\beta \circ \pi \circ \alpha = v \circ g = \Pi$. We then replace (π, f, θ, v) with $(\beta \circ \pi \circ \alpha, \alpha^{-1} \circ f \circ g, \beta \circ \theta, v \circ g)$ and can assume that $\pi = v = \Pi$.

We write locally the non-constant morphism $\theta: \mathbb{P}^1 \rightarrow \mathbb{P}^1$ as $[1 : t] \dashrightarrow [1 : r(t)]$ for some $r(t) \in k(t) \setminus k$. The equation $\Pi \circ f = \theta \circ \Pi$ implies that f is locally given as

$$[1 : t : u] \dashrightarrow [1 : r(t) : s(t, u)]$$

for some rational function $s \in k(t, u)$. As f is not dominant, the two elements $s(t, u)$ and $r(t)$ are algebraically dependent over k . Since $k(t)$ is algebraically closed in $k(t, u)$, this shows that $s \in k(t)$. We can thus write f as

$$[x : y : z] \dashrightarrow [f_0(x, y) : f_1(x, y) : f_2(x, y)]$$

for some homogeneous polynomials $f_0, f_1, f_2 \in k[x, y]$, and can choose $\kappa: \mathbb{P}^1 \rightarrow \mathbb{P}^2$ to be $[u : v] \mapsto [f_0(u, v) : f_1(u, v) : f_2(u, v)]$. \square

5.2. Definition of the Zariski topology on $\text{Bir}(\mathbb{P}^2)$ and basic properties. Following [Dem1970, Ser2010], the notion of families of birational maps is defined, and used in Definition 5.8 for describing the natural Zariski topology on $\text{Bir}(X)$.

Definition 5.7. Let A, X be irreducible algebraic varieties, and let f be a A -birational map of the A -variety $A \times X$, inducing an isomorphism $U \rightarrow V$, where U, V are open subsets of $A \times X$, whose projections on A are surjective.

The rational map f is given by $(a, x) \dashrightarrow (a, p_2(f(a, x)))$, where p_2 is the second projection, and for each k -point $a \in A$, the birational map $x \dashrightarrow p_2(f(a, x))$ corresponds to an element $f_a \in \text{Bir}(X)$. The map $a \mapsto f_a$ represents a map from A (more precisely from the k -points of A) to $\text{Bir}(X)$, and will be called a *morphism* from A to $\text{Bir}(X)$.

Definition 5.8. A subset $F \subseteq \text{Bir}(X)$ is closed in the Zariski topology if for any algebraic variety A and any morphism $A \rightarrow \text{Bir}(X)$ the preimage of F is closed.

If d is a positive integer, we set $\text{Bir}(\mathbb{P}^2)_d := \{f \in \text{Bir}(\mathbb{P}^2), \deg(f) \leq d\}$. We will use the following result, which is [BlaFur2013, Proposition 2.10]:

Lemma 5.9. *A subset $F \subseteq \text{Bir}(\mathbb{P}^2)$ is closed if and only if $F \cap \text{Bir}(\mathbb{P}^2)_d$ is closed in $\text{Bir}(\mathbb{P}^2)_d$ for any d .*

The aim of this whole section 5 is to prove that for each nonnegative integer ℓ the set

$$\text{Bir}(\mathbb{P}^2)^\ell := \{f \in \text{Bir}(\mathbb{P}^2), \text{lgth}(f) \leq \ell\}$$

is closed in $\text{Bir}(\mathbb{P}^2)$. By Lemma 5.9, this is equivalent to proving that $\text{Bir}(\mathbb{P}^2)_d^\ell := \text{Bir}(\mathbb{P}^2)_d \cap \text{Bir}(\mathbb{P}^2)^\ell$ is closed in $\text{Bir}(\mathbb{P}^2)_d$ for any d . We will now describe the topology of $\text{Bir}(\mathbb{P}^2)_d$. A convenient way to handle this topology is through the map $\pi_d: \mathfrak{Bir}(\mathbb{P}^2)_d \rightarrow \text{Bir}(\mathbb{P}^2)_d$ that we introduce in the next definition and whose properties are given in Lemma 5.13 below.

Let us now fix the integer $d \geq 1$. We will constantly use the following piece of notation:

Definition 5.10. Denote by $\mathfrak{Rat}(\mathbb{P}^2)_d$ the projective space associated with the vector space of triples (f_0, f_1, f_2) where $f_0, f_1, f_2 \in k[x, y, z]$ are homogeneous polynomials of degree d . The equivalence class of (f_0, f_1, f_2) will be denoted by $[f_0 : f_1 : f_2]$.

For each $f = [f_0 : f_1 : f_2] \in \mathfrak{Rat}(\mathbb{P}^2)_d$, we denote by ψ_f the rational map $\mathbb{P}^2 \dashrightarrow \mathbb{P}^2$ defined by

$$[x_0 : x_1 : x_2] \mapsto [f_0(x_0, x_1, x_2) : f_1(x_0, x_1, x_2) : f_2(x_0, x_1, x_2)].$$

Writing $\text{Rat}(\mathbb{P}^2)$ the set of rational maps from \mathbb{P}^2 to \mathbb{P}^2 and setting $\text{Rat}(\mathbb{P}^2)_d := \{h \in \text{Rat}(\mathbb{P}^2), \deg(h) \leq d\}$, we obtain a surjective map

$$\Psi_d: \mathfrak{Rat}(\mathbb{P}^2)_d \rightarrow \text{Rat}(\mathbb{P}^2)_d, \quad f \mapsto \psi_f.$$

This map induces a surjective map $\pi_d: \mathfrak{Bir}(\mathbb{P}^2)_d \rightarrow \text{Bir}(\mathbb{P}^2)_d$, where $\mathfrak{Bir}(\mathbb{P}^2)_d$ is defined to be $\Psi_d^{-1}(\text{Bir}(\mathbb{P}^2)_d)$.

Remark 5.11. For each field extension $k \subseteq k'$, we can similarly associate to each $f \in \mathfrak{Rat}(\mathbb{P}^2)_d(k')$ a rational transformation $\psi_f: \mathbb{P}^2_{k'} \dashrightarrow \mathbb{P}^2_{k'}$ defined over k' . This will be needed in the sequel to use a valuative criterium.

We will need the following result:

Proposition 5.12. *The set $\mathfrak{Bir}(\mathbb{P}^2)_d$ is locally closed in $\mathfrak{Rat}(\mathbb{P}^2)_d$ and thus inherits from $\mathfrak{Rat}(\mathbb{P}^2)_d$ the structure of an algebraic variety. Moreover, the following assertions hold:*

- (1) *For each $f \in \overline{\mathfrak{Bir}(\mathbb{P}^2)_d} \setminus \mathfrak{Bir}(\mathbb{P}^2)_d$, the rational map $\psi_f: \mathbb{P}^2 \dashrightarrow \mathbb{P}^2$ is non-dominant.*

- (2) For each field extension $k \subseteq k'$, the set $\mathfrak{Bir}(\mathbb{P}^2)_d(k')$ of k' -points of $\mathfrak{Bir}(\mathbb{P}^2)_d$ is equal to the set $\{f \in \mathfrak{Rat}(\mathbb{P}^2)_d(k'), \psi_f: \mathbb{P}_{k'}^2 \dashrightarrow \mathbb{P}_{k'}^2 \text{ is birational}\}$.

Proof. Even if the first assertion is [BlaFur2013, Lemma 2.4(2)], we recall the argument since this one will be used to prove the rest of the proposition. Denote by $F \subseteq \mathfrak{Rat}(\mathbb{P}^2)_d \times \mathfrak{Rat}(\mathbb{P}^2)_d$ the closed algebraic variety corresponding to pairs $([g_0 : g_1 : g_2], [f_0 : f_1 : f_2])$ such that the “formal composition”

$$g \circ f = [h_0 : h_1 : h_2] = [g_0(f_0, f_1, f_2) : g_1(f_0, f_1, f_2) : g_2(f_0, f_1, f_2)]$$

is a “multiple” (maybe zero) of the identity. This corresponds to ask that $h_0y = h_1x$, $h_0z = h_2x$, $h_1z = h_2y$. We then define $F_0 \subseteq F$ to be the closed subset such that the formal composition is zero. If $(g, f) \in F_0$, let us observe that the formal composition $g \circ f$ is zero, so that the rational map $\psi_f: \mathbb{P}^2 \dashrightarrow \mathbb{P}^2$ is non-dominant. Conversely, if $(g, f) \in F \setminus F_0$, the formal composition $g \circ f$ is non-zero, so that the rational map $\psi_f: \mathbb{P}^2 \dashrightarrow \mathbb{P}^2$ is birational.

The second projection $\text{pr}_2: \mathfrak{Rat}(\mathbb{P}^2)_d \times \mathfrak{Rat}(\mathbb{P}^2)_d \rightarrow \mathfrak{Rat}(\mathbb{P}^2)_d$ yields two closed subvarieties

$$G_0 = \text{pr}_2(F_0) \subseteq G = \text{pr}_2(F) \subseteq \mathfrak{Rat}(\mathbb{P}^2)_d.$$

By what has been said above, ψ_f is non-dominant when $f \in G_0$ and birational when $f \in G \setminus G_0$. It follows that $(G \setminus G_0) \subseteq \mathfrak{Bir}(\mathbb{P}^2)_d$. Since $\deg(\varphi^{-1}) = \deg(\varphi)$ for each $\varphi \in \text{Bir}(\mathbb{P}^2)$ (Lemma 2.20(3)), we even get the equality $\mathfrak{Bir}(\mathbb{P}^2)_d = G \setminus G_0$. This shows that $\mathfrak{Bir}(\mathbb{P}^2)_d$ is locally closed in $\mathfrak{Rat}(\mathbb{P}^2)_d$, and also gives (1). To obtain (2), we observe that the construction made in the proof is defined over k' , and that the inverse of any birational transformation of \mathbb{P}^2 defined over k' is still defined over k' . \square

The following result, which is [BlaFur2013, Corollary 2.9], will be crucial for us since it provides us a bridge from the “weird” topological space $\text{Bir}(\mathbb{P}^2)_d$ to the “nice” topological space $\mathfrak{Bir}(\mathbb{P}^2)_d$ which is an algebraic variety.

Lemma 5.13. *The map $\pi_d: \mathfrak{Bir}(\mathbb{P}^2)_d \rightarrow \text{Bir}(\mathbb{P}^2)_d$ is continuous and closed. In particular, it is a quotient topological map: A subset $F \subseteq \text{Bir}(\mathbb{P}^2)_d$ is closed if and only if its preimage $\pi_d^{-1}(F)$ is closed.*

Recall that our aim is to prove that $\text{Bir}(\mathbb{P}^2)_d^\ell = \{f \in \text{Bir}(\mathbb{P}^2)_d, \text{lgth}(f) \leq \ell\}$ is closed in $\text{Bir}(\mathbb{P}^2)_d$. By Lemma 5.13, this reduces to prove that $\mathfrak{Bir}(\mathbb{P}^2)_d^\ell := \pi_d^{-1}(\text{Bir}(\mathbb{P}^2)_d^\ell)$ is closed in (the algebraic variety) $\mathfrak{Bir}(\mathbb{P}^2)_d$.

We conclude this section by noting that the conjunction of Lemmas 5.9 and 5.13 gives us the following usefull characterisation (already contained in [BlaFur2013, Corollary 2.7]) of closed subsets of $\text{Bir}(\mathbb{P}^2)$:

Lemma 5.14. *A subset $F \subseteq \text{Bir}(\mathbb{P}^2)$ is closed if and only if $\pi_d^{-1}(F \cap \text{Bir}(\mathbb{P}^2)_d) \subseteq \mathfrak{Bir}(\mathbb{P}^2)_d$ is closed for any d .*

5.3. The use of a valuative criterion. Let us set

$$R := k[[t]], \quad \text{and} \quad K := k((t)).$$

We will also write $\overline{K} = \bigcup_{a \geq 1} k((t^{1/a}))$ since this latter field is an algebraic closure of K by Newton-Puiseux theorem [Rui1993, Proposition 4.4].

Definition 5.15. Let $n \geq 1$ be an integer and let $a = a(t) = [a_0 : \cdots : a_n] \in \mathbb{P}^n(K)$ be a K -point of the n -th projective space \mathbb{P}^n . Then, up to multiplying (a_0, \dots, a_n) with some power of t , we may assume that all coefficients a_i belong to R and that the evaluation $(a_0(0), \dots, a_n(0))$ at $t = 0$ is nonzero. This enables us to define non ambiguously the element $a(0) \in \mathbb{P}^n$, also denoted $\lim_{t \rightarrow 0} a(t)$, by

$$a(0) = \lim_{t \rightarrow 0} a(t) := [a_0(0) : \cdots : a_n(0)].$$

Remark 5.16. More generally, if X is a complete k -variety and $x = x(t) \in X(K)$ is a K -point of X , one can define $x(0) = \lim_{t \rightarrow 0} x(t) \in X$ in the following way: The morphism $x: \text{Spec}(K) \rightarrow X$ admits a unique factorisation under the form $x = \tilde{x} \circ \iota$ where $\iota: \text{Spec}(K) \hookrightarrow \text{Spec}(R)$ is the open immersion induced by the natural injection $R \hookrightarrow K$ and where $\tilde{x}: \text{Spec}(R) \rightarrow X$ is a k -morphism (see the valuative criterion of properness given in [Har1977, (II, Theorem 4.7), page 101]).

The following valuative criterion is classical, see e.g. [MFK1994, chap. 2, §1, pp 52-54]. We refer to [Fur2009] for a proof in characteristic zero and to [Bla2016] for a proof in any characteristic.

Lemma 5.17. *Let $\varphi: X \rightarrow Y$ be a morphism between algebraic k -varieties, X being quasi-projective, and Y being projective. Let y_0 be a (closed) point of Y . Then, the two following assertions are equivalent:*

- (1) *We have $y_0 \in \overline{\varphi(X)}$;*
- (2) *There exists a K -point $x = x(t) \in X(K)$ such that the K -point $y = y(t) := \varphi(x(t)) \in Y(K)$ satisfied $y_0 = y(0)$.*

Remark 5.18. Lemma 5.17 is analogue to the case of a continuous map $\varphi: X \rightarrow Y$ between metric spaces where a point y_0 of Y belongs to $\overline{\varphi(X)}$ if and only if there exists a sequence $(x_n)_{n \geq 1}$ of X such that $y_0 = \lim_{n \rightarrow +\infty} \varphi(x_n)$.

Remark 5.19. Applying Definition 5.15 to an element $f = f(t) = [f_0 : f_1 : f_2] \in \mathfrak{Rat}(\mathbb{P}^2)_d(K)$ allows us to define $f(0) \in \mathfrak{Rat}(\mathbb{P}^2)_d$. If we assume furthermore that $f \in \mathfrak{Bir}(\mathbb{P}^2)_d(K) \subseteq \mathfrak{Rat}(\mathbb{P}^2)_d(K)$, note that $f(0)$ necessarily belongs to $\overline{\mathfrak{Bir}(\mathbb{P}^2)_d}$ by Lemma 5.17, so that $\psi_{f(0)}: \mathbb{P}^2 \rightarrow \mathbb{P}^2$ is either birational or non-dominant by Proposition 5.12. Let us recall for clarity that for any $f \in \mathfrak{Rat}(\mathbb{P}^2)_d(K)$, we have defined a K -rational transformation $\psi_f: \mathbb{P}^2_K \dashrightarrow \mathbb{P}^2_K$ in Remark 5.11, and that this transformation is moreover birational if we assume that $f \in \mathfrak{Bir}(\mathbb{P}^2)_d(K)$ by Proposition 5.12(2).

We will prove that $\mathfrak{Bir}(\mathbb{P}^2)_d^\ell$ is closed in $\mathfrak{Bir}(\mathbb{P}^2)_d$. For this, we will prove that its closure $\overline{\mathfrak{Bir}(\mathbb{P}^2)_d^\ell}$ in $\mathfrak{Rat}(\mathbb{P}^2)_d$ is such that $\overline{\mathfrak{Bir}(\mathbb{P}^2)_d^\ell} \cap \mathfrak{Bir}(\mathbb{P}^2)_d = \mathfrak{Bir}(\mathbb{P}^2)_d^\ell$. We begin with the following result which is just a (technical) application of the valuative criterium given above. If k' is an extension field of k , $\text{Aut}_{k'}(\mathbb{P}^2) \simeq \text{PGL}_3(k')$, resp. $\text{Bir}_{k'}(\mathbb{P}^2)$, denotes the group of automorphisms, resp. birational transformations, of \mathbb{P}^2 defined over k' . Actually, we will only consider the cases where $k' = K$ (since we will use the valuative criterium given in Lemma 5.17) and where $k' = \overline{K}$ (since we need an algebraically closed field in order to apply the machinery about the length that we have developped).

Proposition 5.20. *For any $h \in \overline{\mathfrak{Bir}(\mathbb{P}^2)_d}^\ell$ there exists $f \in \mathfrak{Bir}(\mathbb{P}^2)_d(K)$ such that $h = f(0)$, and such that the birational map $\psi_f \in \text{Bir}_{\overline{K}}(\mathbb{P}^2)$ associated to $f \in \mathfrak{Bir}(\mathbb{P}^2)_d(\overline{K})$ has length at most ℓ .*

The proof of Proposition 5.20 relies on the two following lemmas:

Lemma 5.21. *For each $p \in \mathbb{P}^2$, the set $\mathfrak{Jonq}_{p,d} := \Psi_d^{-1}(\text{Jonq}_p \cap \text{Bir}(\mathbb{P}^2)_d)$ is closed in $\mathfrak{Bir}(\mathbb{P}^2)_d$.*

Proof. Up to applying an automorphism of \mathbb{P}^2 , we may assume that $p = [0 : 0 : 1]$. Denote by \mathfrak{L} the projective space (of dimension 3) associated with the vector space of pairs (g_0, g_1) where $g_0, g_1 \in k[x, y]$ are homogeneous polynomials of degree 1. The equivalence class of (g_0, g_1) will be denoted by $[g_0 : g_1]$. Denote by $Y \subseteq \mathfrak{Bir}(\mathbb{P}^2)_d \times \mathfrak{L}$ the closed subvariety given by elements $([f_0 : f_1 : f_2], [g_0 : g_1])$ satisfying $f_0 g_1 = f_1 g_0$. Since the first projection $\text{pr}_1 : \mathfrak{Bir}(\mathbb{P}^2)_d \times \mathfrak{L} \rightarrow \mathfrak{Bir}(\mathbb{P}^2)_d$ is a closed morphism, the lemma follows from the equality $\mathfrak{Jonq}_{p,d} = \text{pr}_1(Y)$. \square

Remark 5.22. Lemma 5.21 asserts that $\mathfrak{Jonq}_{p,d} = \pi_d^{-1}(\text{Jonq}_p \cap \text{Bir}(\mathbb{P}^2)_d)$ is closed in $\mathfrak{Bir}(\mathbb{P}^2)_d$ for each d . By Lemma 5.14, this means that Jonq_p is closed in $\text{Bir}(\mathbb{P}^2)$.

Lemma 5.23. *Any Cremona transformation $g \in \text{Bir}(\mathbb{P}^2)$ of length ℓ admits an expression of the form*

$$g = a_1 \circ \varphi_1 \circ \cdots \circ a_\ell \circ \varphi_\ell \circ a_{\ell+1},$$

where $a_1, \dots, a_{\ell+1} \in \text{Aut}(\mathbb{P}^2)$, $\varphi_1, \dots, \varphi_\ell \in \text{Jonq}_p$, and $\deg(\varphi_i) \leq \deg(g)$ for each i .

Proof. This follows from Theorem 1 and the fact that if φ is a Jonquières transformation such that $g \circ \varphi$ is a predecessor of g , then $\text{Base}(\varphi^{-1}) \subseteq \text{Base}(g)$ (Lemma 1.3(3)), which implies that g has at least $2 \deg(\varphi) - 1$ base-points, so $\deg(g) \geq \deg(\varphi)$ (every element of $\text{Bir}(\mathbb{P}^2)$ of degree $d \geq 2$ has at most $2d - 1$ points, and equality holds if and only if the map is Jonquières [BCM2015, Lemma 13]). \square

Proof of Proposition 5.20. In order to use Lemma 5.17, we realise $\mathfrak{Bir}(\mathbb{P}^2)_d^\ell$ as the image of a morphism of algebraic varieties. Let us fix $p = [0 : 0 : 1] \in \mathbb{P}^2$. By Lemma 5.23, an element f of $\mathfrak{Bir}(\mathbb{P}^2)_d$ belongs to $\mathfrak{Bir}(\mathbb{P}^2)_d^\ell$ if and only if the birational transformation ψ_f admits an expression of the form

$$\psi_f = a_1 \circ \varphi_1 \circ \cdots \circ a_\ell \circ \varphi_\ell \circ a_{\ell+1},$$

where $a_1, \dots, a_{\ell+1} \in \text{Aut}(\mathbb{P}^2)$, $\varphi_1, \dots, \varphi_\ell \in \text{Jonq}_p$, and $\deg \varphi_i \leq d$ for each i .

We now use the closed subvariety $\mathfrak{Jonq}_{p,d} \subseteq \mathfrak{Bir}(\mathbb{P}^2)_d$ given in Lemma 5.21. Define the product $\mathfrak{P} := (\text{PGL}_3)^{\ell+1} \times (\mathfrak{Jonq}_{p,d})^\ell$ and let $\mathbf{Comp} : \mathfrak{P} \rightarrow \mathfrak{Bir}(\mathbb{P}^2)_{d^\ell}$ be the formal composition morphism defined by

$$(a_i)_{1 \leq i \leq \ell+1} \times (\varphi_i)_{1 \leq i \leq \ell} \mapsto a_1 \circ \varphi_1 \circ \cdots \circ a_\ell \circ \varphi_\ell \circ a_{\ell+1}.$$

Let $\Delta \subseteq \mathfrak{Bir}(\mathbb{P}^2)_d \times \mathfrak{Bir}(\mathbb{P}^2)_{d^\ell}$ be the pseudo-diagonal, i.e. the set of pairs (f, g) such that $\psi_f = \psi_g$. Being given by the equations $f_i g_j = f_j g_i$, for all i, j , where $f = [f_0 : f_1 : f_2] \in \mathfrak{Bir}(\mathbb{P}^2)_d$, $g = [g_0 : g_1 : g_2] \in \mathfrak{Bir}(\mathbb{P}^2)_{d^\ell}$, the set Δ is closed into $\mathfrak{Bir}(\mathbb{P}^2)_d \times \mathfrak{Bir}(\mathbb{P}^2)_{d^\ell}$.

Denote by $\text{id} \times \mathbf{Comp} : \mathfrak{Bir}(\mathbb{P}^2)_d \times \mathfrak{P} \rightarrow \mathfrak{Bir}(\mathbb{P}^2)_d \times \mathfrak{Bir}(\mathbb{P}^2)_{d^\ell}$, the morphism sending (f, p) to $(f, \mathbf{Comp}(p))$ and by Δ' the closed variety defined by

$$\Delta' := (\text{id} \times \mathbf{Comp})^{-1}(\Delta) \subseteq \mathfrak{Bir}(\mathbb{P}^2)_d \times \mathfrak{P}.$$

By what has been said above, we have $\mathfrak{Bir}(\mathbb{P}^2)_d^\ell = \text{pr}_1(\Delta')$ where $\text{pr}_1: \mathfrak{Bir}(\mathbb{P}^2)_d \times \mathfrak{P} \rightarrow \mathfrak{Bir}(\mathbb{P}^2)_d$ is the first projection. Setting $\varphi = \iota \circ \text{pr}_1$ where $\iota: \mathfrak{Bir}(\mathbb{P}^2)_d \hookrightarrow \mathfrak{Nat}(\mathbb{P}^2)_d$ is the natural injection, we also have $\mathfrak{Bir}(\mathbb{P}^2)_d^\ell = \varphi(\Delta')$.

Since $h \in \overline{\varphi(\Delta')}$, Lemma 5.17 yields us the existence of an element $(f, p) = (f(t), p(t)) \in \Delta'(K) \subseteq \mathfrak{Bir}(\mathbb{P}^2)_d(K) \times \mathfrak{P}(K)$ such that $h = f(0)$. We observe that the birational map $\psi_f \in \text{Bir}_{\overline{K}}(\mathbb{P}^2)$ associated to $f \in \mathfrak{Bir}(\mathbb{P}^2)_d(\overline{K})$ has length at most ℓ . \square

5.4. The end of the proof of Theorem 3. The main result of the previous section (Proposition 5.20) asserts that any element $h \in \overline{\mathfrak{Bir}(\mathbb{P}^2)_d^\ell}$ is equal to $f(0)$ for a certain element $f \in \mathfrak{Bir}(\mathbb{P}^2)_d(K)$ such that the length of $\psi_f \in \text{Bir}_{\overline{K}}(\mathbb{P}^2)$ is at most ℓ . The main technical result of the present section is Proposition 5.25 which establishes that the limit $\psi_{f(0)}: \mathbb{P}^2 \dashrightarrow \mathbb{P}^2$ is either birational of length $\leq \ell$ or non-dominant (however, in the non-dominant case, we need to prove a stronger statement in order to make an induction). This information is sufficient for showing that $\overline{\mathfrak{Bir}(\mathbb{P}^2)_d^\ell}$ is closed in $\mathfrak{Bir}(\mathbb{P}^2)_d$ thus proving Theorem 3. We begin with the following simple lemma to be used in the proof of Proposition 5.25.

Lemma 5.24. *Let V be a finite dimensional vector space over k and let $u, v \in K \otimes_k V$ be two vectors such that*

- (1) *The vectors u, v are linearly independent over K ;*
- (2) *The vector v belongs to $R \otimes_k V$ and its evaluation $v(0)$ at $t = 0$ is nonzero.*

Then, there exist $\alpha, \beta \in K$ such that:

- (1) *The vector $\tilde{u} := \alpha u + \beta v$ belongs to $R \otimes_k V$;*
- (2) *The vectors $\tilde{u}(0)$ and $v(0)$ are linearly independent over k .*

Proof. Let us complete the vector $e_1 := v(0)$ in a basis e_1, \dots, e_n of V . Decomposing the vectors u, v in this basis, we obtain expressions

$$u = \sum_i u_i e_i, \quad v = \sum_i v_i e_i,$$

where $u_1, \dots, u_n \in K$, $v_1, \dots, v_n \in R$ and $v_1(0) = 1, v_i(0) = 0$ for $i = 2, \dots, n$. The vector $w := u - \frac{u_1}{v_1}v$ is nonzero and admits an expression

$$w = \sum_i w_i e_i,$$

where $w_1, \dots, w_n \in K$ and $w_1 = 0$. Let j be the unique integer such that the vector $\tilde{w} := t^j w$ belongs to $R \otimes_k V$ and its evaluation $\tilde{w}(0)$ at $t = 0$ is nonzero. The vectors $\tilde{w}(0)$ and $v(0) = e_1$ are linearly independent over k because $\tilde{w}(0) \in (ke_2 \oplus \dots \oplus ke_n) \setminus \{0\}$. Since $\tilde{w} = t^j(u - \frac{u_1}{v_1}v)$, it is enough to set $\alpha = t^j$ and $\beta = -t^j \frac{u_1}{v_1}$. \square

Proposition 5.25. *Let $f = f(t) \in \mathfrak{Bir}(\mathbb{P}^2)_d(K) \subseteq \mathfrak{Bir}(\mathbb{P}^2)_d(\overline{K})$ be an element such that the associated birational map $\psi_f \in \text{Bir}_{\overline{K}}(\mathbb{P}^2)$ has length $\ell \geq 0$ and denote by $f(0) \in \overline{\mathfrak{Bir}(\mathbb{P}^2)_d} \subseteq \mathfrak{Nat}(\mathbb{P}^2)_d$ the evaluation of $f = f(t)$ at $t = 0$ (see Definition 5.15 and Remark 5.19). Then, the following implications are satisfied:*

- (1) *If $f(0) \in \overline{\mathfrak{Bir}(\mathbb{P}^2)_d}$, then the birational map $\psi_{f(0)}$ is of length $\leq \ell$.*
- (2) *If $f(0) \in \overline{\mathfrak{Bir}(\mathbb{P}^2)_d} \setminus \mathfrak{Bir}(\mathbb{P}^2)_d$, then the rational map $\psi_{f(0)}$ is equal to $\kappa \circ v$ for some variable $v: \mathbb{P}^2 \dashrightarrow \mathbb{P}^1$ of length $\leq \ell$ and some morphism $\kappa: \mathbb{P}^1 \rightarrow \mathbb{P}^2$.*

In particular, for each linear projection $\pi: \mathbb{P}^2 \dashrightarrow \mathbb{P}^1$, the composition $\pi \circ \psi_{f(0)}: \mathbb{P}^2 \dashrightarrow \mathbb{P}^1$ is either not defined or equal to $\rho \circ v$ for some variable $v: \mathbb{P}^2 \dashrightarrow \mathbb{P}^1$ of length $\leq \ell$ and some endomorphism $\rho: \mathbb{P}^1 \rightarrow \mathbb{P}^1$.

Proof. We prove the result by induction on ℓ .

Case of length $\ell = 0$. The equality $\ell = 0$ corresponds exactly to ask that $\psi_f \in \text{Aut}_K(\mathbb{P}^2)$. We write $f = [ha_0 : ha_1 : ha_2]$, where $h \in K[x, y, z]$ is homogeneous of degree $d - 1$ and $[a_0 : a_1 : a_2] \in \mathfrak{Bir}(\mathbb{P}^2)_1(K) \simeq \text{PGL}_3(K)$. We can moreover assume that the coefficients of h belong to $R \subseteq K$ and that the evaluation $h(0)$ of h at $t = 0$ is non-zero. Similarly, we can assume that a_0, a_1, a_2 have coefficients in R and that at least one of these has a non-zero value at $t = 0$. The element $[a_0(0) : a_1(0) : a_2(0)] \in \mathfrak{Nat}(\mathbb{P}^2)_1$ corresponds to a 3×3 -matrix. If the matrix is of rank 3, the element $f(0) \in \mathfrak{Bir}(\mathbb{P}^2)_d$ corresponds to a linear automorphism $\psi_{f(0)} \in \text{Bir}(\mathbb{P}^2)$ of length 0. If the matrix is of rank 2, then $\psi_{f(0)}$ admits a decomposition under the form $\kappa \circ v$ where $\kappa: \mathbb{P}^1 \rightarrow \mathbb{P}^2$ is a linear morphism and $v: \mathbb{P}^2 \dashrightarrow \mathbb{P}^1$ is a linear variable, i.e. of degree 1. The last case is when the matrix has rank 1, which corresponds to the case where $\psi_{f(0)}: \mathbb{P}^2 \rightarrow \mathbb{P}^2$ is the constant map to some point $a \in \mathbb{P}^2$. Let $\kappa: \mathbb{P}^1 \rightarrow \mathbb{P}^2$ be the constant map to a and let v be any variable of length 0, then we have $\psi_{f(0)} = \kappa \circ v$.

Case of length $\ell \geq 1$.

This implies that the birational transformation $(\psi_f)^{-1} \in \text{Bir}_{\overline{K}}(\mathbb{P}^2)$ admits at least one base-point. The proof is divided into the following steps:

Reduction to the case where all base-points are defined over K :

By assumption, the birational map $\psi_f \in \text{Bir}_{\overline{K}}(\mathbb{P}^2)$ has length ℓ . Replacing t with $t^{\frac{1}{a}}$ for some $a \geq 1$, we can thus assume that all base-points of $(\psi_f)^{-1}$ are defined over K .

Denote by $p \in \mathbb{P}^2(K)$ a base-point of maximal multiplicity of $(\psi_f)^{-1}$.

Reduction to the case where $p = [0 : 0 : 1]$.

Write $p = [p_0 : p_1 : p_2]$ where each p_i belongs to K . Up to multiplying (p_0, p_1, p_2) by t^i for some well chosen (and unique) integer i , we may assume that $p_0, p_1, p_2 \in R$ and that $p_i(0) \neq 0$ for some i . Let us choose coefficients b_{ij} in the field k such that the following matrix has nonzero determinant:

$$M = \begin{pmatrix} b_{00} & b_{01} & p_0(0) \\ b_{10} & b_{11} & p_1(0) \\ b_{20} & b_{21} & p_2(0) \end{pmatrix}.$$

In other words, we have $M \in \text{GL}_3(k)$. This implies that the matrix

$$B(t) = \begin{pmatrix} b_{00} & b_{01} & p_0 \\ b_{10} & b_{11} & p_1 \\ b_{20} & b_{21} & p_2 \end{pmatrix} \in \text{Mat}_3(R)$$

is invertible in $\text{Mat}_3(R)$ (because its determinant is invertible). The evaluation at $t = 0$ of the corresponding automorphism $\beta \in \text{Aut}_K(\mathbb{P}^2) = \text{PGL}_3(K)$ is the element of $\text{PGL}_3(k) = \text{GL}_3(k)/k^*$ given by the class of the matrix $M \in \text{GL}_3(k)$. We can replace f with $\tilde{f} = \beta^{-1} \circ f \in B'_d(K)$ (formal composition), because we have $\tilde{f}(0) = \beta(0)^{-1} \circ f(0)$, where $\beta(0)$ belongs to $\text{PGL}_3(k)$. After this change, the point p is equal to $[0 : 0 : 1] \in \mathbb{P}^2 \subseteq \mathbb{P}^2(K)$.

As in Definition 5.15 (see also Remark 5.19), we write $f = [f_0 : f_1 : f_2]$ where the components $f_i \in R[x, y, z]$ satisfy $(f_0(0), f_1(0), f_2(0)) \neq (0, 0, 0)$.

If $(f_0(0), f_1(0)) = (0, 0)$, then $\psi_{f(0)}$ is the constant map to p : Hence we have $f(0) \notin \mathfrak{Bir}(\mathbb{P}^2)_d$ and the result is trivially true by taking $\kappa: \mathbb{P}^1 \rightarrow \mathbb{P}^2$ the constant map to p and $v: \mathbb{P}^2 \dashrightarrow \mathbb{P}^1$ any variable of length $\leq \ell$. We can thus assume that $(f_0(0), f_1(0)) \neq (0, 0)$, which means that $\psi_{f(0)}$ is not the constant map to p , and can consider the rational map $\Pi \circ \psi_{f(0)}: \mathbb{P}^2 \dashrightarrow \mathbb{P}^1$, given by $[x : y : z] \dashrightarrow [f_0(0)(x, y, z) : f_1(0)(x, y, z)]$. We achieve the proof by studying two cases, depending on whether this rational map is constant or not.

Case A: The rational map $\Pi \circ \psi_{f(0)}$ is not constant – construction of an element of length $\ell - 1$.

Since $p = [0 : 0 : 1]$ is a base-point of maximal multiplicity of $(\psi_f)^{-1} \in \text{Bir}_{\overline{K}}(\mathbb{P}^2)$, by Corollary 3.28 there exists an element $\varphi \in \text{Jonq}_p(\overline{K})$ (i.e. an element of $\text{Bir}_{\overline{K}}(\mathbb{P}^2)$ which preserves the pencil of lines through p) such that $\varphi \circ \psi_f \in \text{Bir}_{\overline{K}}(\mathbb{P}^2)$ is of length $\ell - 1$ and of degree smaller than $\deg(\psi_f) \leq d$.

We may moreover assume that φ satisfies the two following assertions:

- (i) φ is defined over K , i.e. $\varphi \in \text{Jonq}_p(K)$.
- (ii) φ preserves a general line through p , i.e. $\Pi \circ \varphi = \Pi$.

To obtain (i), we could use the fact that all base-points of φ are defined over K (since they are included into the base-points of $(\psi_f)^{-1}$). Alternatively, we can use the same trick as above: Since φ is defined over $k((t^{1/a}))$ for some integer $a \geq 1$, it is enough to replace t with $t^{1/a}$.

To obtain (ii), it is enough to note that any element $\varphi \in \text{Jonq}_p$ may be written as a composition $\alpha \circ \tilde{\varphi}$ where $\alpha \in \text{Aut}(\mathbb{P}^2) \cap \text{Jonq}_p$ and $\tilde{\varphi} \in \text{Jonq}_p$ preserves a general line through p .

Let $g \in \mathfrak{Bir}(\mathbb{P}^2)_d(K)$ be such that $\psi_g = \varphi \circ \psi_f$. Note that the assumption (ii) above shows us that $\Pi \circ \psi_f = \Pi \circ \psi_g$. As before, write $g = [g_0 : g_1 : g_2]$ where the components $g_i \in R[x, y, z]$ satisfy $(g_0(0), g_1(0), g_2(0)) \neq (0, 0, 0)$. The fact that $\psi_{f(0)}$ is not the constant map to p corresponds exactly to saying that $(f_0(0), f_1(0)) \neq (0, 0)$. Replacing φ with its composition with $[x : y : z] \mapsto [t^{-i}x : t^{-i}y : z]$ for some well chosen integer $i \geq 0$ we may replace (g_0, g_1, g_2) with $(t^{-i}g_0, t^{-i}g_1, g_2)$ and then assume that $(g_0(0), g_1(0)) \neq (0, 0)$. We obtain then a rational map

$$\begin{aligned} \nu: \quad \mathbb{P}^2 &\dashrightarrow \mathbb{P}^1 \\ [x : y : z] &\mapsto [f_0(0)(x, y, z) : f_1(0)(x, y, z)] = [g_0(0)(x, y, z) : g_1(0)(x, y, z)] \end{aligned}$$

which satisfies $\nu = \Pi \circ \psi_{f(0)} = \Pi \circ \psi_{g(0)}$ and is thus non-constant by hypothesis.

Applying the induction hypothesis to g (and $g(0)$), the map $\nu = \Pi \circ \psi_{g(0)} = \Pi \circ \psi_{f(0)}$ is equal to $\theta \circ v$, where $v: \mathbb{P}^2 \dashrightarrow \mathbb{P}^1$ is a variable of length at most $\ell - 1$ and $\theta: \mathbb{P}^1 \rightarrow \mathbb{P}^1$ is an endomorphism. Moreover, θ is non-constant since ν is non-constant.

a) If $\psi_{f(0)}$ is a birational map, then $\nu = \Pi \circ \psi_{f(0)}$ is a variable. Since $\nu = \theta \circ v$, this implies that $\theta \in \text{Aut}(\mathbb{P}^1)$ (Lemma 5.5) and thus that $\text{lgth}(\nu) = \text{lgth}(v) \leq \ell - 1$. In particular, $\text{lgth}(\psi_{f(0)}) \leq \ell$ since $\nu = \Pi \circ \psi_{f(0)}$ (Lemma 5.4) as we wanted.

b) If $\psi_{f(0)}$ is not a birational map, then it is non-dominant (see Remark 5.19). The equality $\nu = \Pi \circ \psi_{f(0)} = \theta \circ v$ yields the existence of a morphism $\kappa: \mathbb{P}^1 \rightarrow \mathbb{P}^2$ such that

$\kappa \circ v = \psi_{f(0)}$ (Lemma 5.6). This achieves the proof in this case.

Case B: The rational map $\Pi \circ \psi_{f(0)}$ is constant.

Let $[\lambda : \mu] \in \mathbb{P}^1$ be the constant value of the map $\Pi \circ \psi_{f(0)}$. There exists a homogeneous polynomial $h \in k[x, y, z]$ such that $(f_1(0), f_2(0)) = (\lambda h, \mu h)$. Up to replacing f with $\alpha \circ f$, where $\alpha \in \text{Aut}(\mathbb{P}^2)$ is of the form $[x : y : z] \mapsto [ax + by : cy + dy : z]$, we can assume that $(\lambda, \mu) = (0, 1)$, which implies that $f_0(0) = 0$ and $f_1(0) \neq 0$.

In this case, we have $f(0) \notin \mathfrak{Bir}(\mathbb{P}^2)_d$ and $\psi_{f(0)}$ is the rational map $[x : y : z] \dashrightarrow [0 : f_1(0)(x, y, z) : f_2(0)(x, y, z)]$. Writing $\pi : \mathbb{P}^2 \dashrightarrow \mathbb{P}^1, [x : y : z] \mapsto [y : z]$ the projection away from $[1 : 0 : 0]$, it remains to see that the rational map $\pi \circ \psi_{f(0)} : \mathbb{P}^2 \dashrightarrow \mathbb{P}^1, [x : y : z] \dashrightarrow [f_1(0)(x, y, z) : f_2(0)(x, y, z)]$ is the composition of a variable $\mathbb{P}^2 \dashrightarrow \mathbb{P}^1$ of length $\leq \ell$ and an endomorphism of \mathbb{P}^1 .

To show this, let us note that by Lemma 5.24, there exist $\alpha, \beta \in K$ such that $\tilde{f}_0 = \alpha f_0 + \beta f_1 \in R[x, y, z]$ and $\tilde{f}_0(0)$ does not belong to $k \cdot f_1(0)$.

We observe that the result holds for $\tilde{f} := [\tilde{f}_0 : f_1 : f_2] \in \mathfrak{Bir}(\mathbb{P}^2)_d(K)$. Indeed, $\psi_{\tilde{f}}$ and ψ_f only differ by an element of $\text{Aut}(\mathbb{P}^2)(K)$ that fixes $p = [0 : 0 : 1]$, so $\psi_{\tilde{f}}$ and ψ_f have the same length, p is a base-point of $\psi_{\tilde{f}}^{-1}$ of maximal multiplicity, and all base-points of $\psi_{\tilde{f}}^{-1}$ are defined over K . Moreover, \tilde{f} satisfies Case A. The result then holds for f , since $\pi \circ \psi_{\tilde{f}(0)} = \pi \circ \psi_{f(0)}$. \square

Proof of Theorem 3. We have already explained why Proposition 5.25 implies Theorem 3. Let us however summarise the proof. We want to show that $\text{Bir}(\mathbb{P}^2)^\ell = \{f \in \text{Bir}(\mathbb{P}^2), \text{lgth}(f) \leq \ell\}$ is closed in $\text{Bir}(\mathbb{P}^2)$ for each integer $\ell \geq 0$. By Lemma 5.14, this is equivalent to saying that $\mathfrak{Bir}(\mathbb{P}^2)_d^\ell = \pi_d^{-1}(\text{Bir}(\mathbb{P}^2)_d^\ell)$ is closed in $\mathfrak{Bir}(\mathbb{P}^2)_d$ for each d . This latter point directly follows from Propositions 5.20 and 5.25. \square

REFERENCES

- [Alb2002] MARIA ALBERICH-CARRAMIÑANA: *Geometry of the plane Cremona maps*, Lecture Notes in Mathematics **1769**. Springer-Verlag, Berlin, 2002. [1.1](#), [2.1](#), [2.44](#), [3.1](#), [3.18](#), [3.3](#), [3.22](#), [3.3](#)
- [Ale1916] JAMES W. ALEXANDER: *On the factorization of Cremona plane transformations*. Trans. Amer. Math. Soc. **17** (1916), no. 3, 295–300. [1.1](#)
- [BCM2015] CINZIA BISI, ALBERTO CALABRI, MASSIMILIANO MELLA: *On plane Cremona transformations of fixed degree*. J. Geom. Anal. **25** (2015), no. 2, 1108–1131. [1.4](#), [5.3](#)
- [Bla2011] JÉRÉMY BLANC: *Elements and cyclic subgroups of finite order of the Cremona group*. Comment. Math. Helv. **86** (2011), no. 2, 469–497. [1.1](#)
- [Bla2012] JÉRÉMY BLANC: *Simple Relations in the Cremona Group*. Proc. Amer. Math. Soc. **140** (2012), 1495–1500. [1.1](#)
- [Bla2016] JÉRÉMY BLANC: *Conjugacy classes of special automorphisms of the affine spaces*. Algebra Number Theory **10** (2016), no. 5, 939–967. [5.3](#)
- [BlaCal2016] JÉRÉMY BLANC, ALBERTO CALABRI: *On degenerations of plane Cremona transformations*. Math. Z. **282** (2016), no. 1-2, 223–245. [1.4](#), [2](#), [2.2](#), [2.44](#)
- [BlaCan2016] JÉRÉMY BLANC, SERGE CANTAT: *Dynamical degrees of birational transformations of projective surfaces*. J. Amer. Math. Soc. **29** (2016), no. 2, 415–471. [2](#), [2.2](#), [\(4\)](#), [\(5\)](#), [4.8](#), [4.8](#)
- [BlaDés2015] JÉRÉMY BLANC, JULIE DÉSSERTI: *Degree growth of birational maps of the plane*. Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) **14** (2015), no. 2, 507–533. [1.8](#), [1.3](#), [4.27](#), [4.8](#), [4.33](#)

- [BlaFur2013] JÉRÉMY BLANC, JEAN-PHILIPPE FURTER: *Topologies and structures of the Cremona groups*. Ann. of Math. **178** (2013), no. 3, 1173–1198. [1.8](#), [1.4](#), [5.2](#), [5.2](#), [5.2](#)
- [CC2018] SERGE CANTAT, YVES DE CORNULIER: *Distortion in Cremona groups*. Text in preparation. [1.3](#)
- [Cas1901] GUIDO CASTELNUOVO: *Le trasformazioni generatrici del gruppo cremoniano nel piano*. Atti della R. Accad. delle Scienze di Torino **36**, 861–874, 1901. [1.1](#), [3.3](#)
- [Cor2013] YVES DE CORNULIER: *The Cremona group is not an amalgam*. Appendix of *Normal subgroups in the Cremona group*. Acta Math. **210** (2013), no. 1, 31–94. [1.1](#)
- [Cor1995] ALESSIO CORTI: *Factoring birational maps of threefolds after Sarkisov*. J. Algebraic Geom. **4** (1995), no. 2, 223–254. [1.1](#)
- [Dem1970] MICHEL DEMAZURE: *Sous-groupes algébriques de rang maximum du groupe de Cremona*. Ann. Sci. École Norm. Sup. (4) **3** (1970), 507–588. [1.4](#), [5.2](#)
- [DF2001] JEFFREY DILLER, CHARLES FAVRE: *Dynamics of bimeromorphic maps of surfaces*. Amer. J. Math. **123** (2001), no. 6, 1135–1169. [4.8](#), [\(4\)](#), [4.8](#)
- [Dol2012] IGOR V. DOLGACHEV: *Classical algebraic geometry. A modern view*. Cambridge University Press, Cambridge, 2012. xii+639 pp. ISBN: 978-1-107-01765-8 [4.3](#), [4.3](#)
- [Fur2002] JEAN-PHILIPPE FURTER: *On the length of polynomial automorphisms of the affine plane*. Math. Ann. **322** (2002), no. 2, 401–411. [1.1](#), [1.4](#)
- [Fur2009] JEAN-PHILIPPE FURTER: *Plane polynomial automorphisms of fixed multidegree*. Math. Ann. **343** (2009), no. 4, 901–920. [5.3](#)
- [Giz1980] MARAT KH. GIZATULLIN: *Rational G-surfaces*. Izv. Akad. Nauk SSSR Ser. Mat. **44** (1980), no. 1, 110–144. [4.8](#), [4.8](#)
- [Giz1982] MARAT KH. GIZATULLIN: *Defining relations for the Cremona group of the plane*. Izv. Akad. Nauk SSSR Ser. Mat. **46** (1982), 909–970. [1.1](#)
- [Har1987] BRIAN HARBOURNE: *Rational surfaces with infinite automorphism group and no antipluricanonical curve*, Proc. Amer. Math. Soc. **99** (1987), no. 3, 409–414. [4.8](#)
- [Har1977] ROBIN HARTSHORNE: *Algebraic geometry*, Graduate Texts in Mathematics, no. 52, Springer-Verlag, New York-Heidelberg, 1977. [5.16](#)
- [Isk1985] VASILĬ A. ISKOVSKIKH: *Proof of a theorem on relations in a two-dimensional Cremona group*. Uspekhi Mat. Nauk **40** (1985), no. 5 (245), 255–256. English transl. in Russian Math. Surveys **40** (1985), no. 5, 231–232. [1.1](#)
- [Jun1942] HEINRICH JUNG: *Über ganze birationale Transformationen der Ebene*. J. Reine Angew. Math. **184**, (1942). 161–174. [1.1](#)
- [Lam2001] STÉPHANE LAMY: *L’alternative de Tits pour $\text{Aut}[C^2]$* . J. Algebra **239** (2001), no. 2, 413–437. [1.1](#)
- [Lam2002] STÉPHANE LAMY: *Une preuve géométrique du théorème de Jung*. Enseign. Math. (2) **48** (2002), no. 3–4, 291–315. [1.1](#)
- [Lon2018] ANNE LONJOU: *Graphes associés au groupe de Cremona* <https://arxiv.org/abs/1802.02910> ([document](#)), [1.1](#)
- [MFK1994] DAVID B. MUMFORD, JOHN FOGARTY, FRANCES C. KIRWAN: *Geometric invariant theory*. Third edition. Ergebnisse der Mathematik und ihrer Grenzgebiete (2), **34**. Springer-Verlag, Berlin, 1994. xiv+292 pp. [5.3](#)
- [Rui1993] JÉSUS M. RUIZ: *The basic theory of power series*. Advanced Lectures in Mathematics, Friedr. Vieweg & Sohn, Braunschweig, 1993, x+134 pp. [5.3](#)
- [Ser1980] JEAN-PIERRE SERRE: *Trees*. Translated from the French by John Stillwell. Springer-Verlag, Berlin-New York, 1980. ix+142 pp. [1.1](#)
- [Ser2010] JEAN-PIERRE SERRE: *Le groupe de Cremona et ses sous-groupes finis*. Séminaire Bourbaki. Volume 2008/2009. Astérisque No. **332** (2010), Exp. No. 1000, vii, 75–100. [1.4](#), [5.2](#)
- [vdK53] WOUTER VAN DER KULK: *On polynomial rings in two variables*. Nieuw Arch. Wisk. **1** (1953), 33–41. [1.1](#)
- [Wri1992] DAVID WRIGHT: *Two-dimensional Cremona Groups acting on simplicial Complexes*. Trans. Amer. Math. Soc., vol. **331**, no. 1, (1992). [1.1](#), [1.1](#)

JÉRÉMY BLANC, UNIVERSITÄT BASEL, SPIEGELGASSE 1, CH-4051 BASEL, SWITZERLAND.
E-mail address: `jeremy.blanc@unibas.ch`

JEAN-PHILIPPE FURTER, DPT. OF MATH., UNIV. OF LA ROCHELLE, AV. CRÉPEAU, 17000 LA
ROCHELLE, FRANCE
E-mail address: `jpfurter@univ-lr.fr`