

On birational transformations of pairs in the complex plane

Jérémy Blanc · Ivan Pan · Thierry Vust

Received: 4 April 2008 / Accepted: 8 August 2008 / Published online: 2 December 2008
© Springer Science+Business Media B.V. 2008

Abstract This article deals with the study of the birational transformations of the projective complex plane which leave invariant an irreducible algebraic curve. We try to describe the state of the art and provide some new results on this subject.

Keywords Birational transformation · Decomposition group · Inertia group

Mathematics Subject Classification (2000) 14E07 · 14J26 · 14H50

1 Introduction

1.1 The decomposition and inertia groups

We study the birational transformations of the projective complex plane which leave invariant an irreducible algebraic curve.

We denote by $\text{Bir}(\mathbb{P}^2)$ the group of birational transformations of the complex projective plane $\mathbb{P}^2 = \mathbb{P}^2(\mathbb{C})$: this is the *Cremona group* of \mathbb{P}^2 . If $C \subset \mathbb{P}^2$ is an irreducible curve and $\varphi \in \text{Bir}(\mathbb{P}^2)$, we say that φ *preserves* C (or *leaves* C *invariant*) if φ restricts to a birational transformation of C . If this transformation is the identity, we say that φ *fixes* C .

J. Blanc
Université de Grenoble I,
UFR de Mathématiques, UMR 5582 du CNRS, Institut Fourier, BP 74, 38402, Saint-Martin d'Hères,
France

I. Pan (✉)
Instituto de Matemática, UFRGS, av. Bento Gonçalves 9500, 91540-000, Porto Alegre, RS, Brasil
e-mail: pan.ivan@gmail.com

T. Vust
Section de mathématiques, Université de Genève, 2-4 rue du Lièvre, CP 64, 1211, Genève 4, Switzerland
e-mail: Thierry.Vust@unige.ch

Let $C \subset \mathbb{P}^2$ be an irreducible plane curve. Following Gizatullin [23], we introduce the *decomposition group* of C in $\text{Bir}(\mathbb{P}^2)$, denoted here by $\text{Dec}(\mathbb{P}^2, C) = \text{Dec}(C)$, as the group of Cremona transformations that preserve C . The action ρ of $\text{Dec}(C)$ on C induces a (not necessarily exact) complex

$$1 \longrightarrow \text{Ine}(C) \longrightarrow \text{Dec}(C) \xrightarrow{\rho} \text{Bir}(C) \longrightarrow 1, \tag{1}$$

where $\text{Ine}(C) = \text{Ine}(\mathbb{P}^2, C) := \ker(\rho)$ is the *inertia group* of C in $\text{Bir}(\mathbb{P}^2)$, which is the group of Cremona transformations that fix C .

1.2 Birational geometry of pairs

The above notions can be generalised to pairs (S, C) , where S is a surface and $C \subset S$ an irreducible curve. We say that a birational transformation $\varphi : S \dashrightarrow S'$ is a *birational transformation of pairs* $\varphi : (S, C) \dashrightarrow (S', C')$ if it restricts to a birational transformation $\varphi|_C : C \dashrightarrow C'$, and in this case we say that the two pairs are *birationally equivalent*. The group of birational transformations of a pair (S, C) is denoted by $\text{Dec}(S, C)$ and induces as before a complex

$$1 \longrightarrow \text{Ine}(S, C) \longrightarrow \text{Dec}(S, C) \xrightarrow{\rho} \text{Bir}(C) \longrightarrow 1, \tag{2}$$

which is exactly the complex (1) if $S = \mathbb{P}^2$ (and in this case we omit the surface in the notation). We will say that (2) is the *canonical complex* of the pair (S, C) . Note that $\text{Aut}(S, C) := \text{Aut}(S) \cap \text{Dec}(S, C)$ is the group of automorphisms of S that leave the curve C invariant.

1.3 Outline of the article

The aim of this article is to give a survey about the pairs (\mathbb{P}^2, C) whose decomposition group is not trivial, together with a description of their corresponding canonical complexes; we point out what is known to us about the subject and give some new results.

Sections 2, 3 and 4 deal respectively with curves of genus ≥ 2 , 1 and 0. Sections 5 and 6 express the link between the transformations that preserve or fix curves with respectively the classification of finite subgroups and the dynamics of the elements of $\text{Bir}(\mathbb{P}^2)$.

1.4 Conventions

In the sequel, $g(C)$ will denote the geometric genus of an irreducible curve C . Recall also that a *de Jonquières transformation* is a birational transformation of \mathbb{P}^2 that preserves a pencil of lines. Finally, all our surfaces are assumed to be rational, smooth, projective and irreducible.

2 Curves of genus at least equal to 2

2.1 The main tool: adjoint linear systems

Let $C \subseteq \mathbb{P}^2$ be an irreducible curve with $g(C) \geq 2$. To study the group $\text{Dec}(C)$, we follow an idea of Castelnuovo and Enriques which consists in considering the *adjoint* linear system associated to C : we take an embedded resolution of the singularities of C , say $\sigma : Y \rightarrow \mathbb{P}^2$,

denote by $\tilde{C} \subset Y$ the strict transform of C and consider the linear system $\sigma_*|K_Y + \tilde{C}|$ and its fixed part Δ . By definition, the adjoint system $\text{Adj}(C)$ is the linear system $\sigma_*|K_Y + \tilde{C}| - \Delta$. By Riemann-Roch it has dimension $g(C) - 1 > 0$. The main result is:

Proposition 2.1.1 ([11], [10, Prop. 2.5]) *Let $C \subset \mathbb{P}^2$ be an irreducible curve with $g(C) \geq 2$. If $\varphi \in \text{Bir}(\mathbb{P}^2)$ sends (\mathbb{P}^2, C) onto (\mathbb{P}^2, D) , then it sends $\text{Adj}(C)$ onto $\text{Adj}(D)$. In particular, the group $\text{Dec}(C) = \text{Dec}(\mathbb{P}^2, C)$ stabilizes the linear system $\text{Adj}(C)$.*

One can also define the adjoint of a linear system by taking the adjoint of a general member of the system. Since this construction decreases the degree of the curves, it has to stop after a finite number of iterations when the curves have no adjoint, i.e. when they have genus 0 or 1. This yields:

Proposition 2.1.2 ([10, Prop. 2.12]) *Let $C \subset \mathbb{P}^2$ be an irreducible curve, with $g(C) \geq 2$. There exists a linear system (resp. a pencil) of elliptic or rational curves Λ such that $\text{Dec}(C)$ (resp. $\text{Ine}(C)$) stabilizes Λ .*

2.2 The inertia group of curves of genus ≥ 2

Castelnuovo used the existence of the invariant pencils yielded by Proposition 2.1.2 to bound the order of elements of finite order of $\text{Ine}(C)$:

Theorem 2.2.1 ([11], [24, Chap.VIII, Sect. 2], [13, Book IV, Chap. VII, Sect. 3]) *Let $C \subset \mathbb{P}^2$ be an irreducible curve with $g(C) \geq 2$, and let $\varphi \in \text{Ine}(C)$, $\varphi \neq 1$. Then, either φ is conjugate to a de Jonquières transformation or φ has order 2, 3 or 4.*

In [10], an examination of the two possible cases of pencils yielded by Proposition 2.1.2 leads to a precise description of all pairs (\mathbb{P}^2, C) having a non-trivial inertia group. This generalises Castelnuovo’s theorem. We describe these cases in Examples 2.2.2, 2.2.3 and 2.2.4 below and then state the classification result (Theorem 2.2.5).

Example 2.2.2 ([1,6,13,14,24,27,36])

- (a) Let p_1, \dots, p_7 be 7 points in the plane in general position. The Geiser involution is defined as follows: a general point q in the plane determines a pencil of cubic curves which pass through q and the seven points p_1, \dots, p_7 ; this pencil has a ninth base-point, which is the image of q by the Geiser involution. This involution fixes a non hyperelliptic curve of genus 3 that is a sextic with ordinary double points at p_1, \dots, p_7 and whose smooth model is a plane quartic ([27] or [24]); the blow-up of the seven points conjugates the Geiser transformation to an automorphism of a del Pezzo surface of degree 2 ([1]).
- (b) Let p_1, \dots, p_8 be 8 points in the plane in general position. The Bertini involution is defined as follows: a general point q in the plane determines a linear system of sextic curves which pass through q and are singular at each of the eight points p_1, \dots, p_8 . This linear system has a tenth base-point, which is the image of q by the Bertini involution. This involution fixes a non hyperelliptic curve of genus 4 that is a nonic with ordinary triple points at p_1, \dots, p_8 and whose smooth model lies on a quadratic cone; the blow-up of the eight points conjugates the Bertini transformation to an automorphism of a del Pezzo surface of degree 1 (same references as above).

- (c) Let $C \subset \mathbb{P}^2$ be a curve of degree $g + 2$ with an ordinary g -fold point and which is smooth everywhere else. The de Jonquières involution associated to C is defined in the following way: the restriction of the transformation to a general line passing through the g -fold point of C is the unique involution that preserves this line and fixes the two other points of intersection of C with the line ([1, 15]).

Example 2.2.3 ([6, 14, 18])

Consider the smooth surface S defined by the equation $w^2 = z^3 + F_6(x, y)$ in the weighted projective space $\mathbb{P}(3, 1, 1, 2)$, where F_6 is a homogeneous polynomial of degree 6 with 6 simple roots: it is a particular type of del Pezzo surface of degree 1. The restriction of the map $(w : x : y : z) \mapsto (w : x : y : \omega z)$, where ω is a primitive cube root of 1, defines an automorphism of S of order 3 whose set of fixed points is the union of a point and an irreducible curve \tilde{C} of genus 2. The curve is linearly equivalent to $-2K_S$, hence any birational morphism $S \rightarrow \mathbb{P}^2$ sends this curve onto a sextic with 8 ordinary double points in general position.

Example 2.2.4 ([10])

Let $h \in \mathbb{C}[x]$ be a polynomial of degree $2g + 2$ without multiple roots. Consider the subgroup

$$T_h := \left\{ \begin{pmatrix} a_1 & ha_2 \\ a_2 & a_1 \end{pmatrix} : a_i \in \mathbb{C}(x), a_1^2 - ha_2^2 \neq 0 \right\}$$

of $\mathrm{GL}(2, \mathbb{C}(x))$ and denote by J_h its image in $\mathrm{PGL}(2, \mathbb{C}(x))$. To each $a \in J_h$, we associate a rational map $F_a : \mathbb{C}^2 \dashrightarrow \mathbb{C}^2$ defined by

$$(x, y) \mapsto \left(x, \frac{a_1 y + ha_2}{a_2 y + a_1} \right);$$

this is a de Jonquières transformation whose restriction to the hyperelliptic curve C of equation $(y^2 = h(x))$ is the identity. When $a_1 = 0$, we obtain an involution σ , conjugate to that of Example 2.2.2c).

Note that T_h is isomorphic to the multiplicative group $\mathbb{C}(C)^*$ of the field of rational functions $\mathbb{C}(C)$ on C , from which we deduce that J_h is isomorphic to $\mathbb{C}(C)^*/\mathbb{C}(x)^*$ and that its torsion is generated by σ .

Theorem 2.2.5 ([10, Theorem 1.5])

Let $C \subset \mathbb{P}^2$ be an irreducible curve of genus $g \geq 2$, and assume that $\mathrm{Ine}(C)$ is non-trivial.

Then, either $\mathrm{Ine}(C)$ is a cyclic group of order 2 or 3 generated by one of the transformations from Examples 2.2.2a, 2.2.2b, 2.2.3 or it is equal to the group J_h of Example 2.2.4, where $(y^2 = h(x))$ is the affine equation of C .

In particular, $\mathrm{Ine}(C)$ is Abelian and if it is infinite, then C is hyperelliptic and $\mathrm{Ine}(C)$ is a de Jonquières group, whose torsion is generated by a de Jonquières involution.

Note that Theorem 2.2.5 implies in particular that the elements of order 4 envisaged in Castelnuovo's theorem do not exist. It also implies the following result.

Corollary 2.2.6 Let S be a projective smooth rational surface and let $C \subset S$ be an irreducible curve with $g(C) > 1$. Then, the group of elements of $\mathrm{Aut}(S)$ that fix C has order 1, 2 or 3.

Proof Let us write $G = \mathrm{Ine}(S, C) \cap \mathrm{Aut}(S)$. According to Theorem 2.2.5, we may assume that G preserves a rational fibration $p : S \dashrightarrow \mathbb{P}^1$, and it suffices to show that no element

of G is of infinite order. Suppose to the contrary that some $\varphi \in G$ is of infinite order. After some blow-up we may assume that p is a morphism (since φ acts on the base-point of the fibration). Then, we replace φ by some power, and assume that φ preserves any component of any singular fibre of p . This implies that φ is conjugate to an automorphism of a Hirzebruch surface, which is not possible since it fixes a curve of positive genus. \square

2.3 The decomposition group of curves of genus ≥ 2

Applying the classification of the non-trivial inertia groups of curves of genus at least 2, we deduce

Theorem 2.3.1 *Let $C \subset \mathbb{P}^2$ be an irreducible curve of genus $g \geq 2$, and assume that $\text{Ine}(C)$ is non-trivial. Then, the canonical complex of (\mathbb{P}^2, C) is an exact sequence.*

Proof Theorem 2.2.5 restricts the possibilities for the pair (\mathbb{P}^2, C) .

The exactness of the canonical complex in the case where $\text{Ine}(C)$ is generated by the Geiser or Bertini involution is classical. For a proof (see [34, Thm. 1.8]), we consider the decomposition group as a subgroup of automorphisms of a del Pezzo surface S of degree 2 or 1 and denote by σ the Geiser or the Bertini involution; then each automorphism of the curve Γ fixed by σ extends to an automorphism of S because Γ is canonical in $S/\langle \sigma \rangle$.

In the de Jonquières case (Examples 2.2.2c and 2.2.4), denote by C the curve of degree $g + 2$, by p its g -fold point and by Λ the pencil of lines passing through p . Let j be a birational map of C . Since C is hyperelliptic j stabilizes the trace of Λ on C . Let $x \in \mathbb{P}^2$ be a general point. We can extend j to an element χ of $\text{Dec}(C)$: indeed, take the line $L \in \Lambda$ through p and x and set $L \cap C = \{p, q_x, r_x\}$; we define $\chi(x)$ by the relation

$$(p, x, q_x, r_x) = (p, \chi(x), j(q_x), j(r_x)),$$

where (a, b, c, d) denotes the cross ratio of a, b, c, d .

In the last case we consider the pair (S, \tilde{C}) as in Example 2.2.3 and observe that the restriction homomorphism $\text{Aut}(S, \tilde{C}) \rightarrow \text{Aut}(\tilde{C})$ is surjective: indeed, an automorphism of \tilde{C} extends to an automorphism of $\mathbb{P}(3, 1, 1)$ which lifts to an automorphism of S . \square

Remark 2.3.2 It can be observed that the exact sequence described above is split in the de Jonquières and Geiser cases, and in the case of Example 2.2.3. However it does not split in the Bertini case (there are square roots of the Bertini involution, see [6, Table 1]).

Theorem 2.3.1 completes the classification of pairs (\mathbb{P}^2, C) and canonical complexes such that $\text{Ine}(C) \neq 1$.

What happens when the group $\text{Ine}(C)$ is trivial? Firstly, we can state the following obvious result:

Lemma 2.3.3 *Let $C \subset \mathbb{P}^2$ be an irreducible curve of genus $g \geq 2$, and assume that $\text{Ine}(C) = 1$. Then $\text{Dec}(C)$ is isomorphic to a subgroup of $\text{Bir}(C)$, and is a finite group. In particular, when C is generic, $\text{Ine}(C) = \text{Dec}(C) = 1$.*

Proof It suffices to observe that $\text{Bir}(C)$ is isomorphic to the automorphism group of the normalization of C and to recall that this group is finite when $g(C) \geq 2$ and is trivial if C is generic. \square

The canonical complex is therefore trivially exact for a general curve. However, there exist examples where the map $\text{Dec}(C) \rightarrow \text{Bir}(C)$ is not surjective, see Sections 2.5, 3.3, 4.2 and

4.3. These examples rely on Theorem 2.3.4 below. For each point p that belongs to an irreducible curve $C \subset \mathbb{P}^2$, as a proper or infinitely near point, we denote by $m_p(C)$ the multiplicity of C (or of its strict transform) at p ; if p does not belong to C , we write $m_p(C) = 0$.

Theorem 2.3.4 *Let $C \subset \mathbb{P}^2$ be an irreducible curve of degree n and let $\varphi \in \text{Bir}(\mathbb{P}^2)$.*

Suppose that $3m_p(C) \leq n$ for each point p and that φ sends C onto a curve D of degree $\leq n$.

Then, each base-point q of φ belongs to C as a proper or infinitely near point, and $3m_q(C) = n$. Moreover, the degree of D is n .

Proof We may assume that φ is not an automorphism of \mathbb{P}^2 . Let Λ be the homolooidal net associated to φ (which is the strict pull-back by φ of the linear system of lines of \mathbb{P}^2) and let $\eta : X \rightarrow \mathbb{P}^2$ be a minimal birational morphism that solves the indeterminacies of φ (or equivalently the base-points of Λ). Denote by d the degree of φ (which is the degree of the curves of Λ), by q_1, \dots, q_k the base-points of φ (or Λ), which may be proper or infinitely near points of \mathbb{P}^2 , and by a_i the multiplicity of q_i as a base-point of Λ . We have $a_i \geq 1$ and $m_{q_i} = m_{q_i}(C) \geq 0$.

Consider now the strict transforms $\tilde{\Lambda}$ of Λ and \tilde{C} of C on X . Then, $\tilde{\Lambda}$ is base-point-free and $\tilde{\Lambda}^2 = 1$. Using the adjunction formula we find the classical equality $3(d-1) = \sum_{i=1}^k a_i$. Computing the free intersection of $\tilde{\Lambda}$ and \tilde{C} (which is equal to the degree of the image D of C , and is, by hypothesis, at most equal to n), we find $dn - \sum_{i=1}^k a_i \cdot m_{q_i} \leq n$. This yields, with the above equality:

$$\sum_{i=1}^k n \cdot a_i = 3n(d-1) \leq \sum_{i=1}^k 3m_{q_i} \cdot a_i. \quad (3)$$

Since $3m_{q_i} \leq n$ and $a_i \geq 1$ for $i = 1, \dots, k$, the inequality (3) is an equality. This implies that $\deg(D) = n$ and $3m_{q_i} = n$ for $i = 1, \dots, k$. \square

Corollary 2.3.5 *Let $C \subset \mathbb{P}^2$ be a smooth curve of degree n .*

1. *If $n = 3$, every base-point of each element of $\text{Dec}(C)$ belongs to C , as a proper or infinitely near point.*
2. *If $n > 3$, then every element of $\text{Dec}(C)$ is an automorphism of the plane, i.e. $\text{Dec}(C) = \text{Aut}(\mathbb{P}^2, C)$.*

Proof Apply Theorem 2.3.4, with $m_q(C) = 1$ for any point q that belongs to C as a proper or infinitely near point. \square

The first part of Corollary 2.3.5 can be found in [34, Theorem 1.3] and the second in [34, Cor. 3.6] and [30] (see also [37, p. 181] and [13, Book IV, Chap. VII, Sect. 3, Thm. 11]).

Another important corollary is the following one, which describes the inertia group of a family of classical curves (Halphen curves, Coble curves, ...) as a subgroup of automorphisms of a rational surface. We will use this to provide examples of plane curves whose canonical complex is not exact.

Corollary 2.3.6 *Let $p_1, \dots, p_k \in \mathbb{P}^2$ be k distinct proper points of \mathbb{P}^2 and let $C \subset \mathbb{P}^2$ be an irreducible curve of degree $3n$, with $n > 1$, which has multiplicity n at each \underline{p}_i . Denote by $\pi : X \rightarrow \mathbb{P}^2$ the blow-up of the k points and assume that the strict pull-back \tilde{C} of C by π is a smooth curve. Then,*

1. $\pi^{-1}\text{Dec}(\mathbb{P}^2, C)\pi = \text{Dec}(X, \tilde{C}) = \text{Aut}(X, \tilde{C})$.

2. Let $D \subset \mathbb{P}^2$ be an irreducible curve of degree $\leq 3n$ and let $\varphi : (\mathbb{P}^2, C) \dashrightarrow (\mathbb{P}^2, D)$ be a birational map. Denote by $\eta : Y \rightarrow \mathbb{P}^2$ an embedded minimal resolution of the singularities of D and by $\tilde{D} \subset Y$ the strict transform of D . Then, φ lifts to an isomorphism $\varphi' : (X, \tilde{C}) \rightarrow (Y, \tilde{D})$ such that $\eta\varphi' = \varphi\pi$. Furthermore, the degree of D is $3n$.

Proof Let us prove assertion (2). Theorem 2.3.4 implies that the base-point locus of φ is contained in $\{p_1, \dots, p_k\}$. Then, $\varphi\pi$ is a birational morphism $X \rightarrow \mathbb{P}^2$, that we denote by ν . Since the curve \tilde{C} is equivalent to $-nK_X$, the degree of the curve $\nu(\tilde{C}) = D$ is $3n$ and every (-1) -curve of X intersects \tilde{C} at n points. This implies that ν is an embedded minimal resolution of the singularities of D . The two birational morphisms ν and η differ only by an isomorphism $\varphi' : X \rightarrow Y$, which sends \tilde{C} onto \tilde{D} .

The assertion (1) is a particular case of (2). Indeed, for $\phi \in \text{Dec}(\mathbb{P}^2, C)$, the element $\phi' = \pi^{-1}\phi\pi$ belongs to $\text{Aut}(X, \tilde{C})$ and consequently the group $\pi^{-1}\text{Dec}(\mathbb{P}^2, C)\pi$ is contained in $\text{Aut}(X, \tilde{C})$; the other inclusion is obvious. □

2.4 Examples of different birational embeddings of curves of genus 2 in \mathbb{P}^2

Let C be any abstract smooth curve of genus 2. It is isomorphic to the curve $z^2 = F_6(x, y)$ in the weighted projective plane $\mathbb{P}(1, 1, 3)$, for some form F_6 of degree 6, having 6 simple roots. There exists a birational morphism $C \xrightarrow{\varphi_1} C_0$ where C_0 is a quartic of \mathbb{P}^2 with one ordinary double point, and furthermore there is only one choice of C_0 , up to birational equivalence of the pair (\mathbb{P}^2, C_0) (see [1]). The group $\text{Ine}(C_0)$ is infinite and described in Example 2.2.4 (Theorem 2.2.5); moreover the canonical complex of (\mathbb{P}^2, C_0) is an exact sequence (Theorem 2.3.1).

Let F_4 be any form of degree 4 in two variables (possibly equal to zero), and define S to be the surface with equation $w^2 = z^3 + zF_4(x, y) + F_6(x, y)$ in the weighted projective space $\mathbb{P}(3, 1, 1, 2)$. Since F_6 does not have any multiple roots, S is smooth and hence is a del Pezzo surface of degree 1 [28, Theorem 3.36]. There thus exists a birational morphism $\pi : S \rightarrow \mathbb{P}^2$ that consists of blowing-up 8 points in general position. Sending the curve C into S via the morphism $(x : y : z) \mapsto (z : x : y : 0)$ gives a curve $\tilde{C} \subset S$, equivalent to $-2K_S$, whose image by π is a sextic with eight ordinary double points.

If F_4 is the zero form, let $C_1 \subset \mathbb{P}^2$ denote the image of \tilde{C} by π . Then $\text{Ine}(C_1)$ is isomorphic to $\mathbb{Z}/3\mathbb{Z}$, and the canonical complex of (\mathbb{P}^2, C_1) is an exact sequence (Theorems 2.2.5 and 2.3.1).

If F_4 is not the zero form, then no non-trivial automorphism of S fixes the curve \tilde{C} , which means that $\text{Ine}(S, \tilde{C})$ is trivial (Corollary 2.3.6). The Bertini involution on S (that sends w onto $-w$) leaves \tilde{C} invariant, acts on it as the involution associated to the $g^1_{1/2}$, and generates $\text{Dec}(S, \tilde{C})$ if F_4 is general enough. Moreover $\text{Aut}(\tilde{C})$ is reduced to this involution if and only if no non-trivial automorphism of \mathbb{P}^1 leaves F_6 invariant. It then follows from Corollary 2.3.6 that the canonical complex of (\mathbb{P}^2, C_2) is an exact sequence under these circumstances, where C_2 denotes the image of \tilde{C} by π .

These examples provide three birational embeddings $C \rightarrow C_i \subset \mathbb{P}^2$ that lead to three different canonical complexes for the same abstract curve and also to three birationally different pairs (\mathbb{P}^2, C_i) . Theorem 2.3.4 allows us to improve this result, giving infinitely many such pairs of the last kind. Indeed, let F_4 and F'_4 be two different forms of degree 4, let $\tilde{C} \subset S$ and $\tilde{C}' \subset S'$ be the two embeddings of C into two corresponding del Pezzo surfaces of degree 1, and let $C_2 \subset \mathbb{P}^2$ and $C'_2 \subset \mathbb{P}^2$ be the corresponding sextic curves. If there exists

a birational transformation φ that sends (\mathbb{P}^2, C_2) onto (\mathbb{P}^2, C'_2) , then Corollary 2.3.6 implies that φ lifts to an isomorphism $S \rightarrow S'$. By changing our choice of F_4 , we obtain infinitely many isomorphism classes of del Pezzo surfaces of degree 1, that lead to infinitely many birationally different pairs (\mathbb{P}^2, C_2) such that C_2 is birational to C .

2.5 Examples of different birational embeddings of curves of genus 3 in \mathbb{P}^2

We give another example. Let $C_1 \subset \mathbb{P}^2$ be any smooth quartic curve. The double covering of \mathbb{P}^2 ramified over C_1 is a del Pezzo surface S of degree 2 (see [2]), which is the blow-up $\pi : S \rightarrow \mathbb{P}^2$ of 7 points of \mathbb{P}^2 in general position. Denote by \tilde{C} the image of C_1 on S and by C_2 the curve $\pi(\tilde{C})$. Then, C_2 is a sextic with 7 ordinary double points and $\text{Ine}(C_2) \cong \mathbb{Z}/2\mathbb{Z}$ is generated by the Geiser involution that corresponds to the involution of S associated to the double covering (Theorem 2.2.5). On the other hand, Corollary 2.3.5 implies that $\text{Dec}(C_1) = \text{Aut}(\mathbb{P}^2, C_1)$ and consequently that $\text{Ine}(C_1)$ is trivial. The two curves C_1 and C_2 are birational curves of the plane, but the pairs (\mathbb{P}^2, C_1) and (\mathbb{P}^2, C_2) have different canonical complexes, and in particular are not birationally equivalent.

2.6 Examples of different birational embeddings of curves of genus 4 in \mathbb{P}^2

Let p_1, \dots, p_8 be eight points of the plane and let $S \rightarrow \mathbb{P}^2$ be their blow-up. Assume that S is a del Pezzo surface. Corollary 2.3.6 implies the following observations. Among the linear system Λ of nonics passing through p_1, \dots, p_8 with multiplicity 3, exactly one has a non-trivial inertia group, generated by the Bertini involution of S . The other curves of Λ have a decomposition group that contains the Bertini involution, and for a general curve of Λ this involution generates the decomposition group. Furthermore, the elements of Λ yield infinitely many pairs which are birationally different.

3 Curves of genus one

In Sect. 2, we gave a precise description of all elements of finite order of $\text{Bir}(\mathbb{P}^2)$ that fix a curve of genus ≥ 2 . A precise description also exists for curves of genus 1:

Theorem 3.0.1 ([8, Theorem 2])

Let $C \subset \mathbb{P}^2$ be an irreducible curve with $g(C) = 1$. Let $h \in \text{Ine}(C)$ be an element of finite order $n > 1$. Then, there exists a birational map $\varphi : \mathbb{P}^2 \dashrightarrow S$ that conjugates h to an automorphism α of a del Pezzo surface S , with (α, S, n) given in the following table:

| n | Description of α | Equation of the surface S | In the variety |
|-----|---------------------------|--|----------------------------|
| 2 | $x_0 \mapsto -x_0$ | $\sum_{i=0}^4 x_i^2 = \sum_{i=0}^4 \lambda_i x_i^2 = 0$ | \mathbb{P}^4 |
| 3 | $x_0 \mapsto \zeta_3 x_0$ | $x_0^3 + L_3(x_1, x_2, x_3)$ | \mathbb{P}^3 |
| 4 | $x_0 \mapsto \zeta_4 x_0$ | $x_3^2 = x_0^4 + L_4(x_1, x_2)$ | $\mathbb{P}(1, 1, 1, 2)$ |
| 5 | $x_0 \mapsto \zeta_5 x_0$ | $x_3^2 = x_2^3 + \lambda_1 x_1^4 x_2 + x_1(\lambda_2 x_1^5 + x_0^5)$ | $\mathbb{P}(1, 1, 2, 3)$ |
| 6 | $x_0 \mapsto \zeta_6 x_0$ | $x_3^2 = x_2^3 + \lambda_1 x_1^4 x_2 + \lambda_2 x_1^6 + x_0^6$ | $\mathbb{P}(1, 1, 2, 3)$, |

where $\zeta_n \in \mathbb{C}$ is a primitive n -th root of unity, L_i is a form of degree i and the λ_i are parameters such that S is smooth.

Furthermore, any birational morphism $S \rightarrow \mathbb{P}^2$ sends the fixed curve onto a smooth plane cubic curve.

Theorem 3.0.1 implies in particular the following result:

Corollary 3.0.2 *Let $C \subset \mathbb{P}^2$ be an irreducible curve with $g(C) = 1$. The following conditions are equivalent:*

- (1) *the pair (\mathbb{P}^2, C) is birationally equivalent to a pair (\mathbb{P}^2, D) , where D is a smooth cubic curve;*
- (2) *the group $\text{Ine}(C)$ contains non-trivial elements of finite order;*
- (3) *the group $\text{Ine}(C)$ contains elements of order 2, 3, 4, 5 and 6.*

Proof In order to prove (1) \Rightarrow (3) we observe that in each of the five types of pairs (α, S) of Theorem 3.0.1, one gets an arbitrary elliptic curve as fixed curve. The implication (3) \Rightarrow (2) is obvious and (2) \Rightarrow (1) follows from Theorem 3.0.1. □

The curves of genus 1 having the biggest canonical complex seem in fact to be the cubic curves. We will make this more precise in (3.3). We examine in (3.1) and (3.2) the case of smooth cubic curves and then in (3.3) the other irreducible curves of genus 1.

3.1 The inertia group of smooth cubic curves

Let $C \subset \mathbb{P}^2$ be a smooth cubic curve. For any point $p \in C$, there exist infinitely many elements of $\text{Ine}(C)$ that leave invariant any general line passing through p ; such elements form a group described in Example 2.2.4. There are furthermore many elements of degree 3 in this group ([8, Lemma 4.1]); one of these, that we call σ_p , is the classical de Jonquières involution of Example 2.2.2c (generalised in [23] under the name of R_p to any dimension). The element σ_p is the unique involution that leaves invariant any general line passing through p and fixes the curve C .

By changing the choice of p , all these involutions generate a very large group:

Theorem 3.1.1 ([8, Theorem 1.6])

Let $C \subset \mathbb{P}^2$ be a smooth cubic curve. The subgroup of $\text{Ine}(C)$ generated by all the cubic involutions centred at the points of C is the free product

$$\star_{p \in C} \langle \sigma_p \rangle .$$

Furthermore, since the inertia group of a smooth cubic curve contains elements of order 3, 4, 5 and 6 (Corollary 3.0.2), the free product described in Theorem 3.1.1 is not the whole inertia group. However, there exists an analogue of the Noether-Castelnuovo theorem for this group:

Theorem 3.1.2 ([8, Theorem 1.1]) *The inertia group of a smooth plane cubic curve is generated by its elements of degree 3, which are—except the identity—its elements of lower degree.*

3.2 The decomposition group of smooth cubic curves

Let C be a smooth plane cubic curve. Take three distinct points p, q, r that belong to C as proper or infinitely near points. The linear system of conics passing through these points

defines a birational transformation φ of \mathbb{P}^2 which transforms C onto another smooth cubic curve C' . Composing φ with a linear automorphism mapping C' onto C we obtain a degree 2 element in $\text{Dec}(C)$. Clearly these transformations are the only degree 2 elements in $\text{Dec}(C)$. Moreover, all such transformations may be expressed as a composition of those whose base-point set consists of three proper points of the plane. As for the inertia group, there exists a result analogous to the Noether-Castelnuovo theorem for the decomposition group:

Theorem 3.2.1 ([34, Theorem 1.4])

The decomposition group of a smooth plane cubic curve is generated by its elements of degree 2.

Concerning the action of the decomposition group on the elliptic curve, the following result shows that this is like the whole automorphism group of the curve:

Theorem 3.2.2 ([23, Theorem 6])

Let $C \subset \mathbb{P}^2$ be a smooth cubic curve. The canonical complex of (\mathbb{P}^2, C) is an exact sequence.

Remark 3.2.3 It seems that this sequence is not split.

3.3 Curves of genus 1 that are not equivalent to smooth cubic curves

We recall some classical notions on Halphen curves and surfaces (see [25, 12, 22, 19]).

Definition 3.3.1 A Halphen curve of index n is an irreducible plane curve of degree $3n$, with 9 points of multiplicity n and of genus 1.

A projective rational smooth surface S is a Halphen surface of degree n if the linear system $| -nK_S |$ is a pencil whose general fibre is an irreducible curve of genus 1.

The following classical relation can be verified by hand:

Lemma 3.3.2 *If S is a Halphen surface of degree n , any birational morphism $S \rightarrow \mathbb{P}^2$ sends the general fibres of $| -nK_S |$ onto Halphen curves of index n .*

For $n \geq 2$, the blow-up of the 9 singular points of a Halphen curve of index n is a Halphen surface.

The blow-up of 9 general points is not a Halphen surface. However, for any general set of 8 points of the plane, and for any integer $n \geq 2$, there exists a curve of the plane such that the blow-up of the 8 points and a ninth point on the curve gives a Halphen surface of index n [25].

We now give a simple proof of the following (probably classical) result:

Proposition 3.3.3 *Let $C_1, C_2 \subset \mathbb{P}^2$ be two Halphen curves of index respectively n_1 and n_2 . For $i = 1, 2$, consider the minimal embedded resolution $X_i \rightarrow \mathbb{P}^2$ of C_i (which is the identity if $n_i = 1$). The following assertions are equivalent:*

- (1) *the pairs (\mathbb{P}^2, C_1) and (\mathbb{P}^2, C_2) are birationally equivalent;*
- (2) *there exists an isomorphism $\varphi : X_1 \rightarrow X_2$ that sends the strict transform of C_1 onto the strict transform of C_2 .*

Furthermore, both assertions imply that $n_1 = n_2$.

Proof The second assertion directly implies the first and the equality $n_1 = n_2$. Corollary 2.3.6 shows that the first assertion implies the second. \square

This proposition shows in particular the existence of infinitely many distinct types of pairs (\mathbb{P}^2, C) where C has genus 1; it also raises the following question, which we think is still open:

Question 3.3.4 Let $C_1 \subset \mathbb{P}^2$ be an irreducible curve of genus 1. Does there exist a Halphen curve $C \subset \mathbb{P}^2$ such that the pair (\mathbb{P}^2, C_1) is birationally equivalent to (\mathbb{P}^2, C) ?

We now describe the decomposition and inertia groups of Halphen curves of index ≥ 2 (those of index 1 are the smooth cubic curves, described previously), and show in particular the important difference between index 1 and index ≥ 2 .

Theorem 3.3.5 *Let $C \subset \mathbb{P}^2$ be a Halphen curve of index $n \geq 2$ and assume that the pencil induced by C has no reducible fibre. Then, $\text{Dec}(C)$ contains a normal subgroup of finite index, isomorphic to \mathbb{Z}^8 . In particular, the canonical complex of (\mathbb{P}^2, C) is not exact.*

Assume that C is a general Halphen curve; then $\text{Dec}(C)$ is isomorphic either to $\mathbb{Z}^8 \rtimes \mathbb{Z}/2\mathbb{Z}$ or to \mathbb{Z}^8 . The first case occurs for $n = 2$ and the second one if $n = 3$ or $n \geq 5$.

Proof Let $\pi : S \rightarrow \mathbb{P}^2$ be the blow-up of the nine singular points of C and let $\tilde{C} \subset S$ be the strict transform of C . Corollary 2.3.6 implies that $\text{Dec}(C) = \text{Dec}(\mathbb{P}^2, C)$ is conjugate to $\text{Dec}(S, \tilde{C}) = \text{Aut}(S, \tilde{C})$.

Denote by $D \subset \mathbb{P}^2$ a cubic passing through the singular points of C and by $\tilde{D} \subset S$ the strict pull-back of D . Then \tilde{C} and $n\tilde{D}$ belong to the pencil $| -nK_S |$ and this shows that D is unique.

Note that $\text{Aut}(S)$ acts on the elliptic fibration $\eta : S \rightarrow \mathbb{P}^1$ induced by $| -nK_S |$. Let $G \subset \text{Aut}(S)$ be the subgroup of automorphisms that act trivially on the basis and let G' be the image of $\text{Aut}(S)$ in $\text{Aut}(\mathbb{P}^1)$, such that the following is an exact sequence:

$$1 \rightarrow G \rightarrow \text{Aut}(S) \rightarrow G' \rightarrow 1.$$

We show that G' is finite. Indeed, $\eta(\tilde{D})$ is a fixed point, so we may consider G' as a subgroup of $\text{Aut}(\mathbb{C})$; then G' has at least one finite orbit in \mathbb{C} because there are singular fibres in $| -nK_S |$ (the Euler characteristic of S , which is equal to 12, is the sum of the Euler characteristics of the (singular) fibres of η), which is not possible if G' is infinite.

Now, let $H \subset G$ be the subgroup of elements that act as translations on the general fibre; according to the structure of the automorphism group of an elliptic curve and since G' is finite, H is normal in $\text{Aut}(S)$, of finite index.

A translation on an elliptic curve corresponds to a linear equivalence of a divisor of degree 0. Since no fibre is reducible, there exists an exact sequence (see [22])

$$0 \rightarrow \mathbb{Z}K_S \rightarrow K_S^\perp \rightarrow H \rightarrow 0,$$

where K_S^\perp is the subgroup of $\text{Pic}(S)$ of elements whose intersection with K_S is equal to 0. Since $\text{Pic}(S) \cong \mathbb{Z}^{10}$ and K_S is indivisible, $K_S^\perp \cong \mathbb{Z}^9$ and $H \cong \mathbb{Z}^8$. As $H \subset G \subset \text{Dec}(S, \tilde{C}) \subset \text{Aut}(S)$, the first assertion is proved.

Assume now that C is a general Halphen curve, which implies that $\text{Dec}(S, \tilde{C})$ is equal to G and that the automorphism group of \tilde{C} is equal to $\tilde{C} \rtimes \mathbb{Z}/2\mathbb{Z}$. In particular, the index of H in G is either 2 or 1, depending on whether or not there exists an element of G that acts as an involution with four fixed points on the general fibre. Assume that such an element $\sigma \in G$ exists. Then, it fixes a curve in S which intersects the general fibre in four points. The fibre is equal to $-nK_S$, hence n must divide 4, which implies that $n = 2$ or $n = 4$.

It remains to show that for $n = 2$, such an involution exists. Consider the elliptic fibration $\epsilon : S \dashrightarrow \mathbb{P}^1$ defined by the pencil of plane cubics passing through eight of the nine base-points of the Halphen pencil: the intersection of a general fibre S_η of η with a general fibre of

ϵ is equal to 2, which means that the degree of $\eta \times \epsilon : S \dashrightarrow \mathbb{P}^1 \times \mathbb{P}^1$ is 2; the corresponding involution of S leaves each Halphen curve S_η invariant and has thereon 4 fixed points since the restriction to S_η of ϵ is simply the canonical g_2^1 . \square

Remark 3.3.6 In the case where $n = 2$ and the points are in general position, it may also be observed that the Bertini involution associated to the blow-up of 8 of the 9 points lifts to an automorphism of the surface that acts on each member of the elliptic fibration as an automorphism with four fixed points. Furthermore, the 9 involutions obtained via this map generate the automorphism group of the Halphen surface [12].

4 Rational curves

The case of rational curves is less well described. We can however quote some simple results.

4.1 The line

There exist many elements in the inertia group of the line; for example, any birational map of the form $(x, y) \dashrightarrow \left(\frac{x}{\alpha(y)x + \beta(y)}, y \right)$, where $\alpha, \beta \in \mathbb{C}(y)$, $\beta \neq 0$, fixes the line $x = 0$. It seems that the inertia group of a line is a big and complicated group. Let us make some simple observations:

Proposition 4.1.1 *Let $L \subset \mathbb{P}^2$ be a line, then the canonical complex of (\mathbb{P}^2, L) is a split exact sequence.*

Furthermore, the group $\text{Ine}(\mathbb{P}^2, L)$ is neither finite, nor Abelian and does not leave invariant any pencil of rational curves.

Proof It is obvious that the complex is exact, and split: the group of automorphisms of L extends to a subgroup of $\text{Aut}(\mathbb{P}^2)$, and this yields a section $\text{Aut}(L) \rightarrow \text{Aut}(\mathbb{P}^2, L)$. The other assertions follow from [10, Proposition 4.1]. \square

Does there exist an analogue of the Noether–Castelnuovo theorem, as in the case of smooth cubics?

Question 4.1.2 *Let $L \subset \mathbb{P}^2$ be a line. Is the group $\text{Dec}(\mathbb{P}^2, L)$ (respectively $\text{Ine}(\mathbb{P}^2, L)$) generated by its elements of degree 1 and 2?*

4.2 Coble curves

A Coble curve is an irreducible sextic with 10 double points. There does not exist a sextic singular at ten general points; however Coble curves exist, and are singular members of a Halphen pencil of index 2; furthermore in each such pencil there are in general 12 Coble curves [25]. Corollary 2.3.6 implies that the pair (\mathbb{P}^2, C) where C is a Coble curve is not equivalent to that of a line. Furthermore, we have:

Proposition 4.2.1 *Let $C \subset \mathbb{P}^2$ be a Coble curve, let $\pi : S \rightarrow \mathbb{P}^2$ be the blow-up of its 10 singular points and let $\tilde{C} \subset S$ be the strict pull-back of C by π .*

Then, $\text{Aut}(S) = \text{Dec}(S, \tilde{C}) = \pi^{-1}\text{Dec}(\mathbb{P}^2, C)\pi$.

Proof The curve \tilde{C} is equivalent to $-2K_S$ and since it has negative self-intersection it is the only such curve, consequently $\text{Aut}(S) = \text{Aut}(S, \tilde{C})$. The result then follows directly from Corollary 2.3.6. \square

The description of the automorphisms of a so-called Coble surface obtained by blowing-up the ten singular points of a Coble curve is a classical result of Coble [12], see also [19, Theorem 8, page 107]. It implies in particular the following result:

Proposition 4.2.2 *For any Coble curve C , the group $\text{Dec}(\mathbb{P}^2, C)$ is an infinite countable group. The canonical complex associated to (\mathbb{P}^2, C) is not exact.*

4.3 Other curves of Halphen type

Let S be a Halphen surface of index n obtained by the blow-up $\pi : S \rightarrow \mathbb{P}^2$ of the points p_1, \dots, p_9 . There exist singular fibres of the elliptic fibration $| -nK_S |$, which are thus rational curves with a double point, whose image on \mathbb{P}^2 are curves of degree $3n$ with multiplicity n at the points p_1, \dots, p_9 and multiplicity 2 at some other point p_{10} . The case $n = 1$ gives nodal cubics, which are equivalent to lines; the case $n = 2$ gives Coble curves, and the case $n = 3$ gives other curves. Once again, it seems that in general 12 such curves exist in a general Halphen pencil [25].

Proposition 4.3.1 *Let $C \subset \mathbb{P}^2$ be an irreducible curve of degree $3n$ with multiplicity n at p_1, \dots, p_9 and multiplicity 2 at p_{10} , and assume that $n \geq 3$. Let $\pi : S \rightarrow \mathbb{P}^2$ (respectively $\pi' : S' \rightarrow \mathbb{P}^2$) be the blow-up of p_1, \dots, p_{10} (respectively of p_1, \dots, p_9), and let $\tilde{C} \subset S$ and $\tilde{C}' \subset S'$ be the strict pull-backs of C by π and π' .*

Then, $\text{Aut}(S) = \text{Dec}(S, \tilde{C}) = \pi^{-1}\text{Dec}(\mathbb{P}^2, C)\pi$ and $\text{Aut}(S', \tilde{C}') = \text{Dec}(S, \tilde{C}') = \pi'^{-1}\text{Dec}(\mathbb{P}^2, C)\pi'$.

Furthermore, $\text{Dec}(\mathbb{P}^2, C)$ contains a subgroup of finite index isomorphic to \mathbb{Z}^8 . In particular, the canonical complex associated to (\mathbb{P}^2, C) is not an exact sequence.

Proof As for Corollary 2.3.6, Theorem 2.3.4 implies the equalities $\text{Aut}(S, \tilde{C}) = \text{Dec}(S, \tilde{C}) = \pi^{-1}\text{Dec}(\mathbb{P}^2, C)\pi$. The curve $E_{10} = \pi^{-1}(p_{10})$ is a smooth irreducible rational curve of self-intersection -1 (a (-1) -curve) and its intersection with \tilde{C} is 2; furthermore, it is the unique such curve [29, Theorem 3.3]. Consequently, the whole group $\text{Aut}(S)$ leaves E_{10} invariant; denoting by $\mu : S \rightarrow S'$ the blow-down of this curve (such that $\pi = \pi'\mu$), the group $G = \mu\text{Aut}(S)\mu^{-1}$ is the subgroup of $\text{Aut}(S')$ that fixes the point $(\pi')^{-1}(p_{10}) = \mu(E_{10})$, which is the unique singular point of \tilde{C}' . Since S' is a Halphen surface of index n and \tilde{C}' is a singular member of the fibration, $G = \text{Aut}(S', \tilde{C}')$. This implies the remaining equalities.

The last part follows from Theorem 3.3.5. □

4.4 Other rational curves

Do there exist other examples of pairs (\mathbb{P}^2, C) where C is rational? A famous problem of Coolidge and Nagata asks whether the pair of a rational cuspidal curve is birationally equivalent to the pair of a line (see [13] and [32]).

Definition 4.4.1 Let $C \subset S$ be an irreducible smooth curve on a surface. We denote by $\kappa(S, C)$ the Kodaira dimension of the pair (S, C) : this is the dimension of the image of $X \dashrightarrow \mathbb{P}(H^0(m(C + K_S))^\vee)$ for m large enough. If $|m(C + K_S)| = \emptyset$ for all $m > 0$, the Kodaira dimension is by convention equal to $-\infty$.

For a singular curve $C \subset S$, we write $\kappa(S, C) = \kappa(X, \tilde{C})$, where $X \rightarrow S$ is an embedded resolution of the singularities of C and $\tilde{C} \subset X$ is the strict transform.

Lemma 4.4.2 ([29])

If (S, C) is birationally equivalent to (S', C') then $\kappa(S, C) = \kappa(S', C')$.

Let us quote the following fundamental result, due to Coolidge.

Theorem 4.4.3 ([13, 29])

Let $C \subset \mathbb{P}^2$ be a rational irreducible curve and $L \subset \mathbb{P}^2$ be a line. Then (\mathbb{P}^2, C) is birationally equivalent to (\mathbb{P}^2, L) if and only if $\kappa(\mathbb{P}^2, C) = -\infty$.

We also have a description for Kodaira dimensions 0 and 1:

Theorem 4.4.4 ([29])

Let $C \subset \mathbb{P}^2$ be a rational irreducible curve.

- (1) $\kappa(\mathbb{P}^2, C) = 0$ if and only if (\mathbb{P}^2, C) is birationally equivalent to (\mathbb{P}^2, D) where D is a Coble curve.
- (2) $\kappa(\mathbb{P}^2, C) = 1$ if and only if (\mathbb{P}^2, C) is birationally equivalent to (\mathbb{P}^2, D) , where D is a curve of degree $3n$, with 9 points of multiplicity $n > 2$ and a tenth point of multiplicity 2.

Consequently, finding other rational curves not equivalent to our examples is equivalent to finding rational curves $C \subset \mathbb{P}^2$ with $\kappa(\mathbb{P}^2, C) = 2$.

5 Link between the inertia and decomposition groups and the classification of finite subgroups of the Cremona group

In our description of the decomposition group, and more precisely of the inertia group of a curve of genus ≥ 1 , we provide some subgroups of finite order. Conversely, in the study of the finite subgroups of $\text{Bir}(\mathbb{P}^2)$, the set of birational classes of curves of positive genus fixed (pointwise) is an important conjugacy invariant. For example, this invariant was used to find infinitely many conjugacy classes of elements of order $2n$ of $\text{Bir}(\mathbb{P}^2)$, for any integer n (see [7, Theorem 1.2]).

5.1 Cyclic groups of prime order

The conjugacy class of a finite cyclic subgroup of prime order of $\text{Bir}(\mathbb{P}^2)$ is uniquely determined by the birational equivalence of the curve of positive genus that it fixes (it can fix at most one such curve):

Theorem 5.1.1 ([1, 14]) *Let g, g' be two elements of $\text{Bir}(\mathbb{P}^2)$ of the same prime order, that fix respectively Γ, Γ' , two irreducible curves of positive genus. Then, g and g' are conjugate in $\text{Bir}(\mathbb{P}^2)$ if and only if the curves Γ and Γ' are birational.*

Theorem 5.1.2 ([3]) *An element of $\text{Bir}(\mathbb{P}^2)$ of prime order is not conjugate to a linear automorphism if and only if it belongs to the inertia group of some curve of positive genus.*

5.2 Other groups

Theorem 5.1.2 extends to finite cyclic groups of any order, and almost to finite Abelian groups:

Theorem 5.2.1 ([9], announced in [6])

Let G be a finite cyclic subgroup of order n of $\text{Bir}(\mathbb{P}^2)$. The following conditions are equivalent:

- If $g \in G, g \neq 1$, then g does not fix a curve of positive genus.
- G is birationally conjugate to a subgroup of $\text{Aut}(\mathbb{P}^2)$.
- G is birationally conjugate to a subgroup of $\text{Aut}(\mathbb{P}^1 \times \mathbb{P}^1)$.
- G is birationally conjugate to the group of automorphisms of \mathbb{P}^2 generated by $(x : y : z) \mapsto (x : y : e^{2i\pi/n}z)$.

Theorem 5.2.2 ([9], announced in [6])

Let G be a finite Abelian subgroup of $\text{Bir}(\mathbb{P}^2)$. The following conditions are equivalent:

- If $g \in G, g \neq 1$, then g does not fix a curve of positive genus.
- G is birationally conjugate to a subgroup of $\text{Aut}(\mathbb{P}^2)$, or to a subgroup of $\text{Aut}(\mathbb{P}^1 \times \mathbb{P}^1)$ or to the group $C_{S_{24}}$ isomorphic to $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$, generated by the two elements

$$\begin{aligned} (x : y : z) &\mapsto (yz : xy : -xz), \\ (x : y : z) &\mapsto (yz(y - z) : xz(y + z) : xy(y + z)). \end{aligned}$$

Moreover, this last group is conjugate neither to a subgroup of $\text{Aut}(\mathbb{P}^2)$, nor to a subgroup of $\text{Aut}(\mathbb{P}^1 \times \mathbb{P}^1)$.

However, there are plenty of examples of finite non-Abelian subgroups of $\text{Bir}(\mathbb{P}^2)$ which are not birationally conjugate to a subgroup of $\text{Aut}(\mathbb{P}^2)$ or of $\text{Aut}(\mathbb{P}^1 \times \mathbb{P}^1)$ but such that no curve of positive genus is fixed by any non-trivial element of the group [9, Section 12].

6 The links with the dynamical properties of a Cremona transformation

We can also consider a Cremona transformation as defining a dynamical system. In comparison with the usual case of dynamics defined by automorphisms, the situation here is more complicated due to the presence of indeterminacies and critical points: in the neighbourhood of such points the map does not act in a "natural way".

In [21] and [35] the authors introduce the so-called first dynamical degree of a birational map; this number is invariant by birational conjugation. Let us explain what that degree means in the case of a Cremona transformation φ . Consider the sequence $(\text{deg}(\varphi^n))^{1/n}$ for $n \geq 1$. Since $\text{deg}(\varphi^{n+m}) \leq \text{deg}(\varphi^n) \cdot \text{deg}(\varphi^m)$ it has a limit. The first dynamical degree is then

$$\lambda(\varphi) := \lim_{n \rightarrow \infty} (\text{deg}(\varphi^n))^{1/n}.$$

As shown in [21] the topological entropy $h_{top}(\varphi)$ of φ is at most $\log \lambda(\varphi)$. The equality is conjectured, and proved for a general φ ([4] and [20]).

On the other hand, Diller and Favre propose a more refined approach and consider the sequence of successive degrees $\text{deg}(\varphi^n)$ itself. They classify the plane Cremona transformations (in fact they consider a more general setup) with $\lambda = 1$ in terms of the growth rate of that sequence (see [16, Thm. 0.2]); they show that this growth is at most quadratic in n .

It is natural to ask to what extent the dynamics of a Cremona transformation can be affected by the existence of a genus g stable curve, $g = 0, 1, 2, \dots$ One answer is as follows:

Theorem 6.0.3 ([33, Theorem 1.1])

Let $C \subset \mathbb{P}^2$ be an irreducible curve of genus $g(C) \geq 2$ and let $\varphi \in \text{Dec}(C)$; then $\lambda(\varphi) = 1$ and the sequence $\{\text{deg}(\varphi^n)\}_{n=1}^\infty$ grows at most linearly.

Proof Since $g(C) > 1$, the subgroup $\text{Ine}(C)$ is of finite index in $\text{Dec}(C)$ so we may assume that $\varphi \in \text{Ine}(C)$. If φ is of finite order, we are done. If not, Theorem 2.2.5 yields an explicit description of φ as in Example 2.2.4. Computing the degrees of the powers of φ , we have $\deg(\varphi^n) \leq n(\deg(\varphi) + c)$ for some constant c . This completes the proof. \square

For another proof and some generalizations of this result see [17].

On the other hand, when $g(C) \leq 1$ the number $\lambda(\varphi)$ can be strictly larger than 1. For an example in the rational case we refer the reader to [33, Example 2]. In the case of a smooth cubic curve, the composition of two generic quadratic elements of the decomposition group seems suitable.

Finally, until recently, all known examples of automorphisms of rational surfaces with a first dynamical degree strictly larger than 1 (or equivalently with an action of infinite order on the Picard group) were those which leave invariant a rational or an elliptic curve. A question/conjecture of Gizatullin/Harbourne/McMullen (see [26] and [31]) asked whether this was always the case. A counterexample is announced in [5] which provides the existence of automorphisms of rational surfaces that do not leave invariant any curve.

References

1. Bayle, L., Beauville, A.: Birational involutions of \mathbb{P}^2 . *Asian. J. Math.* **4**(1), 11–17 (2000)
2. Beauville, A.: Complex algebraic surfaces, London Math. Soc. Student Texts, 34 (1996)
3. Beauville, A., Blanc, J.: On Cremona transformations of prime order. *C.R. Acad. Sci., Ser. 1, Math.* **339**(4), 257–259 (2004)
4. Bedford, E., Diller, J.: Energy and invariant measures for birational surface maps. *Duke Math. J.* **128**(2), 331–368 (2005)
5. Bedford, E., Kim, K.: Dynamics of rational surface automorphisms: linear fractional recurrences, arXiv:math/0611297
6. Blanc, J.: Finite Abelian subgroups of the Cremona group of the plane. *C.R. Acad. Sci. Paris Ser. I, Math.* **344**, 21–26 (2007)
7. Blanc, J.: The number of conjugacy classes of elements of the Cremona group of some given finite order. *Bull. Soc. Math. France* **135**(3), 419–434 (2007)
8. Blanc, J.: On the inertia group of elliptic curves in the Cremona group of the plane. *Michigan Math. J.* **56**(2), 315–330 (2008)
9. Blanc, J.: Linearisation of finite Abelian subgroups of the Cremona group of the plane. *Groups Geom. Dyn.* arXiv:math/0704.0537 (to appear)
10. Blanc, J., Pan, I., Vust, T.: Sur un théorème de Castelnuovo. *Bull. Braz. Math. Soc. (N.S.)* **39**(1), 61–80 (2008)
11. Castelnuovo, G.: Sulle trasformazioni cremoniane del piano che ammettono una curva fissa, *Rend. Accad. Lincei* (1892); *Memorie scelte*, Bologna, Zanichelli (1937)
12. Coble, A.B.: Algebraic geometry and theta functions, Amer. Math. Soc. Colloquium Publications, Vol. X (1961)
13. Coolidge, J.L.: A treatise on algebraic curves, Dover Publications, Inc. (1959)
14. de Fernex, T.: On planar Cremona maps of prime order. *Nagoya Math. J.* **174**, 1–28 (2004)
15. Jonquières, E.de : De la transformation géométrique des figures planes, et d'un mode de génération de certaines courbes à double courbure de tous les ordres. *Nouv. Ann.* (2)**3**, 97–111 (1864)
16. Diller, J., Favre, C.: Dynamics of bimeromorphic maps of surfaces. *Amer. J. Math.* **123**(6), 1135–1169 (2001)
17. Diller, J., Jackson, D., Sommese, A.: Invariant curves for birational surface maps. *Trans. Amer. Math. Soc.* **359**(6), 2793–2991 (2007)
18. Dolgachev, I.V., Iskovskikh, V.A.: Finite subgroups of the plane Cremona group. In: *Arithmetic and Geometry-Manin Festschrift*. math.AG/0610595 (2006)
19. Dolgachev, I.V., Ortland, D.: Point sets in projective spaces and theta functions, *Astérisque*, No. 165 (1988), 210 pp. (1989)
20. Dujardin, R.: Laminar currents and birational dynamics. *Duke Math. J.* **131**(2), 219–247 (2006)
21. Friedland, S.: Entropy of algebraic maps. Proceedings of the Conference in Honor of Jean-Pierre Kahane (Orsay, 1993). *J. Fourier Anal. Appl. Special Issue*, 215–228 (1995)

22. Gizatullin, M.H.: Rational G -surfaces. *Izv. Akad. Nauk SSSR Ser. Mat.* **44**(1), 110–144, 239 (1980)
23. Gizatullin, M.H.: The decomposition, inertia and ramification groups in birational geometry. *Algebraic Geometry and Its Applications, Aspects of Mathematics, E*, **25**, 39–45 (1994)
24. Godeaux, L.: *Géométrie algébrique II, géométrie sur une courbe algébrique, géométrie algébrique du plan*, Masson & Cie, éditeurs, Paris (1950)
25. Halphen, M.: Sur les courbes planes du sixième degré à neuf points doubles. *Bull. Soc. Math. France* **10**, 162–172 (1882)
26. Harbourne, B.: Rational surfaces with infinite automorphism group and no antipluricanonical curve. *Proc. AMS.* **99**(3), 409–414 (1987)
27. Hudson, H.: *Cremona transformations in plane and space*, Cambridge University Press (1927)
28. Kollár, J., Smith, K., Corti, A.: Rational and nearly rational varieties. *Cambridge Studies in Advanced Mathematics*, **92** (2004)
29. Kumar, M., Murthy, P.: Curves with negative self-intersection on rational surfaces. *J. Math. Kyoto Univ.* **22**(4), 767–777 (1982/83)
30. Küpper, C.: Ueber das Vorkommen von linearen Scharen $g_n^{(2)}$ auf Curven n^{ter} Ordnung C_p^n , deren Geschlecht p grösser ist als p_1 , das Maximalgeschlecht einer Raumcurve $R_{p_1}^n$, *Prag. Ber.* 264–272 (1892)
31. McMullen, C.: Dynamics on blowups of the projective plane. *Publ. Math. Inst. Hautes. Études Sci.* **105**, 49–89 (2007)
32. Nagata, M.: On rational surfaces. I. Irreducible curves of arithmetic genus 0 or 1. *Mem. Coll. Sci. Univ. Kyoto Ser. A Math.* **32**, 351–370 (1960)
33. Pan, I.: Sur le degré dynamique des transformations de Cremona du plan qui stabilisent une courbe irrationnelle non-elliptique. *C.R. Acad. Sci. Paris, Ser. I* **341**(7), 439–443 (2005)
34. Pan, I.: Sur le sous-groupe de décomposition d’une courbe irrationnelle dans le groupe de Cremona du plan. *Michigan Math. J.* **55**(2), 285–298 (2007)
35. Russakovskii, A., Shiffman, B.: Value distribution for sequences of rational mappings and complex dynamics. *Ind. Univ. Math. J.* **46**, 897–932 (1997)
36. Semple, J.G., Roth, L.: *Introduction to algebraic geometry*. Clarendon Press, Oxford (1949)
37. Snyder, V., Coble, A., Emch, A., Lefschetz, S., Sharp F., Sizam, C.: Selected topics in algebraic geometry, Report of the committee on rational transformations, Nat. Research Council, Chelsea Pub. Company (1970)
38. Wiman, A.: Zur Theorie der endlichen Gruppen von birationalen Transformationen in der Ebene. *Math. Ann.* **48**, 195–240 (1896)